



**YTU FEN EDEBİYAT
FAKÜLTESİ**



**YTU FACULTY of ARTS
and SCIENCES**



TURAN25 BOOK OF ABSTRACT BİLDİRİ ÖZET KİTAPÇIĞI



JUNE 23-24-25
2025



İSTANBUL/TÜRKİYE

www.turansymposium.yildiz.edu.tr



TURAN-Fundamental Sciences Symposium

TURAN-Temel Bilimler Sempozyumu

(TURAN25)

Book of Abstract
Bildiri Özet Kitapçığı

June 23-24-25, 2025

İstanbul/Türkiye

ÖNSÖZ

Temel bilimlerin, bilimin merkezi olduğu; bilimsel keşiflerden teknolojik yeniliklere, mühendislik uygulamalarından endüstriyel çözümlere kadar hayata değer katan her gelişmenin dayanağını oluşturduğu ve insanlığın yarınlarnı şekillendirmede kilit rol oynadığı tartışmasız bir gerçektir. Günümüzde, söz konusu gerçeğin toplumsal ve akademik düzeyde yeniden benimsenmesi; bilimsel ilerleme kavramının doğru biçimde tanımlanması ve teknolojik yeniliklerin tek başına bilimsel gelişme sayılmasının yanlış olduğunun altının çizilmesi büyük önem taşımaktadır. Zira bugün ortaya çıkan her teknolojik yenilik, aslında geçmişte yapılmış kapsamlı temel bilim araştırmalarının ve bu alanda biriken bilginin sonucudur. Temel bilimler araştırmalarının ve eğitiminin bu hayati önemi göz ardı edildiğinde, geleceğin bilimini üretme ve geleceğin teknolojilerini geliştirme potansiyeli daha bugünden yok olmaya başlayacaktır. Bu noktada temel bilimlere hak ettiği saygınlığın yeniden kazandırılması amacıyla yürütülecek girişimlerde, tüm paydaşların sorumluluk üstlenmesi büyük bir gerekliliktir.

Temel bilimlere öncelik verilmesinin önemini her fırsatta dile getiren, 2015 Nobel Kimya Ödülü sahibi Prof. Dr. Aziz Sancar'ın Onursal Başkanlığında, Yıldız Teknik Üniversitesi Fen Edebiyat Fakültesi bünyesinde düzenlenen Uluslararası TURAN Temel Bilimler Sempozyumu (TURAN25), söz konusu gereklilik çerçevesinde şu amaçlarla gerçekleştirilmiştir: temel bilimleri gerek ülkemizde gerek Türk dünyasının genelinde güçlendirme ve gelecekteki bilimsel ve teknolojik gelişmelere öncülük ve liderlik etme vizyonu doğrultusunda, temel bilimlerin önemine yönelik farkındalığı artırmak üzere Fakültemizce ulusal ölçekte yürütülen faaliyetleri Türk Devletleri ile başlayarak uluslararası ölçekte genişletmek; Türk Devletleri arasında temel bilimler alanında gerçekleştirilebilecek ikili anlaşmalara, projelere, araştırma ve eğitim faaliyetlerine zemin hazırlamak ve böylece ülkemiz ile diğer Türk Devletleri arasında siyasi, kültürel, ekonomik ve askeri düzeydeki işbirliklerinin yanı sıra yürütülmekte olan bilimsel iş birliklerine somut katkı sunmak ve Türk Dünyası ile birlikte tüm insanlık için bilim yoluyla fayda ve değer üretmek hedefiyle çıkılacak tarihi bir yolculuğun ilk adımını hep birlikte atmak.

Türk dünyasında ortak değerler etrafında yükselen güçlü bir akademik dayanışmayı temsil eden TURAN25, aralarında Almanya, Azerbaycan, Cezayir, Çin, Kazakistan, Kırgızistan, Macaristan, Özbekistan, Rusya ve Türkiye'nin bulunduğu 10 ülkedeki 50 farklı üniversiteden biyoloji, fizik, kimya ve matematik alanlarında çalışan bilim insanlarını buluşturmuş; Sempozyum kapsamında 6 çağrılı konuşma, 71 sözlü bildiri ve 47 poster sunumu gerçekleşmiştir. Sempozyum'da ayrıca Kapanış

Oturumu kapsamında “İkili İşbirlikleri Paneli” gerçekleştirilmiş; temel bilimlerde uluslararası bilimsel iş birliğini güçlendirmenin, kurumsallaştırmanın ve sürdürülebilir kılmanın yolları tartışılmış ve aşağıda sunulan Sonuç Bildirisi hazırlanmıştır.

TURAN25, Türkiye Yüzyılı vizyonumuz çerçevesinde, temel bilimlerin bilimsel gelişme bağlamındaki önemine dair ulusal düzeyde güçlü bir farkındalık oluşturmak ve temel bilimler eğitiminin, günün ve geleceğin gereksinimleri doğrultusunda yeniden yapılandırılmasına öncülük etmek amacıyla Yıldız Teknik Üniversitesi Fen-Edebiyat Fakültesi bünyesinde gerçekleştirdiğimiz kapsamlı çalışmaların bir parçasıdır. Söz konusu çalışmalar çerçevesinde, yeni nesil bir temel bilimler eğitimi modeli ortaya koymak ve böylece temel bilimler öğrencilerine uluslararası standartlarda bir eğitim sunmak amacıyla “YTÜ-FEF Temel Bilimler Eğitimi Modeli” hayata geçirilmiştir. Temel bilimler bölümlerinde öğrenim gören öğrencilerin yalnızca nitelikli birer bilim insanı ve araştırmacı değil, aynı zamanda özel sektörde yüksek istihdam edilebilirliğe sahip Ar-Ge uzmanları olarak yetişmelerini hedefleyen bu Model, öğretim planlarının güncellenmesini; öğretim planlarının tamamına, Yapay Zekâya Giriş, Temel Bilgisayar Teknolojilerine Giriş, Kariyer Planlama ve Mesleki Etik ile Sosyal Sorumluluk ve Adalet derslerinin Zorunlu Fakülte Dersleri olarak eklenmesini; ayrıca temel bilimler öğrencilerinin disiplin-içi ve disiplinlerarası bilimsel uzmanlıklarını derinleştirmelerine imkân tanıyan yenilikçi sertifika programlarının oluşturulmasını ve farkındalık çalışmalarının gerçekleştirilmesini kapsamıştır.

Temel bilimlerin toplum nezdinde değerini görünür kılmak ve temel bilimlerin gerçekte ne olduğuna dair kavrayışı güçlendirmek üzere yürütülen farkındalık çalışmaları bağlamında, lise öğrencileri için YTÜ Fen-Edebiyat Fakültesi Temel Bilimler Farkındalık Çalıştayları düzenlenmiş; bu çalıştaylarla lise öğrencilerinin ve öğretmenlerinin temel bilimlerin anlamını ve değerini doğru kavramaları, gençlerin üniversite ortamında temel bilimleri yakından tanıyarak bilinçli tercihler yapmaları amaçlanmıştır. Lisans düzeyinde yürütülen farkındalık çalışmaları ise temel bilimler öğrencilerinin daha eğitimleri sırasında kendilerini bilim insanı olarak tanımlamaya başlamalarını ve bu kimliği içselleştirmelerini amaçlamıştır. Öğrencileri bilimsel araştırmaya teşvik eden bu faaliyetler arasında Lisans Bitirme Tezi Poster Sunumları, akademisyenlerin bilimsel araştırmalarını öğrencilerle paylaşmasına imkân tanıyan Fakülte Çarşamba Seminerleri ve farklı üniversitelerden lisans öğrencilerinin TÜBİTAK 2209-A/B projelerini sunduğu 2209A/B Fen ve Matematik Öğrenci Çalıştayları yer almıştır. Bu sayede öğrencilerin, ulusal ve uluslararası temel bilimler topluluğunun bir parçası olduklarını hissetmeleri ve alanlarına aidiyet geliştirmeleri hedeflenmiştir.

TURAN25, temel bilimlere ve temel bilimler eğitime dair Fakültemizce ulusal ölçekte yürütölmekte olan söz konusu çalışmaların, Türk dünyasından başlayarak uluslararası ölçekte genişletilmesi yönünde atılmış güçlü bir adım olmuştur. Bilimsel iş birliklerinin sürdürülebilir şekilde güçlendirilmesi fikri gerek katılımcılarımız gerekse toplumun çeşitli kesimlerince büyük bir heyecanla karşılanmıştır. Bu bağlamda yapılan değerlendirmeler sonucunda, Sempozyumun geleneksel hale getirilmesi, ikincisinin 2027 yılında yine Fakültemiz bünyesinde düzenlenmesi, sonraki yıllarda farklı Türk devletlerinin ev sahipliğinde devam ettirilmesi, Türk dünyasında temel bilimlere ilişkin farkındalığı artırabilmek üzere Yıldız TURAN Akademisi kurulması ve temel bilimler alanına yüksek düzeyde bilimsel katkı yapan bilim insanlarını onurlandırmak için Nobel Ödölüne benzer bir “TURAN Temel Bilimler Ödölü” verme sürecinin başlatılmasına karar verilmiştir.

TURAN25’in, Türk devletleri arasında kalıcı ve güçlü bilimsel iş birliklerine ve Türk dünyasının gelecekte temel bilimler araştırmaları ve eğitime yön verecek potansiyelini gerçekleştirmesine vesile olmasını diliyor, temel bilimlere gönöl vermiş tüm bilim insanlarını şimdiden TURAN27’ye davet etmekten büyük bir onur ve mutluluk duyuyorum.

Saygılarımla,



Prof. Dr. Salim Yüce

Sempozyum Başkanı

Yıldız Teknik Üniversitesi Fen Edebiyat Fakültesi Dekanı

10 Temmuz 2025

TABLE OF CONTENTS

HONORARY CHAIR OF THE SYMPOSIUM	10
CHAIR OF THE SYMPOSIUM	10
SCIENTIFIC COMMITTEE	10
ORGANIZING COMMITTEE (FACULTY)	11
ORGANIZING COMMITTEE (MATHEMATICS)	12
ORGANIZING COMMITTEE (BIOLOGY)	13
ORGANIZING COMMITTEE (CHEMISTRY)	14
ORGANIZING COMMITTEE (PHYSICS)	15
INVITED PLENARY SPEAKERS	17
INVITED SPEAKERS	17
ULUSLARARASI TURAN TEMEL BİLİMLER SEMPOZYUMU SONUÇ BİLDİRİSİ	18
INVITED PLENARY SPEAKERS ABSTRACTS	22
Can We Target the Core Clock Mechanism for the Treatment of Different Diseases?	23
Some Questions of X -Valued Elliptic Equations	24
CERN Today: Pushing the Boundaries of Particle Physics	25
INVITED SPEAKERS ABSTRACTS	26
Maveraünnehir, Horasan, Harezmi Havzasında Muhtelif İlimlerin Birlikte Gelişmesi, Meşhur Alimlerin Yetiştirilmesi, Kıymetli Eserlerin Yazılması Üzerine Bir Analiz	27
A Decade of Machine Learning Research: Evaluation Metrics, Taxonomies and Bibliometrics Analysis	29
On the L_1 Data Dirichlet Problem for the Nonuniform Parabolic Equations of Second Order	30
ORAL PRESENTATION ABSTRACTS	31
MATHEMATICS	32
2-Parameter Generalized Quaternions with Generalized Tribonacci Numbers Components	33
Calculus on Real-variable Split Quaternion-valued Functions	34

Characteristic Roots, Eigenvalues and Eigenvectors of a Certain Type of 4×4 Dual Tridiagonal Matrices	35
Comparative Analysis of Türkiye's Road Traffic Accidents Using Data Mining Techniques	36
The Moment of Inertia of Generated Curves: A Holditch-Type Theorem	37
Do Oil Prices Have a Chaotic Effect on Unemployment and Inflation	38
Genel DSW Sistemine Genişletilmiş Jacobi Eliptik Fonksiyon Yaklaşımı: Tam Çözümler, Kararlılık ve Simülasyonlar	39
Bazı Antiviral Moleküllerin Topolojik Karakterizasyonu	40
Affine Factorable Surfaces in Euclidean 4-Space E^4	41
Monge Surfaces with Their Applications	42
Bazı Özel Afin Yapıların ve Duallerinin Kromatik İndislerinin Bulunması Üzerine	43
Optical Bullets for the (3+1)-Dimensional Perturbed Nonlinear Schrödinger Equation having the Cubic-Quintic Laws and the Spatio-Temporal Dispersion	44
TBMM Maarif Vekâletinin 1920 Yılında Aldığı Geometri Öğretimi Kararları Üzerine	45
Competitive Systems via Hamiltonian Framework and Artificial Neural Estimation	46
Data-Driven Estimation of Krylov Iterations in Riccati Solvers	47
q-Türev Kullanılarak Tanımlanan Yeni Bir Harmonik Fonksiyon Sınıfının Geometrik Özellikleri	48
Some Characterizations of Gradient Ricci-Yamabe Solitons	49
Matematik Lisans Öğrencilerinin Seçmeli Ders Seçimlerinin Oyun Teorisi ile Matematiksel Modellenmesi	50
Improved Fixed Point Approximations via Picard's Three-Step Iteration for Suzuki-Type Non-Expansive Mappings	51
Investigation of Fractal Dimension Algorithms in Different Geometries	52
Machine Learning-Based Classification of Chaotic Behavior in Discrete Dynamical Systems	53
Variation of Parameters and Hölder Stability of Differential Systems with Initial Time Difference	54
W5-Curvature Tensor on a Complex Contact Space Form	55
Schauder Estimates for Elliptic Operators in Banach-Sobolev Spaces	56
Identities on the Generalized Harmonic Numbers of Rank with Order Using Generating Functions	57
Solitary Wave Solutions of the Two-Dimensional Fractional-Order Nonlinear Schrödinger Equation	58
Boundedness Criteria for the G -fractional Integral and G -fractional Maximal Operator on Gegenbauer-morrey Spaces	59
Hemen Hemen Neo Kobalans Sayıları	60
Δ -Convergence For Proximal Point Algorithm And Fixed Point Problem in Generalized $CAT(0)$ Space	61
Impulse-Hydrodynamic Model of Explosive Impact on a Half-Plane with Circular Cavity	62
Model and Theorem of Existence of the Problem of Optimal Control of the Customs Clearance Process	63
Necessary Conditions for Weak and Strong Local Minimum the Problem of the Calculus of Variations with Higher Order Derivatives	64
Numerical Analysis of Stiffened Shell Structures Using an Efficient Three-Dimensional Finite Element	65
Development of a General Algorithm for Solving Stability Problems of Plates on Elastic Bases	66
Modelling Electromagnetic Wave Propagation Through Time-Varying Medium Using the Finite Difference Time Domain Method	67
Numerical Modeling of Stars and Mathematical Analysis	68
New Midpoint and Trapezoid-Type Inequalities for Co-ordinated m -Convex Functions via Variable-Order Fractional Calculus	69
A Study on Dual Quaternions with k -Pell and k -Pell-Lucas Coefficients	70
A Study on Vietoris' Hybrinomials	71
On Multiplicative Magnetic Flux Ruled Surfaces	72
A Study on the Dual Quaternionic Sequence with Vietoris' Components	73
On an Algorithm for a Morphological Analyzer of the Azerbaijani Language	74

The Influence of the Interaction with Fluids of the Hollow Cylinder with Inhomogeneous Initial Stresses on the ZGV Modes that Occur with Axisymmetric Wave Propagation	75
On the Strong Solvability of a Nonlocal Boundary Value Problem for the Laplace Equation in Weighted Grand Sobolev Spaces in Rectangle	76
Hyperbolic Functions on Split Quaternions and Their Characteristics	77
Surface Pencil with Constant Gaussian Curvature Along a Non-null Curve	78
Rectifying Rose Ruled Surfaces	79
The Cross Products on Dual and Dual Lorentzian Spaces	80
PHYSICS	81
Dielectric Relaxation and Conductivity Characteristics of Gamma-Irradiated TlGaTe ₂ and TlInSe ₂ Crystals	82
Electrodeposition and Spray Techniques for Pt/Pd Coated Nickel Foam Electrodes	83
Körük Ventili (Levelling Valve) Hava Akış Karakteristiklerinin Karşılaştırmalı Teknik İncelemesi	84
Sinop İlinde Nükleer Güç Santrali Öncesi Kabalı Nehrinin Sularında Trityum Radyoaktivitesinin Araştırılması	85
Surface Engineering of Ti13Nb13Zr Implants Using Microarc Oxidation and Hydroxyapatite-Based Thermal Spraying	86
A New Force in Nature? The Magnetic Moment of the Muon	87
Dielectric Properties Of CuTiS ₂ Crystals Studied Via Impedance Spectroscopy	88
Effect of Doping and Radiation on the Conductivity Mechanism in Layered GaS Crystals	89
Prospects of Thermal Waters Usage in Small Power Industry	90
(E)-2-(2,4-Dihidroksibenziliden)Tiyosemikarbazon ve (E)-2-[(1H-İndol-3 İl)Metilen]Tiyosemikarbazon Komplekslerinin Yapısal ve Spektroskopik Özelliklerinin DFT Metoduyla İncelenmesi	91
Optimization of Ion-Plasma Nitriding Parameters for Ti-6Al-4V Titanium Alloy	92
Enhancement of Visible-Light-Driven Photocatalytic Activity in Gd-Doped CdZnS Nanocomposites	93
CHEMISTRY	94
Particle Size Analysis of Polymer-Based Metal Nanoparticles Using DLS Method	95
The Study of The Swelling Ratio of Gum Arabic and Acrylamide Hydrogel in Solutions with Different pH	96
Synthesis, Structure, and Properties of New Alicyclic and Aromatic Amidophosphates	97
Development of Lead-Free Epoxy Resin Composites for X-Ray Shielding	98
Alunit Cevherinin RSM Kullanılarak Liçinin Optimizasyonu	99
Synthesis of Green Chemistry-Based Catalyst for Carbon Monoxide Neutralization	100
BIOLOGY	101
Potential Use of Watermelon Seed Oil as an Alternative Linoleic Acid Source Enhanced by Coffee-Derived Caffeic Acid in Cholesterol Treatment	102
Bazı Yonca (<i>Medicago sativa</i> L.) Genotiplerinin iPBS-retrotranspozon Markörleri ile Genetik Çeşitliliğin Değerlendirmesi	103
γ (Gama) Işınlarıyla Muamelenin Bitkilerde Tuz (NaCl) Stresine Tepkisi	104
POSTER PRESENTATION ABSTRACTS	105
MATHEMATICS	106
Ağırlıklı Standart Olmayan Banach Uzaylarında Harmonik Fonksiyonların Yapısal Analizi	107
Structural Properties of the $h_p(X)$ Class of X-Valued Harmonic Functions	108
Exact Solutions of the Time fractional Benney-Luke Equation and the Second Order Benjamin-Ono Equation	109
Lipschitz Stability Analysis of Nonlinear Set-Valued Differential Equations with Initial Time Differences	110
Orlicz-Sobolev Uzaylarında Yerel Olmayan Bir Problemin Güçlü Çözülebilirliği	111

Ülke İçinde Karayolu ile Yük Taşımacılığı	112
Fe-Si-C Amorf Şerit Numunelerinin Araştırılması ve Kristal Fe-Si Muadilleri ile Karşılaştırılması	113
Basicity of a Trigonometric System in Morrey-type Spaces	114
PHYSICS	115
Electron Screening Meets Space: D+D Fusion-Driven Propulsion Systems	116
Constraining Key Resonances in the $^{40}\text{Ca}(p,\gamma)^{41}\text{Sc}$ Reaction via ANC-Assisted DWBA Analysis of Transfer Reactions and its Astrophysical Reaction Rate	117
Contaminated or Clean? How Do Surface Conditions Shape Fusion Rates in Deuterated and Boron Targets?	118
Fusion in the Deep: Pycnonuclear Reactions Under Extreme Stellar Conditions	119
Atmospheric Physics and Natural Transport Paths in the Atmosphere	120
Renormalization Group Studies of Continuous-Spin Complex Systems	121
A Holistic Design Approach to Compact Neutron Generators: Adjustable Energies & Enhanced Yield with D-D Fusion Reactions	122
Re-thinking the $^8\text{Li}(\alpha, n)^{11}\text{B}$ Reaction: Evidence for a Missing Resonance in ^{12}B	123
Aktif Matrisli Hibrit Kompozitlerin Piezoelektrik Özellikleri	124
Amorf Şerit Üretiminde Kalite Kontrol İçin Vickers Sertlik Ölçümlerinde Standart Sapma Potansiyeli	125
Dielectric Performance and Potential Applications of HDPE/nano MgO Composites	126
Electrical Conductivity of Low-Dimensional Electron Gas in Asymmetric Quantum Well Scattering in Ionized Impurities in the Absence of Screening	127
Influence of Gamma Radiation on the Anisotropy of Electrical Conductivity in TlGaTe_2 Crystals	128
Role of the P_z^{1+} State in Ferromagnetism and Topology of MnBi_2Te_4	129
Yapay Zeka Destekli Moleküler Sistemlerin Kuantum Simülasyonu: H_2 Molekülü İçin Qiskit ve OpenFermion ile VQE Uygulaması ve Hata Analizi	130
CHEMISTRY	131
Influence of Water Quality on The Electrochemical Behavior of a Carbon Steel Used in The Combustion Chamber of Boilers in a Petroleum Complex	132
Inhibitive Effect of an Organic Compound on the Corrosion of Carbon Steel in Demineralized Water	133
Synthesis and Photopolymerization of Imidazole-based Photoinitiator	134
Comparative Study of the Performance of Activated Carbon Obtained By Activation and Carbonization of Adsorbent Material Obtained From Avocado Peel with Different Chemicals in Removing Textile Dyes	135
Treatment of Synthetic Textile Wastewater Containing Both Basic Yellow 28 and Basic Blue 3 Dyes by Electrochemical Method	136
The Synthesis And Mesomorphic Investigation of New Chiral Calamitic Liquid Crystal Derived From Cyanobiphenyl Core	137
One-Step Preparation of Composite Phase Change Films via Photoinitiated Oil-in-Oil Emulsion Templating	138
The Preparation and Mesomorphic Investigation of Binary Mixture of a New Chiral Rod-Like Molecule and Benzoic acid Derivative	139
Influence of Monomer Composition on the Morphological and Mechanical Properties of Terpene Derivative Macroporous Polymers	140
Preparation and Characterization of Alumina Enhanced Kaolinite Supported Capric Acid Based Composite Phase Change Materials	141
Synthesis of TX-Based Photoinitiator with Antibacterial Activity	142
Assessment of the Impact of Climate Change Projects on the Sustainable Development of the Country Based on Multi-Criterion Analysis Method	143
Improving Flame Resistance and Polymer Compatibility of Ethylene Propylene-Diene Rubber (SKEPT-60) Through Polyvinyl (PVX) Modification	144

Synthesis of Reagents Based On Natural Oil Acids and Ethylenediamine and Their Study As Components For Preservation Fluids	145
Chitosan-Tannin Based Flocculant for Heavy Metal (Ni^{2+} , Cu^{2+} , and Zn^{2+}) Removal from Industrial Wastewater: Investigating the Influence of pH, Reaction Time, and Stirring Speed	146
Intelligent Identification of Multicomponent Mixture Components Using an Artificial Intelligence	147
Investigation of Concentration Liquids Formulated by Incorporating Amidoamine and Various Fatty Acids into T-30 Oil Distillate	148
Nitration Behavior of Aromatic- and Olefin-Enriched Coking Light Fraction with Ionic Catalytic Systems	149
Phosphoryl-Substituted α -Chloro- and α -Thiocyanatoacetaldehydes in the Reaction with α -Pyridone	150
BIOLOGY	151
Analysis of Codon Usage Bias of Chloroplast Genomes in <i>Camellia</i> Species	152
TURAN-Fundamental Sciences Symposium (TURAN25) Delivered Oral Presentations	153
TURAN-Fundamental Sciences Symposium (TURAN25) Delivered Poster Presentations	157
MAIN SPONSOR	159
SPONSORS	159

HONORARY CHAIR OF THE SYMPOSIUM

Prof. Dr. Aziz SANCAR, 2015 Nobel Laureate in Chemistry, Department of Biochemistry and Biophysics, University of North Carolina, USA

CHAIR OF THE SYMPOSIUM

Prof. Dr. Salim YÜCE, Dean of the Faculty of Arts and Sciences, Yıldız Technical University, Türkiye

SCIENTIFIC COMMITTEE

Prof. Dr. Salim YÜCE, Yıldız Technical University, Türkiye

Prof. Dr. Misir MARDANOV, Institute of Mathematics and Mechanics of ANAS, Azerbaijan

Prof. Dr. Ahmet VELİYEV, Odlar Yurdu University, Azerbaijan

Prof. Dr. Alma DAULETBKOVA, L. N. Gumilyov Eurasian National University, Kazakhstan

Prof. Dr. Avyt ASANOV, Türkiye Manas University, Kyrgyzstan

Prof. Dr. Bilal BİLALOV, Yıldız Technical University, Türkiye

Prof. Dr. Emin KASAP, Ondokuz Mayıs University, Türkiye

Prof. Dr. Fakhreddin MAMMADOV, Odlar Yurdu University, Azerbaijan

Prof. Dr. Fariz EMİROV, Azerbaijan Technical University, Azerbaijan

Prof. Dr. H. Hilmi HACISALİHOĞLU, Türkiye

Prof. Dr. Kadri ARSLAN, Bursa Uludağ University, Türkiye

Prof. Dr. Mehmet Hakkı OMURTAG, Medipol University, Türkiye

Prof. Dr. Murat TOSUN, Sakarya University, Türkiye

Prof. Dr. Nigar ASLANOVA, Azerbaijan University of Architecture and Construction, Azerbaijan

Prof. Dr. Nodirbek YUSUPBEKOV, Tashkent State Technical University, Uzbekistan

Prof. Dr. Nuri KURUOĞLU, Istanbul Gelişim University, Türkiye

Prof. Dr. Surkay D. AKBAROV, Yıldız Technical University, Türkiye

Prof. Dr. Telman GASIMOV, Baku State University, Azerbaijan

ORGANIZING COMMITTEE (FACULTY)

Prof. Dr. Muhammet Kasım ŞENER, Yıldız Technical University, Türkiye

Assoc. Prof. Dr. Yonca SEZER, Yıldız Technical University, Türkiye

Prof. Dr. Hale OCAK GÜMRÜKÇÜ, Yıldız Technical University, Türkiye

Assoc. Prof. Dr. Senem ÖNER BULUT, Yıldız Technical University, Türkiye

ORGANIZING COMMITTEE (MATHEMATICS)

Prof. Dr. Mustafa DÜLDÜL (Head of Department), Yıldız Technical University, Türkiye

Assoc. Prof. Dr. Gülsüm Yeliz SAÇLI (Department Representative), Yıldız Technical University, Türkiye

Prof. Dr. Salih ÇELİK, Yıldız Technical University, Türkiye

Assoc. Prof. Dr. Fatma KARACA, Yıldız Technical University, Türkiye

Assoc. Prof. Dr. Neslihan ÖZDEMİR, Yıldız Technical University, Türkiye

Assoc. Prof. Dr. Nurten GÜRSES, Yıldız Technical University, Türkiye

Assoc. Prof. Dr. Seda KIZILBUDAK ÇALIŞKAN, Yıldız Technical University, Türkiye

Assist. Prof. Dr. Eda YILDIZ YILMAZ, Yıldız Technical University, Türkiye

Assist. Prof. Dr. Elif DEMİR, Yıldız Technical University, Türkiye

Assist. Prof. Dr. Sema AKIN BAŞ, Yıldız Technical University, Türkiye

Assist. Prof. Dr. Faruk DEVELİ, Yıldız Technical University, Türkiye

Research Assistant Dr. Fatma ÇELİKER, Yıldız Technical University, Türkiye

Research Assistant Dr. Hande USLU TUNA, Yıldız Technical University, Türkiye

Research Assistant Hamide Feyza AYKUT, Yıldız Technical University, Türkiye

Research Assistant Çiğdem Zeynep YILMAZ, Yıldız Technical University, Türkiye

Research Assistant Sevilay ÇORUH ŞENOCAK, Ondokuz Mayıs University, Türkiye

Lecturer Kübra ÇETİNERK İstanbul Gelişim University, Türkiye

Lecturer İpek Ebru KARAÇAY, İstanbul Gelişim University, Türkiye

Research Assistant Duygu ÇAĞLAR ÇAY, İstanbul Gelişim University, Türkiye

Research Assistant Edanur ERGÜL ARSLAN, Marmara University, Türkiye

Research Assistant Ümit ILDIZ Marmara University, Türkiye

Research Assistant Pelin DURSUN, İstanbul Ticaret University, Türkiye

Ph. D. Student Mahmure AVEY, Yıldız Technical University, Türkiye

Ph. D. Student Damla BUDAK, Yıldız Technical University, Türkiye

Master Student Zehra Nur BAKIRBAŞ, Yıldız Technical University, Türkiye

ORGANIZING COMMITTEE (BIOLOGY)

Prof. Dr. Semiha ERİŞEN (Head of Department), Yıldız Technical University, Türkiye

Prof. Dr. Sevgi MARAKLI (Department Representative), Yıldız Technical University, Türkiye

Assist. Prof. Dr. Burcu GÜNDÜZ ERGÜN, Yıldız Technical University, Türkiye

Research Assistant Dr. Mine KUÇAK, Yıldız Technical University, Türkiye

Research Assistant Hasan Hüseyin GÜLERCAN, Yıldız Technical University, Türkiye

Research Assistant Şener ÇİNTESUN, Yıldız Technical University, Türkiye

ORGANIZING COMMITTEE (CHEMISTRY)

Prof. Dr. Ali ERDOĞMUŞ (Head of Department), Yıldız Technical University, Türkiye

Prof. Dr. Özlem YAZICI (Department Representative), Yıldız Technical University, Türkiye

Prof. Dr. Dolunay ŞAKAR DAŞDAN, Yıldız Technical University, Türkiye

Prof. Dr. Fatih ÇAKAR, Yıldız Technical University, Türkiye

Prof. Dr. Melda ALTİKATOĞLU YAPAÖZ, Yıldız Technical University, Türkiye

Assoc. Prof. Dr. Bahadır KESKİN, Yıldız Technical University, Türkiye

Assoc. Prof. Dr. Birol IŞIK, Yıldız Technical University, Türkiye

Assoc. Prof. Dr. Göknur YAŞA ATMACA, Yıldız Technical University, Türkiye

Assoc. Prof. Dr. Sevnur KESKİN DOĞRUYOL, Yıldız Technical University, Türkiye

Assoc. Prof. Dr. Türkan BÖRKLÜ BUDAK, Yıldız Technical University, Türkiye

Assist. Prof. Dr. Ceren CAN KARANLIK, Yıldız Technical University, Türkiye

Assist. Prof. Dr. Çiğdem BİLEN, Yıldız Technical University, Türkiye

Assist. Prof. Dr. Sezin ERARPAT BODUR, Yıldız Technical University, Türkiye

Assist. Prof. Dr. Şeyda AYDOĞDU, Yıldız Technical University, Türkiye

Assist. Prof. Dr. Volkan UĞRAŞKAN, Yıldız Technical University, Türkiye

Research Assistant Dr. Büşra ERTÜRK, Yıldız Technical University, Türkiye

Research Assistant Dr. Erdoğan KİRPİ, Yıldız Technical University, Türkiye

Research Assistant Dr. Erem AHMETALİ, Yıldız Technical University, Türkiye

Research Assistant Dr. Fatma Tülay TUĞCU, Yıldız Technical University, Türkiye

Research Assistant Dr. İbrahim E. ÖZYİĞİT, Yıldız Technical University, Türkiye

Lecturer Dr. Ayşe ERDOĞAN ÇAKAR, Yıldız Technical University, Türkiye

Research Assistant Ayşe Enda ZAVALSIZ, Yıldız Technical University, Türkiye

Research Assistant Deniz KILIÇ ÖZTÜRK, Yıldız Technical University, Türkiye

Research Assistant Elif ÖZÇELİK KAZANCIOĞLU, Yıldız Technical University, Türkiye

Research Assistant Mesut BİLGİ, Yıldız Technical University, Türkiye

Research Assistant Sadegül FIRAT, Yıldız Technical University, Türkiye

Research Assistant Sena Nur ALEGÖZ, Yıldız Technical University, Türkiye

Research Assistant Süreyya DEDE, Yıldız Technical University, Türkiye

Research Assistant Ümmügülsüm POLAT KORKUNÇ, Yıldız Technical University, Türkiye

ORGANIZING COMMITTEE (PHYSICS)

Prof.Dr. Zeynep GÜVEN ÖZDEMİR (Head of Department), Yıldız Technical University, Türkiye

Assoc. Prof. Dr. Banu SÜNGÜ MISIRLIOĞLU (Department Representative), Yıldız Technical University, Türkiye

Prof.Dr. Murat ODUNCUOĞLU, Yıldız Technical University, Türkiye

Prof.Dr. Reyhan KAYA, Yıldız Technical University, Türkiye

Assoc. Prof. Dr. Arzu ÇİLLİ, Yıldız Technical University, Türkiye

Assoc. Prof. Dr. Ayşe DURUSOY, Yıldız Technical University, Türkiye

Assoc. Prof. Dr. Birsal CAN ÖMÜR, Yıldız Technical University, Türkiye

Assoc. Prof. Dr. Pınar GÖKDEMİR CHOİ, Yıldız Technical University, Türkiye

Assist. Prof. Dr. Macide CANTÜRK RODOP, Yıldız Technical University, Türkiye

Assist. Prof. Dr. Altan BOZDOĞAN, Yıldız Technical University, Türkiye

Research Assistant Dr. Ayşegül ÇELİK BOZDOĞAN, Yıldız Technical University, Türkiye

Research Assistant Dr. Kıyami ERDİM, Yıldız Technical University, Türkiye

Research Assistant Dr. Pelin KAVAK, Yıldız Technical University, Türkiye

Research Assistant Dr. Ünsal AkDERE, Yıldız Technical University, Türkiye

Research Assistant Ayşenur ŞAHİN, Yıldız Technical University, Türkiye

Research Assistant Emre BEYAZAY, Yıldız Technical University, Türkiye

Research Assistant İsmail Deniz GÜN, Yıldız Technical University, Türkiye

Research Assistant Mesut KAVAL, Yıldız Technical University, Türkiye

Research Assistant Ozan TOKER, Yıldız Technical University, Türkiye

Research Assistant Vildan YILMAZ, Yıldız Technical University, Türkiye

INVITED PLENARY SPEAKERS

İ. Halil KAVAKLI, Can We Target the Core Clock Mechanism for the Treatment of Different Diseases?, Biology

Bilal BİLALOV, Some Questions of X-valued Elliptic Equations, Mathematics

Salim ÇERÇİ, CERN Today: Pushing the Boundaries of Particle Physics, Physics

INVITED SPEAKERS

Kemal Yavuz ATAMAN, Maveraünnehir, Horasan, Harezmi Havzasında Muhtelif İlimlerin Birlikte Gelişmesi, Meşhur Alimlerin Yetiştirilmesi, Kıymetli Eserlerin Yazılması Üzerine Bir Analiz, Tarih

Amir MOSAVİ, A Decade of Machine Learning Research: Evaluation Metrics, Taxonomies and Bibliometrics Analysis, Artificial Intelligence

Ferman MAMEDOV, On the L_1 data Dirichlet problem for the nonuniform Parabolic Equations of Second Order, Mathematics

ULUSLARARASI TURAN TEMEL BİLİMLER SEMPOZYUMU SONUÇ BİLDİRİSİ

Yıldız Teknik Üniversitesi Fen Edebiyat Fakültesi bünyesinde, 2015 Nobel Kimya Ödülü sahibi Prof. Dr. Aziz Sancar'ın Onursal Başkanlığında 23-25 Haziran 2025 tarihleri arasında düzenlenen Uluslararası TURAN Temel Bilimler Sempozyumu (TURAN25) kapsamında, Türk devletlerinde temel bilimler alanında çalışan bilim insanları İstanbul'da bir araya gelmiştir.

Aralarında Almanya, Azerbaycan, Cezayir, Çin, Kazakistan, Kırgızistan, Macaristan, Özbekistan, Rusya ve Türkiye'nin bulunduğu 10 ülkedeki 50 farklı üniversiteden biyoloji, fizik, kimya ve matematik alanlarında çalışan bilim insanlarının katıldığı Sempozyumun üçüncü gününde (25 Haziran 2025) Kapanış Oturumu kapsamında "İkili İşbirlikleri Paneli" gerçekleştirilmiştir.

Panel'de, Fen Edebiyat Fakültesi Dekanı ve Sempozyum Başkanı Prof. Dr. Salim Yüce, Sempozyumun, Yıldız Teknik Üniversitesi Fen Edebiyat Fakültesi bünyesinde, temel bilimler eğitiminin geleceğini inşa etmeye yönelik yürütülmekte olan çalışmaların çok önemli bir parçası olduğunu ifade etmiştir. Yüce, Sempozyumun, temel bilimlerin farklı alanlarından bilim insanlarının bir araya getirdiğini ve bu anlamda tarihte bir ilk olduğunu vurgulamıştır.

Yüce, Sempozyumun amacının, temel bilimlerin gerek ülkemizde gerek Türk Dünyasının genelinde güçlendirilmesi ve gelecekteki bilimsel ve teknolojik gelişmelere öncülük ve liderlik etme vizyonu doğrultusunda, temel bilimlerin önemine yönelik farkındalığı artırmak üzere Yıldız Teknik Üniversitesi Fen Edebiyat Fakültesi bünyesinde ulusal ölçekte yürütülen faaliyetleri Türk Devletleri ile başlayarak uluslararası ölçekte genişletmek; Türk Devletleri arasında temel bilimler alanında gerçekleştirilebilecek ikili anlaşmalara, projelere, araştırma ve eğitim faaliyetlerine zemin hazırlamak ve böylece ülkemiz ile diğer Türk Devletleri arasında siyasi, kültürel, ekonomik ve askeri düzeydeki işbirliklerinin yanı sıra yürütülmekte olan bilimsel iş birliklerine somut katkı sunmak ve Türk Dünyası ile birlikte tüm insanlık için bilim yoluyla fayda ve değer üretmek hedefiyle çıkılacak tarihi bir yolculuğun ilk adımını hep birlikte atmak olduğunu söylemiştir.

Yapay zekâ çağının, toplumun her kesimi için olduğu kadar üniversiteler için de çeşitli meydan okumaları beraberinde getirmekte olduğunu, üniversitelerin “yeni bilgi üretme” ve “yeni bilgi üretecek mezunlar yetiştirme” potansiyelinin yeni bir bakış açısıyla ele alınmak zorunda olduğunu vurgulayan Yüce, bu bağlamda temel bilimler eğitiminin yeniden düşünülmesine gereksinim olduğunu dile getirmiştir.

Yüce, temel bilimler eğitiminin 21. yüzyılda yeniden yapılandırılmasına yönelik yenilikçi bir çerçeve sunan ve Fen Edebiyat Fakültesi bünyesinde geliştirilip uygulamaya konmuş olan yeni nesil “YTÜ FEF Temel Bilimler Eğitimi Modeli”nin, temel bilimlerin, bilimin merkezi olduğu fikrini yeniden tesis etmeyi, temel bilimler eğitiminin niteliğini sürdürülebilir şekilde artırmayı, öğrencilere temel bilimler alanında uluslararası standartlarda yetkinlik ile birlikte disiplinlerarası uzmanlaşma kazandırmayı ve hepsinin ötesinde, toplumsal ve insani fayda odaklı araştırma ve bilgi üretimini merkeze alan bütüncül bir eğitim yaklaşımını hayata geçirmeyi hedeflediğini ifade etmiştir

Azerbaycan Milli Bilimler Akademisi Üyesi ve Eski Milli Eğitim Bakanı Prof. Dr. Misir Mardanov, Türk devletlerindeki değerli bilim insanlarını bir araya getiren ve farklı temel bilimler alanları arasındaki bilimsel etkileşime ve paylaşıma olanak tanıyan TURAN25 Sempozyumunun tarihi ve çok değerli bir girişim olduğunu ve bu kapsamdaki faaliyetlerin güçlendirilmesi, yaygınlaştırılması ve sürdürülebilir kılınması gerektiğini vurgulamıştır. Prof. Dr. Salim Yüce’nin liderliğinde geliştirilmiş ve uygulamaya konmuş olan “YTÜ FEF Temel Bilimler Eğitimi Modeli”nin, temel bilimler eğitiminde yeni bir istikamet gösterdiğini, gelecekte üniversitelerin değişip dönüşeceğini ve bu bağlamda özellikle Model kapsamında geliştirilen ve uzmanlaşmayı hedefleyen yenilikçi sertifika programlarının kritik önem taşıdığını, temel bilimler öğrencilerine alan bilgisi ile birlikte uygulamaya dönük beceri ve yetkinlikler kazandırmanın geleceğin eğitim faaliyetleri açısından çok önemli olduğunu ifade etmiştir.

Panel’de, TURAN25 Sempozyumunda ilk adımı atılmış olan, Türk devletleri arasındaki bilimsel güç birliğini sürdürülebilir şekilde inşa etmenin ve geliştirmeye yönelik girişimlerin ilerletilmesine yönelik yürütülen tartışmalar ve yapılan değerlendirmeler sonucunda,

1. TURAN25 Temel Bilimler Sempozyumunun her iki yılda bir düzenlenerek geleneksel hale getirilmesine,
2. Sempozyumun ikincisinin çok daha geniş bir katılımı 2027 yılında, yine Yıldız Teknik Üniversitesi Fen Edebiyat Fakültesi bünyesinde ve liderliğinde düzenlenmesine,
3. Tüm Türk devletlerinde TURAN27 kapsamındaki çalışmaları yürütmek üzere bilim komiteleri oluşturulmasına,
4. Oluşturulacak bilim komitelerinin faaliyetlerinin Yıldız Teknik Üniversitesi Fen Edebiyat Fakültesi tarafından koordine edilmesine,
5. TURAN27 Sempozyumuna katılımın artırılmasına, özellikle genç bilim insanları ve doktora öğrencilerinin katılıma güçlü biçimde teşvik edilebilmeleri için önlemler alınmasına ve düzenlenecek Sempozyumda genç bilim insanlarını, alanında deneyimli ve kıdemli bilim insanları ile buluşturacak Atölye Çalışmaları gibi uygulamaların gerçekleştirilmesine,
6. Sempozyuma katılım gösteren ülke temsilcilerinin Sempozyum ile ilgili değerlendirmelerini içeren yazılı bir rapor hazırlamalarına, bu raporu YTÜ FEF Sempozyum Düzenleme Kuruluna sunmalarına ve Sempozyum kapsamında yürütülen çalışmalara ilişkin olarak kendi ülkelerinde en yüksek düzeyde bilgilendirme ve tanıtma faaliyetlerinde bulunmalarına,
7. Temel bilimler alanına yüksek düzeyde bilimsel katkı yapan bilim insanlarını onurlandırmak için bir “TURAN Temel Bilimler Ödülü” verme sürecinin başlatılmasına,
8. Temel bilimlerin Türk dünyasında güçlendirilmesine yönelik çalışmaların öncülüğünü yapan Yıldız Teknik Üniversitesi Fen Edebiyat Fakültesi bünyesinde temel bilimler kapsamında araştırma ve eğitim faaliyetlerine yön verecek bir Yıldız TURAN Akademisi kurulmasına yönelik çalışmalar yapılmasına,

9. Geleceğin temel bilimler eğitimini inşa etmek üzere “YTÜ FEF Temel Bilimler Eğitimi Modeli”nin diğer Türk devletlerindeki üniversitelerde de yaygınlaştırılması yönelik çalışmalar yürütülmesine ve Türk devletleri arasında bu kapsamda akademik değişim programları ve iş birlikleri oluşturulmasına,
10. TURAN Temel Bilimler kapsamındaki tüm bu çalışmaların, faaliyetlerin ve ileriye dönük planların tüm Türk devletlerindeki üst düzey yetkililer ile paylaşılmasına ve bu çalışmaların sürdürülebilirliğini ve en yüksek düzeyde fayda üretmesini sağlamak üzere devlet desteğinin talep edilmesine karar verilmiştir.

25 Haziran 2025



Prof. Dr. Salim YÜCE

Sempozyum Başkanı
Fen Edebiyat Fakültesi Dekanı
Yıldız Teknik Üniversitesi



Prof. Dr. Misir MARDANOV

Azerbaycan Cumhuriyeti Bilim ve Eğitim Bakanlığı
Matematik ve Mekanik Enstitüsü Müdürü

INVITED PLENARY SPEAKERS ABSTRACTS

Can We Target the Core Clock Mechanism for the Treatment of Different Diseases?

İbrahim Halil KAVAKLI

*Koc University, Departments of Molecular Biology and Genetics,
Chemical and Biological Engineering, Rumelifeneri Yolu, Sarıyer, İstanbul, Türkiye*

hkavakli@ku.edu.tr

Circadian rhythm is an intrinsic timing system that enables organisms to anticipate and adapt to daily environmental changes, regulating physiological processes such as the sleep-wake cycle. This rhythm is controlled by a molecular circadian clock mechanism driven by a transcriptional and translational feedback loop (TTFL). In mammals, TTFL is orchestrated by the interaction of four key clock proteins: BMAL1, CLOCK, Cryptochromes (CRY), and Periods (PER). BMAL1 and CLOCK form dimers that bind to E-box elements in promoter regions, initiating the transcription of clock-controlled genes (CCGs). Disruption of circadian rhythm has been linked to various diseases, including diabetes, sleep disorders, and neurological conditions. For a long time, it was unclear whether targeting the core clock mechanism could offer therapeutic benefits. Our research, employing *in silico*, *in vitro*, and *in vivo* methods, has identified several small molecules that modulate core clock components. These molecules have demonstrated potential therapeutic effects in animal models for various diseases. In this talk, I will highlight these recent advancements and their implications for disease treatment.

Some Questions of X -Valued Elliptic Equations

Bilal T. BILALOV ^{1,2,3,4}

¹*Odlar Yurdu University, Baku, Azerbaijan*

²*Department of Algebra and Mathematical Logic, Institute of Mathematics and Mechanics, The Ministry of Science and Education of the Republic of Azerbaijan, Baku, Azerbaijan*

³*Department of Mathematics, Yıldız Technical University, Davutpasa, 34220, Istanbul, Türkiye*

⁴*Azerbaijan University of Architecture and Construction, Baku, Azerbaijan*

b_bilalov@mail.ru, bilal.bilalov@yildiz.edu.tr

We consider X -valued elliptic equations with scalar coefficients. First, we provide a brief overview of results obtained in this direction. Then, some new results concerning the solvability problems of X -valued elliptic equations are presented, when X has the so-called Unconditional Martingale Difference (UMD) property. We demonstrate the main scheme for establishing the Fredholmness (or Noetherness) of the considered equations in X -valued Sobolev spaces. Furthermore, we establish interior Schauder-type estimates for m -th order X -valued elliptic operators and investigate the boundedness properties of integral operators with Poisson kernels, associated with some boundary operators under which the complementary conditions hold. These results play a key role in the investigation of the solvability of Boundary Value Problems (BVPs).

Keywords: X -valued elliptic equations, UMD property, Noetherness, X -valued Sobolev spaces, Schauder-type estimates, Poisson integrals.

References

- [1] Ladyzhenskaya, O. A. , Uraltseva, N. N. (1964). Linear and Quasilinear Equations of Elliptic Type. Nauka Moscow [in Russian].
- [2] Bers L., John F., and Schechter M., *Partial Differential Equations*. Lectures in Applied Mathematics, 3A. Providence, R.I.: AMS XIII, 343, 1979.
- [3] Bilalov, B. T., Sadigova, S. R. (2021). *Interior Schauder-type Estimates for Higher-order Elliptic Operators in Grand-Sobolev Spaces*. Sahand Communications in Mathematical Analysis, 18(2), 129-148.
- [4] Bilalov, B. T., Sadigova, S. R. (2024). *On the Fredholmness of the Dirichlet Problem for a Second-order Elliptic Equation in Grand-Sobolev Spaces*. Ricerche Di Matematica, 73, 283-322.

CERN Today: Pushing the Boundaries of Particle Physics

Salim ÇERÇİ

Yildiz Technical University, Faculty of Arts and Sciences, Department of Physics, Istanbul, Türkiye

salim.cerci@yildiz.edu.tr

CERN, the world's largest particle physics laboratory, continues to lead the exploration of fundamental questions about the universe. From high-energy collisions at the Large Hadron Collider (LHC) to the development of next-generation detectors and computing systems, CERN's current research pushes the limits of both technology and theory. This talk will provide an overview of recent developments at the CMS¹ experiment at CERN, including the latest results from LHC Run 3, major detector upgrade efforts such as the High-Granularity Calorimeter (HGCal), and ongoing searches for new physics beyond the Standard Model.² Special emphasis will be placed on the role of international collaboration, technological innovation, and precision measurements in shaping the future of particle physics.

Keywords: Particle physics, CERN, CMS experiment.

References

- [1] CMS Collaboration. (2008). *The CMS experiment at the CERN LHC*, JINST 3 S08004. doi: 10.1088/1748-0221/3/08/S08004.
- [2] CMS Collaboration. (2017). *The Phase-2 Upgrade of the CMS Endcap Calorimeter*, CERN-LHCC-2017-023, CMS-TDR-019, Technical Design Report, 2017. <https://cds.cern.ch/record/2293646>.

INVITED SPEAKERS ABSTRACTS

**Maveraünnehir, Horasan, Harezm Havzasında Muhtelif İlimlerin
Birlikte Gelişmesi, Meşhur Alimlerin Yetiştirilmesi,
Kıymetli Eserlerin Yazılması Üzerine Bir Analiz**

Kemal Yavuz ATAMAN

Müstakil Akademisyen, Sosyal Siyaset Doktoru

kemal.ataman59@gmail.com

Turan coğrafyasının en önemli ilim, kültür havzası Maverāünnehir, Horasan ve Harezm topraklarıdır. Türklerin İslamiyeti kabul etmesinden sonra Müslüman Türk Devletlerinde birçok farklı ilim eş zamanlı olarak gelişmiştir. Özellikle 7-11 yüzyıllarda klasik ilimler, tıp, matematik, fizik, kimya, biyoloji, astronomi alanlarında yetişen Biruni, Harezmi, Fergani, Cürcani, Razi, İbn-i Sina, Hazini, Cabir b. Hayyan, Sabit b. Kurra, Ömer Hayyam, Nasreddin Tusi, Cezeri, Uluğ Bey gibi alimler büyük ilmi keşifler, icadlar yapmışlar, eserleriyle bilim tarihinde yer almışlardır. Aynı dönemlerde tefsir, hadis, fıkıh, kelam, tasavvuf gibi dini ilimlerde İmam Buhari, İmam Tirmizi, İmam Maturidi, Zemahşeri, Nesefi, Mergilani, Serahsi, Ahmed Yesevi, Necmeddin Kübra, Yusuf Hemedani, Bahaeddin Nakşibend gibi alimler eserleriyle, talebeleriyle bir okul ve ekol olmuşlardır. Yine Felsefe, mantık, edebiyat, musiki alanlarında İbn-i Sina, Farabi, Ali Şir Nevayi gibi büyük alimler insanlığı aydınlatmışlardır.

İlmi çalışmalara karşı Türklerin öğrenme, araştırma çabaları ve istekleri, şevk ve motivasyonları müslüman olduktan sonra tezahür etmiştir. Devletlerin başında bulunan Emir, Sultan, Han ve Beğlerden halka doğru toplumun görülen ilim sevgisinin ve gayretinin kaynağı şüphesiz, ilme ehemmiyet veren, insanlığa faydalı olmayı teşvik eden Kuran ve Hadislerdir. Maverāünnehir, Horasan ve Harezm havzasında ilimlerin tekamülü, terakkisi, inkişafı, intişarında çeşitli saikler vardır. Bu topraklara İslam ilk elden seyyidler, sahabe ve tabiin nesliyle gelmiş, yerel halk dini aslından, kaynağından, özünden almıştır. Toplumun imarı ve nizamında Türk tasavvuf tarihinin öncü sufilere belirleyici olmuş, insanlığa faydalı olmak gayesini, ilmin gereğini hayatın önceliği olarak insanlara sunmuşlardır. Klasik ilimlerle uğraşan alimlerin dini bilgilere, dini ilimlerle meşgul olanların klasik ilimlere vukufiyeti her iki sahada tebarüz etmiştir. Mesela, filozof hekimler, matematikçi fakihler, fizikçi ilahiyatçılar dikkat çekmektedir. Nizamiye Medreseleri, Memun Akademisi, Beytü'l Hikme gibi merkezler ilmi faaliyetlerin gelişmesine ve alimler arasında irtibatların artmasına vesile olmuştur.

Savaşlardan sonra fethedilen yerlerin ileri gelen alim ve sanatkarları belirli şehirlere ve başkentlere aktarılmış, saraylarda itibar görmüş ve medreselerde yer almışlardır.

Bölgede hakimiyet tesis eden Karahanlılar, Gazneliler, Selçuklular, Harzemşahlar, Samaniler, moğol istilasından sonraki yüzyıllarda Timuriler, Şeybaniler ilmi çalışmalarını teşvik etmişler, alimlere önem vermişler, sahip çıkmışlar, destek olmuşlardır. Birçok Emir, Sultan, Han ve Beğ iyi eğitilmiş, ilmi birikime sahip şahsiyetlerdir. Saraylar alimlerin buluşma mekanı haline gelmiştir. Devletlerin ekonomik güçlerine göre medreselerin, kütüphanelerin yapılmasına, eserlerin yazılmasına, alimlerin maişetine kaynak ayırılmasına önem verilmiştir. Göçebe hayatı yaşayan Türklerin şehirleşmesiyle halkın ilme ve alimlere desteklerini başlatmış, alimlerin itibarını yükseltmiştir. Ünlü seyyahlar İbni Batuta ve İbni Fadlan seyahatnamelerinde halkın ilme iştiafına işaret etmektedirler. Farklı kültürlerle, dillere, ırklara, dinlere mensup insanların birlikte yaşadığı Semerkant, Buhara, Gurgench, Hive, Belh, Merv gibi şehirler ilmi çalışmaların ve alimlerin merkezi haline gelmiştir. Vakıf müessesesinin gelişmesiyle ilim yolcularına, alimlere, medreselere, eserlere zenginler ve eşraf hamilik yapmıştır. Coğrafyanın özelliğinden, zirai, ticari ve içtimai ihtiyaçlar ve sebeplerden dolayı matematik, astronomi ilimleri gelişmiştir. Türklerin seyahat emniyetini sağladığı zamanlarda İpek Yolu ve Hint Baharat Yolu ilim, irfan, sanat yolu haline gelmiş, alimler güven içinde ilim ve kültür merkezlerine yolculuklar yapmışlar, eserlerin intikali, alimlerin münasebetleri kolaylaşmıştır. Bu havzada Mezopotamya, Mısır, Hind, İran, Roma, Eski Yunan, Çin ve Endülüs medeniyet birikimlerinden faydalanılmış, ilim ve kültür merkezleriyle bilgi akışı sürmüştür.

Bu çalışmada ilim havzası Maveraünnehir, Horasan ve Harezm'de klasik, dini ve içtimai ilimlerin gelişmesi, büyük alimlerin yetişmesi ve kıymetli eserlerin aynı zamanlarda yazılmasının saikleri ve sebepleri değerlendirilerek bir analiz yapılmaya çalışılacaktır.

Anahtar Kelimeler: İlim, alim, Turan, Maveraünnehir, Horasan, Harezm.

A Decade of Machine Learning Research: Evaluation Metrics, Taxonomies and Bibliometrics Analysis

Amir MOSAVİ

Obuda University, Hungary

amir.mosavi@uni-obuda.hu

Machine learning (ML) has revolutionised many scientific and industrial domains in the last ten years, spurring automation and innovation. However, thorough evaluation is necessary to ensure the efficacy and dependability of ML models, so it is crucial to comprehend important metrics and methodologies. With an emphasis on bibliometric analysis, taxonomies, and evaluation metrics, this presentation explores the trends in machine learning research over the last ten years. We classify machine learning algorithms according to their approaches and examine key evaluation metrics that have influenced model evaluation in different fields. Furthermore, within the ML community, bibliometric analysis identifies key research trends, significant publications, and networks of collaboration. In addition to growing ethical concerns and the need for standardised benchmarking, our findings show changes in evaluation methodologies, the continued dominance of deep learning, and the growing emphasis on explainability and fairness.

Keywords: Machine learning, artificial intelligence.

References

- [1] Nie, Z., Xiao, Z., Wang, T., Chronopoulos, A. T., Andonie, R., and Mosavi, A. (2025). *TIPS: A Text Interaction Evaluation Metric for Learning Model Interpretation*. Expert Systems with Applications, 128184.
- [2] Dénes-Fazakas, L., Simon, B., Hartvég, Á., Szilágyi, L., Kovács, L., Mosavi, A., and Eigner, G. (2024). *Personalized food consumption detection with deep learning and Inertial Measurement Unit Sensor*. Computers in Biology and Medicine, 182, 109167.

On the L_1 Data Dirichlet Problem for the Nonuniform Parabolic Equations of Second Order

Ferman MAMEDOV^{1,2}

¹*Institute of Mathematics and Mechanics of MSE, Baku AZ1141, B. Vahabzade, 9, Azerbaijan*

²*Azerbaijan State Oil and Industry University, Baku AZ1010, Azadlig, 16/21, Nasimi*

farman-m@mail.ru

For the nonuniform parabolic equation of second order the first boundary value is considered. The data on the right hand side of the equation

$$\partial_t u - Lu = f; \quad Q_T = [0, T) \times D$$

$$u(0, z) = g(z), \quad z \in D$$

and the initial value functions f, g are of the L_1 class. The sufficient conditions are proposed to the degeneration in the spartial part of the equation in terms of the ball conditions on R^n . For the purpose are proved new inequalities of the Poincare-Sobolev type having nonuniform degenerate gradient. For the subject we refer to the works^{1,2}.

Keywords: Non-uniform parabolic equation, Sobolev-Poincare inequality, existence and uniqueness.

References

- [1] Mamedov, F., Mansurro, S. (2022). *Sobolev Inequality with Non-uniformly Degenerating Gradient*, Elektronik J. Qual. Th. Diff. Equ.
- [2] Mamedov, F. (2022). *On Harnack Inequality and Hölder Continuity to the Degenerate Parabolic Equations*. J. Diff. Equ., 340, 521-556.

ORAL PRESENTATION ABSTRACTS

TURAN-FUNDAMENTAL SCIENCES SYMPOSIUM
TURAN-TEMEL BİLİMLER SEMPOZYUMU

Honorary Chair/Onursal Başkan: Prof. Dr. Aziz SANCAR

MATHEMATICS

2-Parameter Generalized Quaternions with Generalized Tribonacci Numbers Components

Zehra İŞBİLİR¹, Nurten GÜRSES²

¹Department of Mathematics, Faculty of Arts and Sciences, Düzce University, 81620, Düzce, Türkiye

²Department of Mathematics, Faculty of Arts and Sciences, Yıldız Technical University, 34220, Istanbul, Türkiye

zehraisbilir@duzce.edu.tr

In this presentation, our main goal is to introduce the combination of generalized quaternions and generalized Tribonacci numbers which are quite a big special recurrence family for third-order special recurrence numbers. We define the non-negative and negative subscripted generalized quaternions with generalized Tribonacci numbers components. Then, we give some special cases and achieve some properties with respect to these type new quaternions such as; Binet formulas, generating functions, exponential generating functions, Poisson generating functions, summation formulas, matrix formulas, and special determinant properties.

Keywords: Generalized Tribonacci numbers, generalized quaternions, generalized quaternions with generalized Tribonacci numbers components.

References

- [1] Cerda-Morales, G. (2017). *On a Generalization for Tribonacci Quaternions*. Mediterr. J. Math., 14(239), 1-12.
- [2] İşbilir, Z., Gürses, N. and Tosun, M. (2025). *On the 3-Parameter Generalized Quaternions with Generalized Tribonacci Numbers Components*. Filomat, 39(9), 3003-3027.
- [3] Şentürk, T. D., Ünal, Z. (2022). *3-Parameter Generalized Quaternions*. Comput. Methods Funct. Theory, 22(3), 575–608.
- [4] Hamilton, W. R. (1844). *On Quaternions; or On a New System of Imaginaries in Algebra*. Philos. Mag., 25(3), 489-495.
- [5] Jafari, M., Yaylı, Y. (2015). *Generalized Quaternions and Their Algebraic Properties*. Commun. Fac. Sci. Ank. Series A1, 64(1), 15-27.
- [6] Soykan, Y. (2020). *On Sum Formulas for Generalized Tribonacci Sequence*. Journal of Scientific Research and Reports, 26(7), 27-52.
- [7] Soykan, Y. (2020). *A Study on Generalized (r, s, t) -Numbers*, MathLAB Journal, 7, 101-129.
- [8] Soykan, Y. (2021). *On the Recurrence Properties of Generalized Tribonacci Sequence*. Earthline J. Math. Sci., 6(2), 253-269.
- [9] Soykan, Y. (2020). *Summing Formulas for Generalized Tribonacci Numbers*. Univers. J. Math. Appl., 3(1), 1-11.
- [10] Soykan, Y. (2020). *Generalized Tribonacci Numbers: Summing Formulas*. International Journal of Advances in Applied Mathematics and Mechanics, 7, 57-76.

Calculus on Real-variable Split Quaternion-valued Functions

Edanur ERGÜL ARSLAN¹, Salim YÜCE²

¹Mathematics, Marmara University, Istanbul, 34722, Türkiye

²Mathematics, Yıldız Technical University, Istanbul, 34220, Türkiye

edanur.ergul@marmara.edu.tr

In this study, we investigate fundamental analytical concepts within the framework of split quaternions, focusing specifically on real-variable split quaternion-valued functions. Although the literature includes various studies on split quaternion-variable functions, the definitions and explanations of limit, continuity, and certain derivative concepts remain either sufficiently undeveloped or ambiguous. Addressing this gap, we define the concepts of limit, derivative, P-limit, P-derivative, and integral for split quaternion-valued functions of a real variable, and give several theorems with proofs, including a theorem that determines the conditions under which the classical limit and the P-limit coincide. Additionally, numerous examples are provided to explain the concepts of P-limit, P-derivative and integral of elementary functions in particular.

Keywords: Split Quaternions, real-variable split quaternion-valued functions, limit, derivative, integral.

References

- [1] Cockle, J. (1849). *On Systems of Algebra Involving More than One Imaginary; and on Equations of the Fifth Degree*. Philosophical Magazine Series 1, 35, 434-437.
- [2] Yüce, S. (2020). Sayılar ve Geometri. Pegem Academy Publishing.
- [3] Levent, K., Yaylı, Y. (2007). *Split Quaternions and Rotations in Semi Euclidean Space E_2^4* . Journal of the Korean Mathematical Society, 44(6), 1313-1327.
- [4] Masrouri, N., Yaylı, Y., Faroughi, M. H. and Mirshafizadeh, M. (2011). *Comments on Differentiable over Function of Split Quaternions*. Revista Notas de Matemática, 7(2), 128-134.
- [5] Libine, M. (2011). *An Invitation to Split Quaternionic Analysis*. In: Sabadini, I., Sommen, F. (eds) Hypercomplex Analysis and Applications. Trends in Mathematics. Springer, Basel, 161-180.
- [6] Zill, D. G., Shanahan, P. D. (2003). *A First Course in Complex Analysis with Applications*. Jones and Bartlett Publishers.

Characteristic Roots, Eigenvalues and Eigenvectors of a Certain Type of 4×4 Dual Tridiagonal Matrices

Pelin DURSUN¹, Nurten GÜRSES²

¹*Istanbul Ticaret University, Faculty of Humanities and Social Science, Department of Mathematics,
34310, Istanbul, Türkiye*

²*Department of Mathematics, Faculty of Science and Arts, Yildiz Technical University,
34220, Istanbul, Türkiye
pdursun@ticaret.edu.tr*

Dual number matrices serve as effective tools in modeling problems in brain science and multi-agent formation control. In the literature one can find determinants, characteristic polynomials, eigenvalues, and eigenvectors of dual number matrices. An $n \times n$ dual number matrix may have exactly n eigenvalues, none, or infinitely many. In this presentation, by taking a certain type of 4×4 dual number tridiagonal matrix, we demonstrate that it has exactly four eigenvalues, which directly result from structured combinations of the eigenvalues of its real matrix components. Finally, we present illustrative examples that support and clarify our theoretical findings.

Keywords: Dual numbers, characteristic polynomial, eigenvalues, tridiagonal matrix.

References

- [1] Yüce, S. (2020). Sayılar ve Geometri. Pegem Academy Publishing, Ankara.
- [2] Öteleş A., Akbulak, M. (2013). *Positive Integer Powers of Certain Complex Tridiagonal Matrices*. Appl. Math. Comput., 219(21), 10448–10455.
- [3] Qi L., Cui, C. (2023). *Eigenvalues and Jordan Forms of Dual Complex Matrices*. Commun. Appl. Math. Comput., 1–17.
- [4] Qi L., Cui, C. (2024). *Eigenvalues of Dual Hermitian Matrices with Application in Formation Control*. SIAM J. Matrix Anal. Appl., 45(4), 2135–2154.
- [5] Ling C., Qi L. (2024). *Quasi-Determinant and Right Eigenvalues of Dual Quaternion Matrices*. arXiv:2404.19348.

Comparative Analysis of Türkiye's Road Traffic Accidents Using Data Mining Techniques

Sema Nur ÇEPE, Sema Akın BAŞ

Yildiz Technical University, Faculty of Arts and Science, Department of Mathematics,

Istanbul, Türkiye

nur.cepe@std.yildiz.edu.tr

Traffic accidents, especially involving unprotected vehicles such as motorcycles and mopeds, are significant public safety concern due to their high fatality and injury rates. In this study, motorcycle accidents in Türkiye were analyzed by applying classification techniques in the field of data mining. Three algorithms, namely Logistic Regression, Naive Bayes and Random Forest, were applied to the data on fatal and injury traffic accidents provided by the Traffic Department of the General Directorate of Security. The performance of the studied models was evaluated using multiple evaluation criteria in order to determine the most accurate method for predicting accident outcomes. The analysis showed that the Random Forest algorithm achieved the highest classification accuracy for both fatal and non-fatal accidents. In addition, the main risk factors affecting the occurrence of accidents, including driver errors, accident time and environmental conditions, were identified. These results provide helpful information for decision makers who want to improve motorcycle safety and create focused prevention strategies.

Keywords: Data mining, traffic accidents, motorcycle accidents, Python.

References

- [1] Turkish Statistical Institute (TUIK). (2024). *Traffic Accident Statistics*. Retrieved from <https://www.tuik.gov.tr>
- [2] General Directorate of Security (EGM). (2024). *Traffic Accident Data*. Retrieved from <https://www.egm.gov.tr>

The Moment of Inertia of Generated Curves: A Holditch-Type Theorem

Kübra ÇETİNER¹, Salim YÜCE²

¹Computer Programming, Istanbul Gelisim Vocational School, Istanbul Gelisim University, Istanbul,
Türkiye

²Department of Mathematics, Faculty of Arts and Sciences, Yıldız Technical University, Istanbul,
Türkiye

kcetinberk@gelisim.edu.tr

Holditch's Theorem states that when a fixed-length segment rotates inside a convex closed curve, a point on the segment traces a closed curve whose enclosed area is smaller than that of the original curve by πpq . In our previous work, the moment of inertia of parallel curves was examined in this context.

In this study, we focus on generated curves, which are constructed from a base curve using specific geometric rules but are not equidistant from it. We calculate the moment of inertia of these curves. Based on the results obtained, we present a Holditch-type theorem involving the moment of inertia. The aim is to explore the relationship between generated curves and their inertial properties, offering a new interpretation parallel to classical geometric results.

Keywords: Holditch's theorem, Holditch surface, generated curves, moment of inertia.

References

- [1] Çetinberk, K., Yüce, S. (2024). *The Moment of Inertia of Parallel Curves and New Sight into Holditch's Theorem*. SISA 2024: 6th International Symposium on Innovations in Scientific Areas, Türkiye.
- [2] Plata, D. R. (2019). *On Holditch's Theorem and Related Kinematics*. Doctoral dissertation, Universitat de Valencia. Facultat de Ciències Matemàtiques.
- [3] Monterde, J., Rochera, D. (2019). *Holditch's Theorem in 3D Space*. Results in Mathematics, 74(3), 110.

Do Oil Prices Have a Chaotic Effect on Unemployment and Inflation

Melike BİLDİRİCİ¹, Merve ÇOLAK²

¹ *Department of Economics, Faculty of Economics and Administrative Sciences, Yildiz Technical University, 34220, Istanbul, Türkiye*

² *Graduate School of Social Sciences, Yildiz Technical University, 34220, Istanbul, Türkiye*
mervecolk95@gmail.com

Oil prices have a significant impact on macroeconomic variables on a global scale. Although unemployment and inflation rates, in particular, are sensitive to fluctuations in oil prices, the nature of this relationship can be non-linear and chaotic. Research exploring the chaotic nature of this relationship remains limited. In linear time series analyses, chaotic behaviors are often treated as stochastic, and unexplained observations are dismissed as random anomalies. However, while some papers examined oil prices through the pass-through mechanism, research on their chaotic structure remains limited, with notable contributions from Adrangi et al.(2001), Komijani et al(2011). In this article, the effects of oil prices on unemployment and inflation will be examined within the framework of a chaotic dynamic structure. Within the scope of the analysis, the rescaled range (R/S) of Lo and the Mandelbrot-Wallis methods will determine fractionality and long-term dependence. Renyi, Shannon, Tsallis and HCT entropy tests will be used to determine entropy. The Kolmogorov-Sinai complexity (KSCM) method will identify evidence of the complexity of variables. The Hurst exponents will be used to investigate evidence of mean reversal, chaos, and the Brownian movement. The Lyapunov exponent (LE) will discover evidence of chaos. The aim is to reveal how changes in oil prices can create disorder or chaos in unemployment and inflation dynamics, using the most up-to-date and advanced tools of time series analysis.

Keywords: Oil prices, chaos theory, unemployment, inflation, entropy.

References

- [1] Adrangi, B., Chatrath, A. and Raffiee, K. (2001). *Chaos in Oil Prices?* Energy Economics.
- [2] Komijani, A., et al. (2011). *Nonlinear Dynamics of Oil Markets*. Applied Energy.

Genel DSW Sistemine Genişletilmiş Jacobi Eliptik Fonksiyon Yaklaşımı: Tam Çözümler, Kararlılık ve Simülasyonlar

Nisa ÇELİK

Bursa Uludağ Üniversitesi, Fen Edebiyat Fakültesi, Matematik Bölümü, 16059, Nilüfer, Bursa, Türkiye

nisa@uludag.edu.tr

Bu çalışmada, su dalgalarının bir modeli olarak kabul edilen genel Drinfel'd–Sokolov–Wilson (DSW) sisteminin ilerleyen dalga çözümleri ve dalga dinamiği araştırıldı. DSW sistemi, dalga dönüşümüyle daha basit bir adi diferansiyel denklem sistemine dönüştürüldü. Ardından genişletilmiş Jacobi eliptik fonksiyon açılım yöntemi ve eliptik rasyonel fonksiyon açılım yöntemi uygulanarak tam çözümler elde edildi. Çözümlerin davranışları, farklı parametreler altında Maple™ programı ile yapılan iki ve üç boyutlu sayısal simülasyonlarla analiz edildi. Jacobi eliptik fonksiyonlar, fizik ve mühendislikte geniş uygulamalara sahip olup su dalgaları, gelgitler ve akustik dalgaların modellenmesinde kullanılmaktadır. Son olarak elde edilen çözümlerin yeteneğini ortaya koymak amacıyla kararlılık özelliği uygulanarak çözümler test edildi.

Anahtar Kelimeler: Drinfel'd-Sokolov-Wilson (DSW) sistemi, Jacobi eliptik fonksiyon, tam çözüm.

Referanslar

- [1] Zhang, H. (2007). *Extended Jacobi Elliptic Function Expansion Method and its Applications*. Communications in Nonlinear Science and Numerical Simulation, 12, 627-635.
- [2] Wang, Qi, Chen Y., Hongqing Z. (2005). *A New Elliptic Function Rational Expansion Method and its Application To (1+1)-Dimensional Dispersive Long Wave equation*. Chaos Solitons & Fractals, 477-482.
- [3] Wazwaz, A. M. (2009). *Partial Differential Equations and Solitary Waves Theory*. Nonlinear Physical Science, Saint Xavier University, Chicago, IL 60655.
- [4] Bashar, Md. H., Yiasir S. M. A., Islam S. M. R, and Rahman M. M. (2022). *Wave Solutions of the Couple Drinfel's-Sokolov-Wilson Equation: New Wave Solutions and Free Parameters Effect*. Journal of Ocean Engineering and Science. <https://doi.org/10.1016/j.joes.2022.05.003>.

Bazı Antiviral Moleküllerin Topolojik Karakterizasyonu

Betül ÖZTÜRK, Elif HANEDAN, and Fatma Özen ERDOĞAN

Bursa Uludağ Üniversitesi, Fen-Edebiyat Fakültesi, Matematik Bölümü, 16059, Bursa, Türkiye

betullozturk.97@gmail.com

Bu çalışmada, seçilen bazı antiviral bileşiklerin¹, yapısal özellikleri, graf teori temelli yaklaşımlarla analiz edilmiştir. Her bir molekül, atomların köşelere ve bağların kenarlara karşılık geldiği moleküler bir graf olarak modellenmiştir. Çalışmada ayrıca, temel moleküler grafların yanı sıra, kenar-komşuluk grafları gibi türetilmiş yapılar da incelenmiştir. Moleküler graflar için, graf teoride sıkça kullanılan temel topolojik indeksler²⁻³ ile Platt sayısı⁴ hesaplanmış ve aralarındaki ilişkiler karşılaştırmalı olarak araştırılmıştır. Ayrıca, bu indeksler ile moleküllerin belirli fizikokimyasal özellikleri arasındaki bağlantılar korelasyon analizleriyle ortaya konmuştur.

Elde edilen bulgular, istatistiksel ve görsel analizlerle desteklenmiş olup, topolojik indeksler arasında yapısal benzerliklerin ve farklılıkların bulunduğunu göstermektedir. Bu karşılaştırmalı analiz; moleküler yapıların sayısal temsiline ilişkin anlayışı derinleştirmekte ve graf teori kuramına dayalı bu yaklaşımın yapısal sınıflandırma ile öngörüselleştirme alanlarındaki potansiyelini vurgulamaktadır.

Anahtar Kelimeler: Antiviral bileşikler, topolojik indeksler, platt sayısı, kenar-komşuluk grafi.

Referanslar

- [1] De Clercq, E., Li, G. (2016). *Approved Antiviral Drugs over the Past 50 Years*. Clinical microbiology reviews, 29(3), 695-747.
- [2] Gutman, I. (2013). *Degree-Based Topological Indices*. Croatica chemica acta, 86(4), 351-361.
- [3] Nikolić, S., Trinajstić, N. (1995). *The Wiener Index: Development and Applications*. Croatica Chemica Acta, 68(1), 105-129.
- [4] Sarika, S. G., Manjusha, R. (2025). *Graph Theory and Molecular Behavior: Utilizing Platt Numbers and Polynomial Approximations for Boiling Point Estimations*. Engineering Reports, 7(2), e13124.

Affine Factorable Surfaces in Euclidean 4-Space E^4

Emine KOPARAL, Betül Bulca SOKUR

Bursa Uludağ University, Arts and Science Faculty, Department of Mathematics, Bursa, Türkiye

eminekop16@gmail.com

In this study the affine factorable surfaces in Euclidean 4-space were defined. We have obtained the Gaussian, normal and mean curvature of the affine factorable surfaces in E^4 . Further, the classification of surfaces with flat, normal flat and minimal affine factorable surfaces was given.

Keywords: Affine factorable surfaces, Gaussian curvature, mean curvature.

References

- [1] Aminov, Yu. (1994). *Surfaces in E^4 with a Gaussian Curvature Coinciding with a Gaussian Torsion up to the Sign*. Math. Notes, 56, 1211-1215.
- [2] Aydın, M. E., Erdur, A., and Ergüt, M. (2020). *Affine Factorable Surfaces in Isotropic Spaces*. TWMS J. Pure Appl. Math., 11, 72-88.
- [3] Bulca, B., Arslan, K. (2013). *Surfaces Given with Monge Patch in E^4* . Journal of Mathematical Physics, Analysis, Geometry, 4, 435-447.
- [4] Medjahdi, B., Zoubir, H. (2021). *Affine Factorable Surfaces in the Three-Dimensional Simply Isotropic Space*. IAENG Int. J. of Appl. Math., 51(2), 366-371.
- [5] Meng, H., Liu, H. (2009). *Factorable Surfaces in 3-Minkowski Space*. Bull. Korean Math. Soc., 46(1), 155-169.
- [6] Saad, M., Abdel-Aziz, H., and Ali, H. (2023). *Investigation of Affine Factorable Surfaces in Pseudo-Galilean Space*. Wseas Transactions on Mathematics., 22, 666-673.
- [7] Yu, Y., Liu, H. (2007). *The Factorable Minimal Surfaces*. In: Proceedings of The Eleventh International Workshop on Diff. Geom., 11, 33-39
- [8] Zong, P., Xiao, L., and Liu, H.L. (2015). *Affine Factorable Surfaces in Three-Dimensional Euclidean Space*. Acta Math. Sinica Chinese Serie 58(2), 329-336.

Monge Surfaces with Their Applications

Beyza SARUHAN¹, Kadri ARSLAN², and Betül BULCA SOKUR²

¹Bursa Uludağ University, Institute of Science, Department of Maths., Bursa, Türkiye

²Bursa Uludağ University, Art and Science Faculty, Department of Maths., Bursa, Türkiye

16beyzasaruhan@gmail.com

Analytical surfaces have an important place in differential geometry. These surfaces are frequently used in geometric design. The ability to describe such surfaces with parametric equations is very common for modeling with "computer-aided geometric design". In this study, a characterization of Monge surfaces, which is an important type of analytical surfaces, is discussed. Monge surfaces are surfaces that are examples of carved surfaces. These surfaces are quite impressive in terms of aesthetics in architectural design and exterior cladding of buildings.

Keywords: Monge surface, carved surface.

References

- [1] Brander, D., Gravesen, J. (2018). *Monge Surfaces and Planar Geodesic Foliations*. J. Geom. 109, 1-14.
- [2] Krivoschapko S. N., Ivanov V. N. (2015). *Encyclopedia of Analytical Surfaces*. Springer Int. Pub. Switzerland.
- [3] Salkowski E. (1909). *Zur Transformation von Raunkurven*. Math. Ann. 66, 517-557.
- [4] Gil-oulbé1, M., Ndomilep, A. J. I. (2020). *Geometry and Classification of Carved Monge Surfaces*. Journal of Physics: Conference Series, 1687, 1-10.
- [5] Gray, A. (1993). *Modern Differential Geometry of Curves and Surfaces*. CRC Pres Inc., USA, 398 pp.
- [6] Monge, G. (1850). *Application De L'Analyse a La Geometrie*. Bachelier, Paris.

Bazı Özel Afın Yapıların ve Duallerinin Kromatik İndislerinin Bulunması Üzerine

Abdurrahman DAYIOĞLU, Fatma ÖZEN ERDOĞAN, and Basri ÇELİK

Bursa Uludağ Üniversitesi Fen-Edebiyat Fakültesi, Matematik Bölümü, Bursa, Türkiye

dayioglu@uludag.edu.tr

Bu çalışmada, klasik graf teorisinin temel kavramlarından olan köşe ve kenar renklendirme ilkeleri afın yapılara uyarlanmış; bu doğrultuda yeni renklendirme algoritmaları geliştirilmiş ve en uygun renklendirme sayıları mertebe derecelerine bağlı olarak belirlenmiştir. Bazı özel afın yapılarıdaki nokta ve doğruların renklendirilmesine yönelik çeşitli genel sonuçlar ve teoremler elde edilmiştir. Araştırma kapsamında, geometrik yapılar graf teorisinin köşe ve kenar renklendirme yaklaşımları çerçevesinde ele alınarak, bu iki alan arasındaki yapısal ilişkiler derinlemesine incelenmiştir. Özellikle köşe ve kenar renklendirme prensiplerinin afın yapılara genişletilmesi, kombinatoryal matematik ile geometri arasındaki etkileşime yeni bir bakış açısı kazandırmakta ve mevcut renklendirme tekniklerine katkı sağlayacak nitelikte özgün sonuçlar sunmaktadır.

Anahtar Kelimeler: Afın düzlem, kromatik indis.

Referanslar

- [1] Erdős, P. (1981). *On The Combinatorial Problems Which I Would Most Like to See Solved*. Combinatorica, 25–42. <https://doi.org/10.1007/BF02579174>.
- [2] Beutelspacher, A., Jungnickel, D., and Vanstone, S. A. (1989). *On The Chromatic Index of a Finite Projective Space*. Geom Dedicata 32, 313–318. <https://doi.org/10.1007/BF00147923>.
- [3] Araujo-Pardo, G., Kiss, G., Rubio-Montiel, C., and Vázquez-Avila, A. (2017). *On Chromatic Indices of Finite Affine Spaces*. Ars Mathematica Contemporanea, 16(1), 67-79.
- [4] Xu, L., Feng, T. (2023). *The Chromatic Index of Finite Projective Spaces*. J. Combin. Des., 31, 432–446. <https://doi.org/10.1002/jcd.21904>.
- [5] Meszka M. (2013). *The Chromatic Index of Projective Triple Systems*. J. Combin. Designs, 21, 531–540. <https://doi.org/10.1002/jcd.21368>.
- [6] Ozeki, K. (2022). *Kempe Equivalence Classes of Cubic Graphs Embedded on the Projective Plane*. Combinatorica, 42 (Suppl 2), 1451–1480. <https://doi.org/10.1007/s00493-021-4330-2>.

Optical Bullets for the (3+1)-Dimensional Perturbed Nonlinear Schrödinger Equation having the Cubic-Quintic Laws and the Spatio-Temporal Dispersion

Neslihan ÖZDEMİR¹, Müslüm ÖZİŞİK²

¹Department of Mathematics, Yildiz Technical University, Istanbul, Türkiye

²Department of Mathematical Engineering, Yildiz Technical University, Istanbul, Türkiye

ozdemirn@yildiz.edu.tr

This study includes the soliton dynamics governed by the (3+1)- dimensional perturbed nonlinear Schrödinger equation having the cubic-quintic laws and the spatio-temporal dispersion. This extended form of the nonlinear Schrödinger equation is particularly significant for modeling the propagation of ultrashort optical pulses in nonlinear media, where conventional dispersion plays a minimal role, but higher-order and nonlinear effects become dominant. To obtain optical bullets, we employ the addendum to the Kudryashov's method. By applying suitable constraints on model parameters, we establish the conditions necessary for the existence of various optical bullets. In addition, a comprehensive analysis of modulation instability is conducted to further elucidate the stability characteristics of the model. The results provide valuable theoretical insights into the complex behavior of pulse propagation in advanced photonic structures, with implications for the design and optimization of nonlinear optical waveguides and fiber communication systems.

Keywords: The Schrödinger model, optical bullets, cubic-quintic laws, modulation instability analysis.

References

- [1] Zakharov, V. E. E., Manakov, S. V. (1974). *On the Complete Integrability of a Nonlinear Schrödinger Equation*. Theoretical and Mathematical Physics, 19(3), 551-559.
- [2] Agrawal, G. P. (2000). *Nonlinear Fiber Optics*. In Nonlinear Science at the Dawn of the 21st Century (pp. 195-211). Berlin, Heidelberg: Springer Berlin Heidelberg.
- [3] Zayed, E. M., Gepreel, K. A., and Alngar, M. E. (2021). *Addendum to Kudryashov's Method for Finding Solitons in Magneto-Optics Waveguides to Cubic-Quartic NLSE with Kudryashov's Sextic Power Law of Refractive Index*. Optik, 230, 166311.
- [4] Ozisik, M., Secer, A., Bayram, M., and Aydin, H. (2022). *An Encyclopedia of Kudryashov's Integrability Approaches Applicable to Optoelectronic Devices*. Optik, 265, 169499.
- [5] Wang, G. A. (2021). *New (3 + 1)-dimensional Schrödinger Equation: Derivation, Soliton Solutions and Conservation Laws*. Nonlinear Dyn 104, 1595–1602.
- [6] Wang, G., Wang, X., Guan, F., and Song, H. (2023). *Exact Solutions of an Extended (3+1)-dimensional Nonlinear Schrödinger Equation with Cubic-Quintic Nonlinearity Term*. Optik, 279, 170768, ISSN 0030-4026.
- [7] Zayed, E. M., Ahmed, M. S., Arnous, A. H., and Yıldırım, Y. (2025). *Novel Soliton Solutions of the (3+1)-dimensional Stochastic Nonlinear Schrödinger Equation in Birefringent Fibers*. Chaos, Solitons & Fractals, 194, 116152, ISSN 0960-0779.

TBMM Maarif Vekâletinin 1920 Yılında Aldığı Geometri Öğretimi Kararları Üzerine

Basri ÇELİK

Bursa Uludağ Üniversitesi, Fen-Edebiyat Fakültesi, Matematik Bölümü, Bursa, Türkiye

basri@uludag.edu.tr

Bu çalışmada TBMM Maarif Vekâleti Hicri 1339 (Miladi 1920) yılında ilkokullarda geometri öğretiminin nasıl olması gerektiğine dair aldığı kararlar¹ ve bu karar metninin daha önce yayınlanmış benzerleri² üzerinde durulacaktır. Aradan 100 yıldan daha fazla bir zaman geçmesine rağmen, geometri öğretiminde günümüzde uygulanması arzulanan yöntemlerin Payitaht İstanbul'un işgal altında olduğu, ülkenin her yerinde savaşın hüküm sürdüğü bir dönemde alınan bu kararlarla uygulamaya konulması eğitime verilen önemin göstergesidir.

Anahtar Kelimeler: Geometri öğretimi, matematik tarihi, Osmanlı Devleti'nde eğitim.

Referanslar

- [1] Maarif Vekâleti. (Hicri 1339). *Mekâtib-i İbtidaiyyede Hendesenin Usûl-i Tedrisi*. Matbaa-i Amire, İstanbul, Adet 7.
- [2] Hakkı, İ. (Hicri 1329). *Hendeseyi Nasıl Tedris Etmeli?* Tedrisat Mecmuası, Sene 4, Cilt 4, Sayı 22, 133-139.

Competitive Systems via Hamiltonian Framework and Artificial Neural Estimation

Hande USLU TUNA¹, Bedia Merih ÖZÇETİN²

¹*Department of Mathematics, Faculty of Art and Sciences, Yıldız Technical University, Istanbul, Türkiye*

²*Department of Avionics, Faculty of Applied Sciences, Istanbul Gelisim University, Istanbul, Türkiye*

usluh@yildiz.edu.tr

In this study, the dynamical behavior of competitive Lotka-Volterra systems has been investigated by analyzing the curvature properties of their integral curves in a Hamiltonian framework¹. In the literature²⁻³, the three and four species systems have been explored in terms of their geometric properties, especially those reflecting ecological interactions. In recent advances in the analysis of large-scale Lotka-Volterra systems and species persistence, artificial neural networks (ANNs) have been applied to approximate and reconstruct integral curves and curvature properties from partial or noisy observations⁴. Furthermore, Hamiltonian conservation laws have been integrated with data-driven learning techniques to create a unified framework, thus providing a comprehensive representation of both structural and empirical aspects of nonlinear population dynamics⁵. By integrating mathematical theory with machine learning, this study contributes both structural and computational perspective on the behavior of nonlinear systems modeled by Lotka-Volterra equations.

Keywords: Curvatures, competitive systems, Hamiltonian frames, artificial neural networks.

References

- [1] Plank, M. (1995). *Hamiltonian structures for the n-dimensional Lotka-Volterra equations*. J. Math. Phys, 36, 3520–3534.
- [2] Hsu, S., Ruan, S. and Yang, T. (2015). *Analysis of three species Lotka-Volterra food web models with omnivory*. Journal of Mathematical Analysis and Applications, 426(2), 659-687.
- [3] Vaidyanathan, S. (2015). *Global Chaos Synchronization of the Lotka-Volterra Biological Systems with Four Competitive Species via Active Control*. International Journal of PharmTech Research, 8(6), 206-217.
- [4] Clenet, M., Massol, F. and Najim, J. (2023). *Equilibrium and surviving species in a large Lotka-Volterra system of differential equations*. J. Math. Biol. 87 (13).
- [5] Ouahabi, R., Boulezaz, C. (2024). *A novel approach to synchronizing a biological snap oscillator within a fixed time and expanding the method to various chaotic systems*. J Supercomput 80, 18237-18259.

Data-Driven Estimation of Krylov Iterations in Riccati Solvers

Yaprak DERİCİOĞLU^{1,2}

¹Bitlis Eren University, Faculty of Arts and Science, Department of Mathematics, Bitlis, Türkiye

²Yildiz Technical University, Faculty of Arts and Sciences, Department of Mathematics,
Istanbul, Türkiye

yguldogan@beu.edu.tr

This study proposes a machine learning approach to estimate the iteration count needed by Krylov subspace methods in solving differential Riccati equations (DREs). Krylov methods, particularly those using the Arnoldi process, are efficient for large-scale control problems¹⁻². However, stopping criteria based on fixed tolerances may not suit all systems.

We construct a dataset by tracking system features and residual norms during the Arnoldi process³, then train regression models to predict the necessary iteration count⁴. Results show that these models improve convergence estimation and computational efficiency, supporting the integration of data-driven techniques into classical solvers⁵.

Keywords: Riccati equation, Krylov methods, Arnoldi iteration, machine learning, regression, iteration prediction.

References

- [1] Benner, P., Li, J.-R., and Penzl, T. (2008). *Numerical Solution of Large-Scale Lyapunov Equations, Riccati Equations, and Linear-Quadratic Optimal Control Problems*. Numerical Linear Algebra with Applications, 15(9), 755–777.
- [2] Simoncini, V. (2007). *A New Iterative Method for Solving Large-Scale Lyapunov Matrix Equations*. SIAM Journal on Scientific Computing, 29(3), 1268–1288.
- [3] Druskin, V., Knizhnerman, L. (1995). *Krylov Subspace Approximation of Eigenpairs and Matrix Functions in Exact and Computer Arithmetic*. Numerical Linear Algebra with Applications, 2(3), 205–217.
- [4] Xu, W., Li, M., and Zhou, J. (2021). *Machine Learning Techniques for Estimating Iteration Numbers in Numerical Solvers*. Applied Intelligence, 51(5), 3035–3047.
- [5] Karniadakis, G. E., Kevrekidis, I. G., Lu, L., Perdikaris, P., Wang, S., and Yang, L. (2021). *Physics-Informed Machine Learning*. Nature Reviews Physics, 3(6), 422–440.

q-Türev Kullanılarak Tanımlanan Yeni Bir Harmonik Fonksiyon Sınıfının Geometrik Özellikleri

Serkan ÇAKMAK

*Istanbul Gelişim Üniversitesi, İktisadi İdari ve Sosyal Bilimler Fakültesi, Yönetim Bilişim Sistemleri
Bölümü, İstanbul, Türkiye*

secakmak@gelisim.edu.tr

Bu çalışma, q -türev operatörleriyle tanımlanan yeni bir harmonik fonksiyon sınıfının geometrik özelliklerini incelemektedir. Ele alınan sınıfın katsayı sınırları, distorsiyon teoremleri, konvekslik ve yıldızlılık yarıçapları belirlenmiş; ayrıca konveks kombinasyon ve konvolüsyon altında kapalılık özellikleri analiz edilmiştir. Elde edilen sonuçlar, q -analogu harmonik fonksiyonların yapısını daha iyi anlamaya katkı sağlamaktadır.

Anahtar Kelimeler: Harmonik fonksiyonlar, q -türev, geometrik fonksiyon teorisi, konvekslik, yıldızlılık.

Referanslar

- [1] Clunie, J., Sheil-Small, T. (1984). *Harmonic Univalent Functions*. Annales Academiae Scientiarum Fennicae. Series A. I. Mathematica, 9, 3–25.
- [2] Owa, S., Nunokawa, M., Saitoh, H., and Srivastava, H. M. (2002). *Close-to-convexity, Starlikeness, and Convexity of Certain Analytic Functions*. Applied Mathematics Letters, 15(1), 63–69.
- [3] Jackson, F. H. (1909). *On q -functions and a Certain Difference Operator*. Transactions of the Royal Society of Edinburgh, 46(2), 253–281.
- [4] Jackson, F. H., Fukuda, T., and Dunn, O. (1910). *On q -definite Integrals*. Quarterly Journal of Pure and Applied Mathematics, 41(15), 193–203.
- [5] Ahuja, O. P., Çetinkaya, A. (2019). *Use of Quantum Calculus Approach in Mathematical Sciences and its Role in Geometric Function Theory*. AIP Conference Proceedings, 2095, 020001.
- [6] Duren, P. L. (2001). *Univalent Functions*. Springer, 259.
- [7] Silverman, H. (1998). *Harmonic Univalent Functions with Negative Coefficients*. Journal of Mathematical Analysis and Applications. 220(1), 283–289.
- [8] Fejér, L. (1925). *Über die Positivität von Summen*. Acta Scientiarum Mathematicarum (Szeged), 1, 75–86.
- [9] Singh, R., Singh, S. (1989). *Convolution Properties of a Class of Starlike Functions*. Proceedings of the American Mathematical Society, 106(1), 145–152.

Some Characterizations of Gradient Ricci-Yamabe Solitons

Fatma KARACA

Department of Mathematics, Yildiz Technical University, Istanbul, 34220, Türkiye

fatma.karaca@yildiz.edu.tr

In this study, we investigate doubly warped product manifolds admitting gradient Ricci-Yamabe solitons. We find the main relations for a doubly warped product manifold to be a gradient Ricci-Yamabe soliton and demonstrate this result for a physical model of the universe, namely, for the doubly warped spacetime.

Keywords: Gradient Ricci-Yamabe soliton, doubly warped product, spacetime.

Acknowledgement: This research has been supported by Istanbul Beykent University Scientific Research Projects Coordination Unit. Project Number: 2023-24-BAP-03, 2025.

References

- [1] Güler, S., Crasmareanu, M. (2019). *Ricci-Yamabe Maps for Riemannian Flows and Their Volume Variation and Volume Entropy*. Turkish Journal of Mathematics, 43(5), 2631-2641.
- [2] Karaca, F. (2023). *Gradient Ricci-Yamabe Solitons on Warped Product Manifolds*. Filomat, 37(7), 2199-2207.
- [3] Ramos, M. P. M., Vaz, E. G. L. R., and Carot, J. (2003). *Double Warped space-times*. J. Math. Phys. 44, 4839-4865.
- [4] Ünal, B. (2001). *Doubly Warped Products*. Differential Geom. Appl., 15, 253-263.

Matematik Lisans Öğrencilerinin Seçmeli Ders Seçimlerinin Oyun Teorisi ile Matematiksel Modellenmesi

Adem Cengiz ÇEVİKEL, Burcu ALTINTAŞ

Matematik Bölümü, Yıldız Teknik Üniversitesi, 34220, İstanbul, Türkiye

burcu.altintas1@std.yildiz.edu.tr

Bu çalışmada, matematik lisans öğrencilerinin seçmeli ders tercihlerinde etkili olan faktörler oyun teorisi çerçevesinde modellenmiştir. Öğrencilerden anket yoluyla toplanan verilerle, en çok tercih edilen iki ders “oyuncu” olarak belirlenmiş ve öğrencilerin karar kriterleri “strateji” olacak biçimde sınıflandırılmıştır. Analiz sürecinde hem sıfır toplamli hem de sıfır toplamli olmayan oyun modelleri kullanılmış; oyun modellerine stratejik baskınlık, minimax yöntemi ve Nash dengesi gibi çözüm yöntemleri uygulanmıştır. Elde edilen sonuçlar, öğrencilerin ders seçimlerinde bireysel ve sosyal etkenlerin rol oynadığını göstermektedir. Bu modelleme ile müfredat planlamalarında öğrenci davranışlarını stratejik bakış açısıyla değerlendirerek daha verimli ders programları oluşturulmasına katkı sunmayı amaçlamaktadır.

Anahtar Kelimeler: Oyun teorisi, seçmeli ders, Nash dengesi, sıfır toplamli oyun, sıfır toplamli olmayan oyun, stratejik karar.

Teşekkür Bölümü: Bu çalışma 2210-A Yurt İçi Genel Yüksek Lisans Burs Programı kapsamında TÜBİTAK Bilim İnsanı Destek Programları Başkanlığı tarafından desteklenmektedir.

Referanslar

- [1] Romp, G. (1997). Game Theory: Introduction and Applications. New York: Oxford University Pres.
- [2] Zagare, F. C. (1989). Game Theory, Concepts and Applications. Beverly Hills, California: Sage Publications, Fourth Printing, 11.
- [3] Tezcan, H., Gümüş, Y. (2008). *Üniversite Öğrencilerinin Seçmeli Ders Tercihlerine Etki Eden Faktörlerin Araştırılması*. Gazi Üniversitesi Gazi Eğitim Fakültesi Dergisi, 28(1), 1–17.
- [4] Aslım, G., Yaşar, A., Tekindal, M. A., Çelik, E. N. ve Yiğit, A. (2023). *Türkiye’de Veteriner Fakültesi’nde Seçmeli Ders Tercihlerine Etki Eden Faktörler*. Kafkas Üniversitesi Veteriner Fakültesi Dergisi.

Improved Fixed Point Approximations via Picard's Three-Step Iteration for Suzuki-Type Non-Expansive Mappings

Lale CONA¹, Suzan IŞIK²

¹*Faculty of Engineering and Natural Sciences, Mathematical Engineering,
Gümüşhane University, Türkiye*

²*Graduate Education Institute, Mathematics, Gümüşhane University, Türkiye*

suzankarabey@hotmail.com

In this study, we investigate the weak and strong convergence behavior of Picard's three-step iteration scheme applied to Suzuki generalized non-expansive mappings in the context of uniformly convex Banach spaces. By employing essential tools such as the Opial condition, the demiclosedness principle, and asymptotic center theory, we establish several convergence theorems. Our results demonstrate that under appropriate assumptions, the iterative sequence converges strongly and weakly to fixed points of enriched Suzuki-type non-expansive operators. The proposed iteration method enhances the approximation process in comparison to classical schemes, offering improved convergence performance. This work contributes to the ongoing development of fixed point theory and provides a more flexible and efficient framework for solving nonlinear problems in Banach spaces.

Keywords: Fixed point, Suzuki generalized non-expansive mapping, three-step Picard iteration, strong convergence, weak convergence.

References

- [1] Suzuki, T. (2008). *Fixed Point Theorems and Convergence Theorems for Some Generalized Non-expansive Mappings*. J. Math. Anal. Appl., 340, 1088-1095.
- [2] Ali, F., Ali, J., and Rodríguez-Lopez, R. (2022). *Approximation of Fixed Points and the Solution of a Nonlinear Integral Equation*. Nonlinear Functional Analysis and Applications, 26(5), 869-885.
- [3] Opial Z. (1967). *Weak Convergence of the Sequence of Successive Approximations for Nonexpansive Mappings*. Bulletin of the American Mathematical Society, 73(4), 591-597.

Investigation of Fractal Dimension Algorithms in Different Geometries

İpek Ebru KARAÇAY¹, Salim YÜCE²

¹Computer Programming, Istanbul Gelisim Vocational School, Istanbul Gelisim University, Avcılar,
Istanbul, Türkiye

²Department of Mathematics, Faculty of Arts and Sciences, Yıldız Technical University, Esenler,
Istanbul, Türkiye

iekaracay@gelisim.edu.tr

Fractal geometry is a sub-branch of mathematics used to mathematically describe the complex and irregular structures observed in nature. The concept of fractal dimension, also referred to as a complexity measure, is employed to quantify these intricate structures. This measure, which goes beyond classical Euclidean geometry, reveals the degree of complexity underlying phenomena such as biological signals, natural patterns, financial time series, and neurological data. Owing to its non-integer, fractional nature, the fractal dimension provides a powerful tool for characterizing the geometry of nature. Numerous algorithms have been developed to calculate fractal dimensions. In particular, the Katz and Higuchi algorithms stand out for their high precision in analyzing time series data. The Katz algorithm defines complexity as the ratio between the total length of a signal and its maximum deviation, whereas the Higuchi algorithm estimates dimensionality using a log-log regression across multiple subsequences. Both of these algorithms are widely used in the analysis of biological data, especially electroencephalographic (EEG) signals.

This paper aims to investigate these two fractal dimension algorithms within the framework of Galilean and Lorentzian (Minkowski) space-time geometries, in order to explore the physical interpretation of dimensional complexity. By embedding fractal analysis within these distinct geometric models, the study seeks to offer a novel perspective on how space-time assumptions can influence the calculation and interpretation of fractal dimensions.

Keywords: Fractal, Katz fractal dimension, Higuchi fractal dimension, Minkowski plane, Galilean plane.

References

- [1] Mandelbrot, B. B. (1983). *The Fractal Geometry of Nature*. Henry Holt and Company ISBN:0716711869.
- [2] Higuchi, T. (1988). *Approach to an Irregular Time Series on the Basis of the Fractal Theory*. Physica D: Nonlinear Phenomena 31(2), 277-283.
- [3] Katz, M. J. (1988). *Fractals and the Analysis of Waveforms*. Computers in biology and medicine, 18(3), 145-156.

Machine Learning-Based Classification of Chaotic Behavior in Discrete Dynamical Systems

Radva ANSARY, Derya SEKMAN

*Department of Mathematical Engineering, Faculty of Chemistry and Metallurgy, Yıldız Technical
University, 34220, Istanbul, Türkiye*

radva.ali@std.yildiz.edu.tr

Discrete dynamical systems that exhibit chaotic behavior form the foundation of many natural and engineering systems due to their sensitivity to initial conditions and unpredictability. This study aims to classify the parameter-dependent dynamical behaviors of one-dimensional discrete chaotic systems using machine learning methods. The study focuses on distinguishing chaotic and non-chaotic regimes of various discrete dynamical system models through the application of machine learning algorithms.

Keywords: Discrete dynamical systems, chaotic behavior, machine learning, classification.

References

- [1] Devaney, R. L. (2020). A First Course in Chaotic Dynamical Systems: Theory and Experiment. CRC Press, Boca Raton.
- [2] Uzun, S. (2022). *Machine Learning-Based Classification of Time Series of Chaotic Systems*. European Physical Journal Special Topics, 231, 493–503.
- [3] Akmeşe, Ö. F., Emin, B., Alaca, Y., Karaca, Y., and Akgül, A. (2024). *The Time Series Classification of Discrete-Time Chaotic Systems Using Deep Learning Approaches*. Mathematics, 12(19), 3052.

Variation of Parameters and Hölder Stability of Differential Systems with Initial Time Difference

Coşkun YAKAR¹, Dilara KARSLIOĞLU²

¹*Department of Mathematics, Gebze Technical University, Kocaeli, Türkiye*

²*Department of Mathematics, Yeditepe University, Istanbul, Türkiye*

dilara.karslioglu@yeditepe.edu.tr

In this paper, we have investigated the relation between an unperturbed differential equations and a perturbed differential system that have different initial positions and an initial time difference by using the variation-of-parameter techniques employed to obtain integral formulae. To establish Hölder stability with initial time difference for nonlinear differential systems, the variational system associated with the unperturbed differential system must be used.

Keywords: Differential equations, initial time difference, Hölder stability, variation of parameters.

References

- [1] Dannan, F. M., Elaydi, S. (1986). *Lipschitz Stability of Nonlinear Systems of Differential Equations*. Journal of Mathematical Analysis and Applications, 113(2), 562-577.
- [2] Yakar, C., Deo, S. G. (2006). *Variation of Parameters Formulae with Initial Time Difference for Linear Integrodifferential Equations*. Applicable Analysis, 85(4), 333-343.
- [3] Yakar, C., Shaw, M. D. (2008). *Initial Time Difference Stability in Terms of Two Measures and a Variational Comparison Result, Dynamics of Continuous. Discrete & Impulsive Systems A. Mathematical Analysis*, 15(3), 417-425.

W5-Curvature Tensor on a Complex Contact Space Form

Aysel TURGUT VANLI, Gamze ALKAYA

Department of Mathematics, Gazi University, Ankara, Türkiye

gamzealkaya90@gmail.com

This article aims to examine the curvature properties of the W5-curvature tensor on a complex contact space form. We examine a complex contact space form under the W5-curvature tensor's W5-flat, ξ -W5-flat, GH-W5-flat, GH-W5-semi-symmetric, W5Q and W5(ξ , K)S conditions. We also demonstrate that there is no a complex contact space form for W5-flat, ξ -W5-flat, GH-W5-flat, GH-W5-semi-symmetric, W5Q and W5(ξ , K)S.

Keywords: Complex contact manifold, GH-sectional curvature, complex contact space form, W5-curvature.

References

- [1] Blair, D. E., Molina, V. M. (2011). *Bochner and Conformal Flatness on Normal Complex Contact Metric Manifolds*. Ann. Global Anal. Geom., 39(3), 249–258.
- [2] Turgut Vanli, A., Unal, I. (2017). *Conformal, Concircular, Quasi-conformal and Conharmonic Flatness on Normal Complex Contact Metric Manifolds*. International Journal of Geometric Methods in Modern Physics, 14(5), 1750067.

Schauder Estimates for Elliptic Operators in Banach-Sobolev Spaces

Bilal T. BİLALOV^{1,2}, Eminaga MAMEDOV², and Yonca SEZER¹

¹*Department of Mathematics, Yıldız Technical University, Davutpaşa, 34220 İstanbul, Türkiye*

²*Department of Algebra and Mathematical Logic, Institute of Mathematics and Mechanics, The*

Ministry of Science and Education of the Republic of Azerbaijan, Baku, Azerbaijan

ysezer@yildiz.edu.tr

In this work, we establish a Schauder-type estimate up to the boundary for elliptic operators with continuous coefficients on a finite-dimensional bounded domain, within a non-separable Banach-Sobolev space generated by a rearrangement-invariant Banach function space. This paper extends the results¹⁻³. Our approach uses an abstract functional-analytic framework—combining interior estimates, extendibility properties of functions, and the theory of elliptic operators—instead of classical function-theoretic methods.

Keywords: Banach function space, rearrangement-invariant space, Banach-Sobolev space, elliptic operator, Schauder-type estimate.

References

- [1] Mamedov, E. M., Cetin, S. (2024). *Interior Schauder-type Estimates for m -th Order Elliptic Operators in Rearrangement-Invariant Sobolev Space*. Turkish Journal Mathematics. 48(4), 793–816.
- [2] Bilalov, B. T., Sadigova, S. R. (2021). *On the Fredholmness of the Dirichlet Problem for a Second-order Elliptic Equation in Grand-Sobolev Spaces*. Ricerche di Matematica. doi:org/10.1007/s11587-021-00599-9.
- [3] Bilalov, B. T., Sadigova, S. R. (2021). *Interior Schauder-type Estimates for Higher-Order Elliptic Operators in Grand-Sobolev Spaces*. Sahand Communications in Mathematical Analysis. 1(18-2), 129-148.

Identities on the Generalized Harmonic Numbers of Rank with Order Using Generating Functions

Ömer DURAN¹, Neşe ÖMÜR¹, and Sibel KOPARAL²

¹ Department of Mathematics, Kocaeli University, 41380, İzmit, Kocaeli, Türkiye

² Department of Mathematics, Bursa Uludağ University, 16059, Bursa, Türkiye

omer20841@gmail.com

In this study, we investigate new identities involving the generalized harmonic numbers of rank r with order $m = (m_0, m_1, \dots, m_r) \in (\mathbb{Z}^+)^{r+1}$. By employing generating function techniques, we derive several summation formulas, recurrence relations, and connections with classical special numbers such as Stirling numbers and r -derangement numbers. Our approach generalizes existing results and provides a unified framework that links classical special numbers with harmonic numbers and generalizations. These findings contribute to the combinatorial and analytical understanding of generalized harmonic numbers and their applications in number theory.

Keywords: Generalized harmonic numbers, generating functions, combinatorial identities.

References

- [1] Benjamin, A. T., Gaebler, D., and Gaebler, R. (2003). *A Combinatorial Approach to Hyperharmonic Numbers*. *Integers: Electron. J. Combin. Number Theory*, 3, 1-9.
- [2] Duran, Ö., Ömür, N., and Koparal, S. (2020). *On Sums with Generalized Harmonic, Hyperharmonic and Special Numbers*. *Miskolc Mathematical Notes*, 21(2), 791-803.
- [3] Duran, Ö., Ömür, N., and Koparal, S. (in submission). *Generalized Harmonic Numbers of Rank r with Order $m = (m_0, m_1, \dots, m_r)$ and Their Applications*.
- [4] Gertsch, A. (1997). *Generalized Harmonic Numbers*. *Journal of Number Theory*, 324, 7-10.
- [5] Ömür, N., Koparal, S. (2022). *Sums Involving Generalized Harmonic and Daehee Numbers*. *Notes on Number Theory and Discrete Mathematics*, 28(1), 92-99.
- [6] Shapiro, L. W., Getu, S., Woan, W. J., and Woodson, L. C. (1991). *The Riordan Group*. *Discrete Applied Mathematics*, 34(1-3), 229-239.
- [7] Wang, C., Miska, P., and Mező, I. (2017). *The r -derangement Numbers*. *Discrete Mathematics*, 340, 1681-1692.
- [8] Wang, Y., Xin, G. (2019). *Hankel Determinants for Convolution Powers of Catalan Numbers*. *Discrete Mathematics*, 342, 2694-2716.

Solitary Wave Solutions of the Two-Dimensional Fractional-Order Nonlinear Schrödinger Equation

Günay ASLAN

Istanbul Kültür University, 34158, Istanbul, Türkiye

g.aslan@iku.edu.tr

In this study, solitary wave solutions of the two-dimensional fractional-order nonlinear Schrödinger (2DfNLS) equation are investigated using the Petviashvili method. Both single and double soliton profiles are constructed by numerically solving the 2DfNLS equation, which takes the following form:

$$iu_t - \beta(-\nabla)^s u - V(\mathbf{x})u - \gamma|u|^{2\sigma}u = 0, \quad \mathbf{x} = (x, y), \quad t > 0.$$

Here, $u = u(x, y, t)$ denotes the wave function, $\nabla = \partial_{xx} + \partial_{yy}$ is the two dimensional Laplacian operator, and $V(\mathbf{x})$ represents the potential function. In this equation, the nonlinearity is characterized by $\sigma > 0$, with $\sigma = 1$ corresponding to the cubic case. The fractional operator $(-\nabla)^s$, where $0 < s < 1$, introduces nonlocal effects into the model and generalizes the classical nonlinear Schrödinger equation (recovered when $s = 1$), a well-known model for the propagation of weakly nonlinear waves in dispersive media. The equation may describe either a focusing or defocusing regime, depending on the signs of the parameters β and γ .

The aim of this study is to provide a comprehensive analysis of soliton solutions for the 2DfNLS equation. The Petviashvili method, employed in this work, emerges as an effective numerical scheme for computing nonlinear stationary solitons, particularly useful for analyzing stable localized structures. The method was originally introduced for solitary wave solutions of nonlinear wave equations in plasma physics¹, and its convergence properties and optimal parameter conditions were studied in detail². The findings offer valuable insights into the dynamics of soliton-like wave phenomena in various physical systems.

Keywords: Fractional Schrödinger equation, soliton dynamics, Petviashvili method.

References

- [1] Petviashvili, V. I. (1976). *Equation of an Extraordinary Soliton*. Plasma Physics, 2, 469– 472.
- [2] Pelinovsky, D. E., and Stepanyants, Y. A. (2004). *Convergence of Petviashvili's Iteration Method for Numerical Approximation of Stationary Solutions of Nonlinear Wave Equations*. SIAM Journal on Numerical Analysis, 42, 1110– 1127.

Boundedness Criteria for the G -fractional Integral and G -fractional Maximal Operator on Gegenbauer-morrey Spaces

Elman I.İBRAHİMOV

Ministry of Science and Education Republic of Azerbaijan, Institute of Mathematics and Mechanics

elmanibrahimov@yahoo.com

In my report we establish necessary and sufficient conditions for the boundedness of the Gegenbauer fractional (G -fractional) integral on Gegenbauer-Morrey (G -Morrey) spaces associated with the Gegenbauer differential operator. A similar problem is studied for the G -maximal function. In the paper, we prove a Adams type theorem on the boundedness of the Gegenbauer fractional integral J_G^α and maximal function generated by Gegenbauer differential operator

$$G \equiv G_\lambda \equiv (x^2 - 1)^{\frac{1}{2}-\lambda} \frac{d}{dx} (x^2 - 1)^{\lambda+\frac{1}{2}} \frac{d}{dx}, \quad x \in (1, \infty), \quad \lambda \in \left(0, \frac{1}{2}\right),$$

introduced in². The result obtained is an analog of the corresponding theorem obtained for Riesz potentials in¹.

Keywords: G -fractional integral, G -maximal operator, G -Morrey spaces.

References

- [1] Adams D. R. (1975). *A note on Riesz Potentials*. Duke Math., 42, 765-778.
- [2] Durand L., Fishbane P. M., and Simmons L. M. (1976). *Expansion Formulas and Addition Theorems for Gegenbauer Functions*. J. Math. Phys, 17(11), 1993-1948.

Hemen Hemen Neo Kobalans Sayıları

B. Ecem AKGÜÇ CİMBİZ, Ahmet TEKCAN

Bursa Uludağ Üniversitesi, Fen Edebiyat Fakültesi, Matematik Bölümü, Bursa, Türkiye

ecemakg2802@gmail.com

Bu çalışmada hemen hemen neo kobalans sayıları tanımlanmış ve bu sayıların genel terimleri, balans sayılarına bağlı olarak elde edilmiştir. Daha sonra bu sayıların Pell, Pell-Lucas, üçgensel ve kare üçgensel sayılar ile olan ilişkisi ortaya çıkartılmıştır. Son olarak bu sayıların ilk n -terim toplamaları elde edilmiştir.

Keywords: Balans sayıları, hemen hemen balans sayıları, Pell sayıları, üçgensel sayılar.

References

- [1] Barbeau, E. J. (2003). Pell's Equation. Springer-Verlag, New York.
- [2] Behera, A., Panda, G. K. (1999). *On the Square Roots of Triangular Numbers*. The Fibonacci Quarterly, 37(2), 98-105.
- [3] Chailangka, N., Pakapongpun, A. (2021). *Neo Balancing Numbers*. Int. Jour. of Math. and Comp. Sci., 16(4), 1653-1664.
- [4] Liptai, K., Luca, F., Pinter, A., and Szalay, L. (2009). *Generalized Balancing Numbers*. Indagationes Math., 20(1), 87-100.
- [5] Panda, G. K., Ray, P. K. (2009). *Some Links of Balancing and Cobalancing Numbers with Pell and Associated Pell Numbers*. Bulletin of the Inst. of Maths, Academia Sinica, 6(1)(2011), 41-72.
- [6] Ray, P. K. (2009). *Balancing and Cobalancing Numbers*. Ph.D. thesis, Department of Mathematics, National Institute of Technology, Rourkela, India.
- [7] Panda, G. K., Panda, A. K. (2015). *Almost Balancing Numbers*. Jour. of the Indian Math. Soc., 82(3-4), 147-156.
- [8] Panda, A. K. (2017). *Some Variants of the Balancing Sequences*. Ph.D. thesis, National Institute of Technology Rourkela, India.
- [9] Tekcan, A., Akgüç, E. (2025). *Almost Neo Cobalancing Numbers*. Notes on Number Theory and Dis. Maths., 31(1), 113-126.

Δ -Convergence For Proximal Point Algorithm And Fixed Point Problem in Generalized $CAT(0)$ Space

Muhammet KNEFATI¹, Vatan KARAKAYA²

¹*Department of Mathematics, Yildiz Technical University, Istanbul, Türkiye*

²*Department of Mathematical Engineering, Yildiz Technical University, Istanbul, Türkiye*
mknefati@gmail.com

This paper investigates generalized $CAT(0)$ spaces by introducing their definition and examining key structural properties. A generalized CN^* inequality is formulated within this framework, and it is subsequently employed to demonstrate the strong convergence and Δ –convergence of the Karakaya-iteration process towards fixed points of nonexpansive mappings. Building upon, the Karakaya -iteration method is further applied to address a minimization problem, with particular emphasis. Two numerical examples are presented: the first within a $CAT_p(0)$ space that does not satisfy the $CAT(0)$ condition, and the second within a $CAT(0)$ space that is not a Hilbert space.

Keywords: $CAT(0)$ spaces, proximal point algorithm , Δ –convergence.

References

- [1] Bridson, M. R., Haefliger, A. (2013). *Metric Spaces of Non-positive Curvature*, 319. Springer Science & Business Media, Berlin.
- [2] Khamsi, M., Shukri, S. (2017). *Generalized $CAT(0)$ Spaces*. Bulletin Belgian Math. Soc.-Simon Stevin, 24(3), 417–426.
- [3] Kirk, W., Panyanak, B. (2008). *A Concept of Convergence in Geodesic Spaces*. Nonlinear Anal.: Theory, Methods & Appl., 68(12), 3689–3696.

Impulse-Hydrodynamic Model of Explosive Impact on a Half-Plane with Circular Cavity

Nail TUKTAMYSHOV

Russian Federation, Kazan, Kazan State University of Architecture and Civil Engineering

nail1954@gmail.com

This study considers the explosion of a surface cord charge with variable cross-section in a half-plane containing a circular cavity. The impulse hydrodynamic model [1,2] is used, where the medium is treated as an ideal incompressible fluid. The main objective is to determine the width of the ejection cavity, where the particle velocity at the edge reaches the critical value v_0 .

The free surface is assumed to coincide with the x -axis, and the thickness of the charge is defined by the function $y = f(x)$. The pressure is proportional to the charge thickness: $P = kf$, where k is a known constant. The potential along the x -axis is given by $\varphi = -kf(x)/\rho$, where ρ is the density of the medium. The complex potential is introduced as $w(z) = \varphi(x, y) + i\psi(x, y)$. The goal is to find the function $w(z)$ in the lower half-plane $\text{Im} z < 0$ with a circular cavity, satisfying the boundary conditions: $\varphi = -kf(x)/\rho$ for $x \in (-\infty, \infty)$ and $\varphi = 0$ on the circle $z - ih = a_0 \exp(i\theta)$ ($0 \leq \theta < 2\pi$), a_0 is the radius of the cavity, and h is the depth of its center. By conformal mapping $z_1 = z_1(z)$, the domain is transformed into an annular region. The function $w(z_1)$, analytic in the ring, is represented as $w(z_1) = \tilde{A} \ln z_1 + \tilde{w}(z_1)$, where $\tilde{A} = -\frac{1}{\ln q} \frac{1}{2\pi} \int_0^{2\pi} (u_0(\gamma) - u_1(\gamma)) d\gamma$, $u_0(\gamma) = \text{Re} w(e^{i\gamma})$, $u_1(\gamma) = \text{Re} w(qe^{i\gamma})$. The function $\tilde{w}(z_1)$ is defined by the Villat formula [3]: $\tilde{w}(z_1) = \frac{i\omega_1}{\pi^2} \left(\int_0^{2\pi} u_0(\gamma) \zeta\left(\frac{\omega_1}{\pi i} \ln z_1 - \frac{\omega_1}{\pi} \gamma\right) d\gamma - \int_0^{2\pi} (u_1(\gamma) - \tilde{A} \ln q) \zeta_3\left(\frac{\omega_1}{\pi i} \ln z_1 - \frac{\omega_1}{\pi} \gamma\right) d\gamma \right) + i\beta_0$, where $\zeta(z_1)$ and $\zeta_3(z_1)$ are the Weierstrass zeta functions, β_0 is an arbitrary real constant, and $q = \exp(-\pi\omega_3 / i\omega_1)$.

The resulting expression for the complex potential allows determining the geometry of the ejection cavity and the velocity distribution in the medium and on the cavity boundary. Numerical results are of practical importance.

Keywords: Explosion, pressure impulse, potential.

References

- [1] Vlasov, O. E. (1957). Fundamentals of Explosion Dynamics. Moscow: VIA, 377.
- [2] Kuznetsov, V. M. (1977). Mathematical Models in Blasting. Novosibirsk: Nauka, 260.
- [3] Akhiezer, N. I. (1970). Elements of the Theory of Elliptic Functions. Moscow: Nauka, 304.

Model and Theorem of Existence of the Problem of Optimal Control of the Customs Clearance Process

Saidov ABDUSOBİR¹, Saidov IMRON², and Abdurakhmonov TOHİR³

¹*Tashkent University of Information Technologies named after Muhammad al-Khwarizmi,
Tashkent, Uzbekistan*

²*Yildiz Technical University, Istanbul, Türkiye*

³*Institute of Mathematics named after V. Romanovsky of the Academy of Sciences of
the Republic of Uzbekistan*

abdusobir59@gmail.com

The article examines the problem of modeling the multi-stage process of customs clearance of foreign trade goods. A mathematical model for controlling the considered process has been developed. The essence of customs risks is presented and the theorem on the existence of a solution to the problem of optimal control of the customs clearance process is proven.

The problem of optimal control of the customs clearance process is considered as a linear programming problem with variable coefficients of the objective function¹. An analysis of scientific papers on the study of similar problems showed that a sufficient number of studies have been conducted and certain methods have been developed for solving the problem of linear programming with variable coefficients. For example, in the work (Pinsker A.G.) the problem of linear programming with variable coefficients was investigated by introducing the concept of resolving combinations of coefficients of the objective function². Based on the results of this study, within the framework of the present research, the following theorem is proven:

Theorem. The task of optimal management of the customs clearance process with non-empty set of plans and a targeted function of customs risks limited from above has a solution.

Keywords: Customs clearance, mathematical modeling, customs risks.

References

- [1] Mukhtorov, I., Abdurakhmonov, T., and Saidov, A. (2024) *Mathematical Model of the Customs Clearance Process Taken into Account of Customs Risks*. Uzbek Mathematical Journal, 2024, 68(4), 107-115. doi: 10.29229/uzmj.2024-4-12.
- [2] Pinsker, A. (1979). *A Linear Programming Problem with Variable Coefficients of the Purpose Function*. Sib Math J 20, 466–468. doi.org/10.1007/BF00969958.
- [3] Van der Waerden, B. L. (1976). *Hamilton's Discovery of Quaternions*. Math. Mag., 49(5), 227-234.

Necessary Conditions for Weak and Strong Local Minimum the Problem of the Calculus of Variations with Higher Order Derivatives

Narmina GUBATOVA

*Azerbaijan State Oil and Industry University(ASOIU), 16/21 Azadliq Ave, Baku
French-Azerbaijani University(UFAZ), Nizami 183, Baku
Institute of Mathematics and Mechanics, B. Vahabzade 9, Baku*

narmina2606@gmail.com

We consider the following problem of the calculus of variations with higher order derivatives in space $C^n([t_0; t_1])$ and $PC^n([t_0; t_1])$

$$J(x) = \int_{t_0}^{t_1} L(t, x, \dot{x}, \ddot{x}, \dots, x^{(n)}) dt \rightarrow \min, \quad (P)$$

$$x^{(k)}(t_0) = x_{k0}, \quad x^{(k)}(t_1) = x_{k1}, \quad k = 0, \dots, n-1.$$

Theorem 1. Let $\hat{x} \in PC^n([t_0; t_1])$ provides a strong local minimum in the problem (P), with the integrant L being n times continuously differentiable in some neighborhood of the extended graph . Then, $\hat{x}$ satisfies the Euler–Poisson equation and the Weierstrass condition. Moreover, if the second derivative $L_{x^{(n)}x^{(n)}}$ exists, then the Legendre condition holds.

Theorem 2. Let $\hat{x} \in C^{2n}([t_0; t_1])$ provides a weak local minimum in the problem (P), with the integrant L being $n+2$ times continuously differentiable in some neighborhood of the extended graph. Then, $\hat{x}$ satisfies the Euler–Poisson equation and Legendre condition. If the strong Legendre condition holds then Jacobi conditions also holds.

Keywords: Calculus of variations, problem with higher order derivatives.

References

- [1] Alekseev, V., Galeev, E., and Tikhomirov, V. (2011). Collection of Optimization Problems, FIZMATLIT.
- [2] Galeev, E. (2016). Methods of Optimization. Baku branch of Moscow State University.
- [3] Galeev E., Tikhomirov V. (2000). Optimization. Theory. Examples. Problems, Editorial URSS.

Numerical Analysis of Stiffened Shell Structures Using an Efficient Three-Dimensional Finite Element

Djamal HAMADI, Maria LEGOUÏRAH, and Abdulrahman AL-NADHARI

LARGHYDE Laboratory, Civil Engineering and Hydraulics Department, Biskra University,

B. P 145.RP.07000, Biskra, Algeria

d.hamadi@univ-biskra.dz

Shell structures are one of the most distinctive engineering constructions in architectural design due to their efficiency in supporting loads through their curved geometries and providing a huge free surface without columns. These structures are commonly used in large-scale architectural structures such as sports stadiums and airports, where they provide a balance between technical aesthetics and practical functions¹. Usually, they are subjected to important external loads with a huge covered surface. These cause large deflection of the structure and threaten its structural stability. To overcome these inconveniences, the shell can be reinforced with stiffeners to improve its structural stability. Unfortunately, with this type of complex structure in geometrical shape and boundary conditions, the analytical methods can not be used to check the stability, in addition to calculating the internal forces. Hence; numerical methods will be involved to solve the problem. In this paper, the numerical analysis is carried out with ABAQUS software, using a three-dimensional finite element "C3D8IH" for both the shell and the stiffener to overcome the incompatibility of degrees of freedom encountered when using two different elements². The applications are carried out on stiffened shell structures, and the results obtained showed that the stiffeners significantly improve the stability of the structure and the efficiency of the element used is confirmed. This approach and the results obtained provide innovative design recommendations to improve the safety and rigidity of structures that cover huge surfaces.

Keywords: Numerical analysis, stiffened shell, three-dimensional element, complexe geometrical shape.

References

- [1] Legouirah, M., Hamadi, D., and Al-Nadhari, A. M. (2024). *The Efficiency of Ring Stiffener Shape on the Deformation of Cylindrical Shell Structures–Numerical Analysis with Solid Finite Element*. Asian Journal of Civil Engineering, 1-10. <https://doi.org/10.1007/s42107-024-01134-5>
- [2] Karpov, V. V., Semenov, A. A. (2021). *Structural Anisotropy Method for Shells with Orthogonal Stiffeners*. Elsevier, 34, 3206-3221. <https://doi.org/10.1016/j.istruc.2021.09.027>

Development of a General Algorithm for Solving Stability Problems of Plates on Elastic Bases

Shahin GULİYEV, Rizvan SHUKUROV

Azerbaijan Technological University, Ganja, Azerbaijan

sh.guliyev@atu.edu.az

The development of a general algorithm for solving stability problems of anisotropic plates is devoted¹. The basics of the method of discretization of additional load are considered in the article². The proposed discrete method for solving the problem of stability of elastic systems has a certain area of application, the main indicators of which follow from the fundamental characteristics of the method. These include, first of all, the necessary initial information. It is in this method is an additional load. To obtain it, it is necessary to have differential equations describing the deformation of the system in the given and adjacent states. Since such a condition almost does not meet any restrictions when setting the stability in the "small", the scope of application of the additive load method is very extensive. It includes not only simple elastic systems made of individual elements, but also complex systems formed by the articulation of elements. It is important to note that only differential equations are needed, not their solutions.

To determine the Green's function in complex cases, the variational-difference method is used. They solve a wide class of plane and spatial problems. In this case, the Green's function by its nature, as related to the unit load, is an invariant part of the calculation, it does not depend on the load of the system and is preserved at any given loads. Varying the grid of points in determining the Green's function, we have revealed a regularity, the use of which greatly facilitates computational operations. Thus, the proposed method of solving stability problems of elastic systems allows solving stability problems of plates on an elastic base of systems by the same general algorithm, has a wide enough area of application and significant advantage over existing methods.

Keywords: Stability, plates, elastic foundation, Green's function.

References

- [1] Guliyev, Sh. M., Shukurov, R. E., Huseynzade, H. S., Huseynova, L. N., and Hasanov, A. B. (2024). *Development of a General Algorithm for Solving the Stability Problem of Anisotropic Plates*. Eastern-European Journal of Enterprise Technologies, 128(7), 16-23.
- [2] Guliyev, Sh. M. (2021). *Resheniye Nekotoryx Zadach Ustoychivosti Uprugix Sistem Po Metodu Diskretizatsii Dobavochnoy Nagruzki*. Materialy mejdunarodnoy nauchno-prakticheskoy konferentsii "Osnovnyye problemy reytinga vuzov", Ganja, Azerbaijan, 2, 12-13.

Modelling Electromagnetic Wave Propagation Through Time-Varying Medium Using the Finite Difference Time Domain Method

Deniz GÜNDOĞAN, Sebahattin EKER

Informatics Institute, Istanbul Technical University, Istanbul, Türkiye

erso22@itu.edu.tr

Electromagnetic wave propagation in time-varying medium using the Finite Difference Time Domain method is presented. Electromagnetic wave propagation in space varying medium like plasma slab was studied earlier. Sudden created plasma medium is chosen for the time varying medium. Permittivity of the medium is changing with time during the wave propagation. In order to compute the electric and magnetic field values we used the Maxwell Equation's. Finite Difference method used for the analysis of Maxwell Equation's. Maxwell Equation's in time domain are partial differential equations. Finite difference time domain method properly formulated and discretized the Maxwell Equation's in time and space to lead the solution step-by step.

Keywords: Time-varying medium, wave propagation, finite difference time domain method.

References

- [1] Taflove, A. (1995). Computational Electrodynamics: The Finite Difference Time Domain Method. Artech House Inc. , Norwood, MA, USA, 1995.
- [2] Yee, K. (1966). Numerical Solution of Initial Boundary Value Problems Involving Maxwell Equations in Isotropic Media. IEEE Trans Antennas Propag., 14(3), 302-307.
- [3] Kalluri, D. K. (2010). Electromagnetics of Time Varying Complex Media. Second edition, CRC press, LLC, Boca Raton, FL, USA.

Numerical Modeling of Stars and Mathematical Analysis

Handan ÇELİK, Derya SEKMAN

*Department of Mathematical Engineering, Faculty of Chemistry and Metallurgy, Yıldız Technical
University, 34220 Istanbul, Türkiye*

handancelik615@gmail.com

In this study, which aims to demonstrate the relationship between stars and mathematics, the mathematical background of three formulas related to stars has been examined. The goal of this research is to show how consistent the values obtained from the examined formulas are with the GaiaDR3 data. The formulas investigated within the scope of the study are the Stefan-Boltzmann Law, the mass-luminosity relation, and the absolute magnitude equations. These theoretically examined relations were applied to both directly measured and model-derived values in the GaiaDR3 data set to evaluate their success. In this context, the mathematical derivations of the Stefan-Boltzmann Law and the absolute magnitude formulas were demonstrated, and the mass-luminosity relation was applied using regression to illustrate its connection with mathematics. The results obtained from the implementation carried out in the Python environment were visualized through diagrams and various graphs, providing a comprehensive understanding of the subject. As a result of the applications performed, it was observed that the values obtained from the three formulas generally show consistency with the GaiaDR3 dataset. However, some discrepancies were also noted in certain formulas, indicating areas for further investigation and refinement.

Keywords: Stefan-Boltzmann law, absolute magnitude, mass-luminosity relation, stars, numerical modeling.

References

- [1] Crepeau, J. (2009). *A Brief History of the T₄ Radiation Law*. Proceedings of the ASME 2009 Heat Transfer Summer Conference collocated with the InterPACK09 and 3rd Energy Sustainability Conferences, San Francisco, California, USA.
- [2] Wang, J., Zhong, Z. (2018). *Revisiting the Mass-luminosity Relation with an Effective Temperature Modifier*. Astronomy & Astrophysics, 619, DA.
- [3] LoPresto, M. C., Hagoort, N. (2011). *Determining Planetary Temperatures with the Stefan-Boltzmann Law*. The Physics Teacher, 49, 113–116.

New Midpoint and Trapezoid-Type Inequalities for Co-ordinated m -Convex Functions via Variable-Order Fractional Calculus

Abdüllatif YALÇIN¹, Erdal GÜL², and Ahmet Ocak AKDEMİR³

^{1,2}*Yıldız Technical University, Faculty of Arts and Science Department of Mathematics,
Istanbul, Türkiye*

³*Ağrı İbrahim Çeçen University, Faculty of Science and Letters, Department of Mathematics,
Ağrı, Türkiye*

abdullatif.yalcin@std.yildiz.edu.tr

This paper presents novel midpoint and trapezoid-type inequalities for co-ordinated m -convex functions by employing variable-order Riemann–Liouville fractional integrals. Building on classical results for convex mappings, we extend these foundational inequalities to a more general class of m -convex functions defined on bidimensional rectangular domains, thereby providing tighter and more flexible bounds. The analysis utilizes the adaptability of variable-order fractional operators to achieve more refined results compared to constant-order fractional integrals. Specifically, the proposed inequalities generalize existing findings in the literature, encompassing both classical Riemann integrals and constant-order fractional integrals as special cases. We further demonstrate that the derived inequalities reduce to known results under appropriate parameter choices, highlighting their versatility. These contributions advance the theoretical landscape of fractional inequalities and are expected to have significant applications in numerical approximation, integral equations, and optimization methods, particularly in contexts where generalized convexity plays a crucial role.

Keywords: Variable-Order fractional integrals, co-ordinated convex mapping, midpoint-type inequalities, trapezoid-type inequalities, co-ordinated m -convex functions.

Acknowledgement: This study is produced from the findings of the Ph. D. thesis of first author.

References

- [1] Samko, S. G., Ross, B. (1993). *Integration and Differentiation to a Variable Fractional Order*. Integral transforms and special functions, 1(4), 277-300.
- [2] Özdemir, M. E., Set, E. (2011). *Some New Hadamard Type Inequalities for Co-Ordinated m -convex and (α, m) -convex Functions*. Hacettepe journal of mathematics and statistics, 40(2), 219-229.

A Study on Dual Quaternions with k -Pell and k -Pell-Lucas Coefficients

Çiğdem Zeynep YILMAZ, Gülsüm Yeliz SAÇLI

*Department of Mathematics, Faculty of Arts and Sciences, Yildiz Technical University, 34220, Istanbul,
Türkiye*

cigdem.yilmaz@yildiz.edu.tr

Generalized forms of positive integer sequences for any positive integer k have been the subject of extensive research^{1,2}. In 2013, Catarino defined k -Pell numbers. In this article, the Binet's formula, the generating function, some identities and properties of k -Pell numbers are examined³. In a different work, Catarino and Vasco conducted a study involving k -Pell numbers⁴. In 2013, definition of the k -Pell-Lucas sequence is given by Catarino and Vasco, and some properties, the generating function of k -Pell-Lucas numbers are provided⁵. In 2006, Majernik defined dual quaternions⁶. In 2011, Ercan and Yüce presented some properties of dual quaternions⁷. In 2016, Aydın and Yüce combined Pell numbers with dual quaternions⁸. Following a similar approach, we define new algebraic structures called k -Pell and k -Pell-Lucas dual quaternions. Subsequently, we examine their Binet formulas, generating functions, and various identities and properties associated with these sequences.

Keywords: k -Pell number, k -Pell-Lucas number, dual-quaternion.

References

- [1] Falcon, S., Plaza A. (2007). *On the Fibonacci k -numbers*. Chaos Solitons Fractals, 32(5), 1615-1624.
- [2] Yazlik, Y., Taskara, N. (2012). *A Note on Generalized k -Horadam Sequence*. Comput. Math. Appl., 63(1), 36-41.
- [3] Catarino, P. (2013). *On Some Identities and Generating Functions for k -Pell Numbers*. Int. J. Math. Anal., 7(38), 1877-1884.
- [4] Catarino, P., Vasco, P. (2013). *Some Basic Properties and a Two-by-two Matrix Involving the k -Pell Numbers*. Int. J. Math. Anal., 7(45), 2209-2215.
- [5] Catarino, P., Vasco, P. (2013). *On Some Identities and Generating Functions for k -Pell-Lucas Sequence*. Appl. Math. Sci., 7(98), 4867-4873.
- [6] Majernik, V. (2006). *Quaternion Formulation of the Galilean Space-time Transformation*. Acta Physica Slovaca, 56(1), 9-14.
- [7] Ercan, Z., Yüce, S. (2011). *On Properties of the Dual Quaternions*. European Journal of Pure Applied Mathematics, 4(2), 142-146.
- [8] Torunbalcı Aydın, F., Yüce, S. (2016). *Dual Pell Quaternions*. Journal of Ultra Scientist of Physical Sciences, 28(7), 328-339.

A Study on Vietoris' Hybrinomials

Nurten GÜRSES¹, Duygu ÇAĞLAR ÇAY²

¹*Department of Mathematics, Faculty of Arts and Sciences, Yıldız Technical University,
Istanbul, Türkiye*

²*Department of Software Engineering, Faculty of Engineering and Architecture,
Istanbul Gelisim University, Istanbul, Türkiye
dcaglar@gelisim.edu.tr*

Number sequences, which play a significant role in mathematics, are used to model and understand various situations across different scientific fields. In the literature, Fibonacci, Pell, Lucas, and Pell–Lucas are among the well-known and commonly studied number sequences. These sequences have been the subject of various studies focusing on their recurrence relations, Binet-like formulas, and algebraic properties. Vietoris numbers have also attracted attention in recent years and have been studied in several works. Number polynomials, developed in parallel with number sequences, provide a solid theoretical framework for generalizing and studying these sequences in greater depth. In this study, by combining hybrid numbers and Vietoris' polynomials, we introduce Vietoris' hybrinomials. Furthermore, we examine the recurrence relations, generating function, and Binet-like formulas.

Keywords: Vietoris' numbers, Vietoris' polynomials, hybrid number, hybrinomials.

References

- [1] Cação I., Falcão, M. I., and Malonek, H. R. (2019). *On Generalized Vietoris' Number Sequences*. Discrete Applied Mathematics, 269, 77-85.
- [2] Vietoris, L. (1958). *Über ber das Vorzeichen Gewisser Trigonometrischer Summen*. Springer.
- [3] Ruscheweyh, S., Salinas, L. (2004). *Stable Functions and Vietoris' Theorem*. Journal of Mathematical Analysis and Applications, 291 (2), 596-604.
- [4] Vietoris, L. (1984). *Eine Verallgemeinerung der Gleichung $(n + 1)! = n!(n + 1)$ und zugehörige vermutete Ungleichungen*. Monatshefte für Mathematik, 97(2), 157–160.
- [5] Cação, I., Falcao, M. I., and Malonek, H. (2017). *Hypercomplex Polynomials, Vietoris' Rational Numbers and a Related Integer Numbers Sequence*. Complex Analysis and Operator Theory, 11 (5), 1059–1076.
- [6] Catarino, P., Almeida, R. (2022). *A Note on Vietoris' Number Sequence*. Mediterranean Journal of Mathematics, 19 (1), 1–19.
- [7] Gürses, N., Çağlar Çay, D. (2025). *Toward the Determination of Vietoris-like Polynomials*. Journal of New Theory, Accepted.
- [8] Özdemir, M. (2018), *Introduction to Hybrid Numbers*. Advances in Applied Clifford Algebras, 28(1), Article number: 11.
- [9] Gürses, N., Saçlı, Y., and Yüce, S. (2024). *On Vietoris' Hybrid Number Sequence*. Turkish Journal of Mathematics, 48(4), 658-672.

On Multiplicative Magnetic Flux Ruled Surfaces

Emel KARACA

Dept. Of Mathematics, Ankara Hacı Bayram Veli University, Ankara, Türkiye

emel.karaca@hbv.edu.tr

Multiplicative Lorentz forces and multiplicative magnetic striction curves acquired by the multiplicative geodesic Frenet frame on the multiplicative striction curve of the multiplicative ruled surface introduced by the multiplicative spherical curve in a multiplicative magnetic field are expressed, and multiplicative magnetic flux ruled surfaces (or non-Newtonian magnetic flux ruled surfaces) are constructed by multiplicative magnetic vector fields along multiplicative magnetic striction curves. The main idea of this paper is to acquire magnetic flux ruled surfaces and their basic properties by employing multiplicative calculus. Finally, some examples are provided to validate the results for multiplicative magnetic flux surfaces.

Keywords: Magnetic curves, non-Newtonian analysis, magnetic striction curve.

References

- [1] Has, A., Yilmaz, B. (2024). *A Non-Newtonian Magnetic Curves in Multiplicative Riemann Manifolds*. Phys. Scr., 99, 045239.
- [2] Georgiev, S. G. (2022). *Multiplicative Differential Geometry*. Chapman and Hall/CRC, New York.
- [3] Özdemir, Z. (2021). *A Geometrical and Physical Interpretation of Quaternionic Generalized Magnetic Flux Tubes*. Chaos, Solitons & Fractals, 143, 110541.

A Study on the Dual Quaternionic Sequence with Vietoris' Components

Gülsüm Yeliz SAÇLI¹, Nurten GÜRSSES¹, Salim YÜCE¹,
Regina De ALMEIDA², and Paula CATARINO²

¹ Yildiz Technical University, Faculty of Arts and Sciences, Department of Mathematics,
34220, Istanbul, Türkiye

² University of Trás-os-Montese Alto Douro, Department of Mathematics,
5001-801, Vila Real, Portugal

yeliz.saccli@yildiz.edu.tr

In this study, a dual quaternionic sequence is constructed using the rational Vietoris numbers as coefficients. This newly defined sequence serves as a commutative four-dimensional quaternionic generalization of the classical sequence. Fundamental properties such as the conjugate and norm are examined in detail. Furthermore, two-term and three-term recurrence relations, several important identities, a Binet-like formula, and the generating function are derived.

Keywords: Vietoris' sequence, recurrence relation, Binet's formula, generating function, dual quaternion.

References

- [1] Majernik, V. (2006). *Quaternion Formulation of the Galilean Space-time Transformation*. Acta Physica Slovaca, 56(1), 9-14.
- [2] Ercan, Z., Yüce, S. (2011). *On Properties of the Dual Quaternions*. European Journal of Pure and Applied Mathematics, 4(2), 142-146.
- [3] Almeida, R. De, Catarino, P. (2021). *On a Quaternionic Sequence with Vietoris'*. Filomat, 35, 1065-1086.
- [4] Almeida, R. De, Catarino, P. (2022). *A Note on Vietoris' Number Sequence*. Mediterranean Journal of Mathematics, 19(41).
- [5] Şentürk, G. Y. (2023). *A Generalized Quaternionic Sequence with Vietoris' Number Components*. Filomat, 37(28), 9753-9768.
- [6] Gürses, N., Saçlı, G. Y., and Yüce, S. (2024). *On Vietoris' Hybrid Number Sequence*. Turkish Journal of Mathematics, 48(4), 658-672.
- [7] Almeida, R. De, Catarino, P. (2024). *Elliptic Biquaternionic Sequence with Vietoris' Numbers as Its Components*. In: Mathematical Methods for Engineering Applications. ICMASE 2023. Springer Proceedings in Mathematics & Statistics, vol 439. Springer, Cham.

On an Algorithm for a Morphological Analyzer of the Azerbaijani Language

Fariz İMRANOV¹, Aida TAGİYEVA^{2,3}

^{1,2}*Department of Electronical Academy, Presidium of ANAS, Baku, Azerbaijan*

³*National Aviation Academy, Baku, Azerbaijan*

fariz@anas.az

This work presents the algorithmic analysis and mathematical modeling of the morphological structures of the Azerbaijani language. Our objective is to precisely model the application of affixes in Azerbaijani, a member of the agglutinative language family, using formal mathematical functions and rule-based systems. Morphological transformations applied to a word root K are represented as functions $f: K \rightarrow K \times S$, where S is the set of affixes constructed based on the language's phonetic regularities and morphological rules. Phonetic changes, such as the vowel harmony function $h: V \rightarrow V$ (operating over sets of vowels) and consonant alternation transformations $\sigma: C \rightarrow C'$ (between sets of consonants), are expressed within a formal mathematical framework¹.

The algorithm models morphological functions through mathematical logic and discrete structures to generate word roots combined with appropriate affixes. The resulting root-affix pairs are stored in a structured database, which is used for evaluating morphological analysis systems and supporting machine learning applications^{2,3}.

The proposed rule-based approach is significant for low-resource languages, including Azerbaijani, as an effective tool for the mathematical description and automated analysis of generative morphology. Furthermore, it opens new perspectives on morphological analysis of natural language based on discrete mathematical models^{3,4}.

Keywords: Mathematical modeling, discrete mathematics, Azerbaijani language, morphological analysis, rule-based system, natural language processin.

Acknowledgement: This work was supported by the Azerbaijan Science Foundation: Grant N AEF-MCG-2023-1(43)-13/05/1-M-05.

References

- [1] Mahmudov, M. (2024). Linguistic Problems of Artificial Intelligence. Baku: Elm və Təhsil P. H., 376.
- [2] Mahmudov, M. (2013). Computational Linguistics. Baku: Elm və Təhsil Publishing House, 356.
- [3] Valiyeva, K. (1996). Automatic Syntactic Analysis and Synthesis of Texts. Baku.
- [4] Shiraliyev, M. (1980). Grammar of the Azerbaijani Language. Baku: Maarif.
- [5] Bektaev, K. (1995). Kazakh-Russian Dictionary. Almaty: Ana Tili.

The Influence of the Interaction with Fluids of the Hollow Cylinder with Inhomogeneous Initial Stresses on the ZGV Modes that Occur with Axisymmetric Wave Propagation

Emin T. BAGİROV¹, Surkay D. AKBAROV^{1,2}

¹ Institute of Mathematics and Mechanics of Science and Education Ministry Republic of Azerbaijan,
Baku, Azerbaijan

²Department of Mechanical Engineering, Faculty of Mechanical Engineering, Yıldız Technical
University, 34349, Istanbul, Türkiye

akbarov@yildiz.edu.tr

In the authors' studies^{1,2}, it was found that in the case where the hollow cylinder has axisymmetric inhomogeneous initial stresses caused by the hydrostatic pressure on the outer surface of the cylinder, the axisymmetric waves may have the zero group velocity (ZGV) modes (or points). This means that there are such values of wave frequency and length at which the group velocity is zero under the non-zero wave number. The question of how the magnitude of the initial inhomogeneous stresses in the cylinder affects the frequency and velocity of the ZGV points was studied in¹ only for the case where the material of the cylinder is steel. In the present investigation, this study is developed for the case where the material of the cylinder is Lucite (Plexiglas) and the fluid is water. The investigations are carried out using the exact linearized 3D equations and relations of the theory of elastic waves in bodies with initial stresses and the linearized equations of the flow of barotropic non-viscous compressible fluids.

Keywords: Axisymmetric waves, ZGV modes, initial inhomogeneous stresses, inviscid fluid, immersed hollow cylinder, hydrostatic pressure, wave dispersion.

References

- [1] Akbarov, S. D. Bagirov, E. T. (2024). *Dispersion of Axisymmetric Longitudinal Waves Propagating in an Inhomogeneous Prestressed Hollow Cylinder Immersed in an Inviscid Compressible Fluid*. Journal of Sound and Vibration, 569, 118097.
- [2] Akbarov, S. D. Bagirov, E. T. (2024). *The Dispersion of The Axisymmetric Longitudinal Waves Propagating in the Bi-layered Hollow Cylinder with the Initial Inhomogeneous Thermal Stresses*. Waves in Random and Complex Media, 34(2), 471-509.

On the Strong Solvability of a Nonlocal Boundary Value Problem for the Laplace Equation in Weighted Grand Sobolev Spaces in Rectangle

Telman GASYMOV¹, Baharchin AKHMADLI²

¹*Department of Mathematics and Mechanics, Baku State University*

²*Institute of Mathematics and Mechanics, Baku, Azerbaijan*

a_becherchin@mail.ru

The following nonlocal problem for the Laplace equation in a rectangular domain is considered:

$$u_{xx} + u_{yy} = 0, \quad 0 < x < 2\pi, \quad 0 < y < h, \quad (1)$$

$$u(x, 0) = \varphi(x), u(x, h) = \psi(x), \quad 0 < x < 2\pi, \quad (2)$$

$$u_x(0, y) = 0, u(0, y) = u(2\pi, y), \quad 0 < y < h. \quad (3)$$

Such problems have specific features in comparison with problems with local conditions. For the Laplace equation in an unbounded domain, a similar problem was considered in¹⁻², where the classical solution of the problem is studied.

In this paper, we study problem (1)-(3) in a weighted grand Sobolev space with a weight from the Muckenhoupt class. The notion of a strong solution of this problem is defined.

Keywords: Laplace equation, nonlocal problem, weighted grand Sobolev space, strong solution.

Acknowledgement: This work was supported by the Azerbaijan Science Foundation-Grant №AEF-MCG-2023-1(43)-13/06/1-M-06.

References

- [1] Moiseev, E. I. (1999). *On the Solution of a Nonlocal Boundary Value Problem by the Spectral Method*. Differents, Uravneniya, 35(8), 1094-1100.
- [2] Lerner, M. E., Repin, O. A. (1999). *On Frankl' Type Problems for Some Elliptic Equations with Degeneration of Various Types*. Differents, Uravneniya, 35(8), 1087-1093.

Hyperbolic Functions on Split Quaternions and Their Characteristics

Hamide Feyza AYKUT, Salim YÜCE

Yıldız Technical University, Faculty of Arts and Science, Department of Mathematics, Istanbul, Türkiye
fevza.aykut@yildiz.edu.tr

In the literature, exponential function¹ has been defined on split quaternions using Taylor series expansion, and the definition discussed and restated considering the characteristics of split quaternions. The characterization and properties of exponential function on split quaternion have been investigated² via a similar approach. The exponential function on split quaternion matrices has been defined and its properties have been discussed³.

In this study, hyperbolic functions have been defined with split quaternion variables and their properties and characteristics have been given considering the characteristics of split quaternion variables. Additionally, the hyperbolic functions have been divided into real and scalar parts to facilitate their applicability in composite functions.

Keywords: Split quaternion, hyperbolic function, characteristics, exponential function.

References

- [1] Kula, L., Yayli, Y. (2007). *Split Quaternions and Rotations in Semi-Euclidean Space E_2^4* . Journal of the Korean Mathematical Society, 44(6), 1313–1327.
- [2] Aykut, H. F., Yüce, S. (2023). *Exponential, Logarithmic and Trigonometric Functions on Split Quaternions (Master's Thesis)*. Yıldız Technical University.
- [3] Alagöz, Y., Oral, K. H., and Yüce, S. (2012). *Split Quaternion Matrices*. Miskolc Mathematical Notes, 13(2), 223-232.

Surface Pencil with Constant Gaussian Curvature Along a Non-null Curve

Sevilay Çoruh ŞENOCAK, Ergin BAYRAM

Mathematics, Ondokuz Mayıs University, Samsun, Türkiye

sevilay.coruhsenocak@omu.edu.tr

This study obtains surface pencil passing through a given curve with constant Gaussian curvature along this curve in Minkowski 3-space. There results are presented for spacelike curves with spacelike or timelike normal, as well as for timelike curves. The surface pencil passing through these curves are expressed parametrically. Additionally, sufficient conditions are given for these type of surface pencil. Examples are included to aid in visualizing the equations found.

Keywords: Surface pencil, Gaussian curvature, Frenet frame, Minkowski space.

References

- [1] Akutagawa, K., Nishikawa, S. (1990). *The Gauss Map and Spacelike Surfaces with Prescribed Mean Curvature in Minkowski 3-space*. Tohoku Mathematica IJournal, Second Series, 42(1), 67-82.
- [2] Bayram, E. (2020). *Verilen Bir Eğri Boyunca Gauss Eğriliği Sabit Olan Yüzeyler*. Afyon Kocatepe Üniversitesi Fen ve Mühendislik Bilimleri Dergisi, 20(5), 819-823.
- [3] Ergün, E., Bayram, E., and Kasap, E. (2014). *Surface Pencil with a Common Line of Curvature in Minkowski 3-space*. Acta Mathematica Sinica (English Series), 30(12), 2103-2118.
- [4] Bayram, E., Güler, F., and Kasap, E. (2012). *Parametric Representation of a Surface Pencil with a Common Asymptotic Curve*. Computer-Aided Design, 44(7), 637-643.
- [5] Ikawa, T. (1985). *On Curves and Submanifolds in an Indefinite-Riemannian Manifold*. Tsukuba journal of mathematics, 9(2), 353-371.
- [6] O'Neill, B. (1983). *Semi-Riemannian Geometry with Applications to Relativity*. Academic press.
- [7] Yüce, S., Çoruh, S. (2023). *A Comprehensive Insight Into Lorentzian Geometry*. Pegem Press.
- [8] Wang, G. J., Tang, K., and Tai, C.L. (2004). *Parametric Representation of a Surface Pencil with a Common Spatial Geodesic*. Computer Aided Design, 36(5), 447-59.

Rectifying Rose Ruled Surfaces

Damla BUDAK, Salim YÜCE

Department of Mathematics, Yildiz Technical University, 34220, Istanbul, Türkiye

damla.budak@std.yildiz.edu.tr

In this study, a special type of ruled surface called the rectifying rose ruled surface is investigated in the three-dimensional Euclidean space. This surface is constructed using a rose curve as the base curve and a direction vector defined as an angular linear combination of the unit tangent and binormal vectors of this curve.

After deriving the parametric equations of the surface, fundamental differential geometric properties such as the normal of the surface, the coefficients of the first and second fundamental forms, and the Gaussian curvature are analyzed. Additionally, global properties such as the striction curve and the drall are examined. The geodesic frame defined on the surface is also introduced, and the associated geodesic Frenet formulas are calculated.

In addition to theoretical investigations, the surfaces are visualized using MATLAB, and the effect of the angular variation of the direction vector on the surface geometry is studied. Rectifying rose ruled surfaces generated with different angular values are analyzed and compared.

Keywords: Rose curve, ruled surfaces, rose ruled surfaces.

References

- [1] Grandi, G. (1722). *I. Florum Geometricorum Manipulus Regiae Societati Exhibitus*. Philosophical Transactions of the Royal Society of London, 32(378), 355-371.
- [2] Grandi, G. (1728). *Flores Geometrici ex Rhodonearum et Cloeliarum Curvarum Descriptione Resultantes*. Typis Regiae Celsitudinis.
- [3] Yüce, S. (2024). *Öklid Uzayında Diferansiyel Geometri*. Pegem Akademi Yayıncılık, 10.Baskı.
- [4] Althibany, N. (2021). *Classification of Ruled Surfaces Family with Common Characteristic Curve in Euclidean 3-Space*. Turkish Journal of Science, 6(2), 61–70.

The Cross Products on Dual and Dual Lorentzian Spaces

Zehra Nur BAKIRBAŞ, Salim YÜCE

Department of Mathematics, Faculty of Arts and Sciences, Yıldız Technical University,

Istanbul, Türkiye

nur.bakirbas@std.yildiz.edu.tr

In the literature, the cross products on the dual spaces $\mathbb{D}^3$ and $\mathbb{D}^7$ and dual Lorentzian space $\mathbb{D}_1^3$ have been defined by using determinant. These cross products are expressed by inner and cross products on the spaces $\mathbb{R}^3, \mathbb{R}^7$ and $\mathbb{R}_1^3$. The cross products on the spaces $\mathbb{D}^3$ and $\mathbb{D}_1^3$ have been restated with the anti-symmetric matrices and their properties have been investigated via this approach.

In this study, the cross products on the dual spaces $\mathbb{D}^4$ and $\mathbb{D}^5$ and the dual Lorentzian spaces $\mathbb{D}_1^4, \mathbb{D}_1^5$ and $\mathbb{D}_1^7$ have been stated. Moreover, the cross product on these spaces have restated with the anti-symmetric matrices and their properties have been given via this approach.

Keywords: Cross product, anti-symmetric matrix, dual space, dual Lorentzian space.

References

- [1] Yüce, S. (2020). Sayılar ve Geometri. Pegem Academy Publishing, Ankara.
- [2] Çakır, H., Bektaş, Ö., and Yılmaz, R. (2023). *Dual Octonions and Rigid Body Kinematics*. Mathematical Methods in the Applied Sciences, 47(7), 5957-5974.
- [3] Uğurlu, H. H., Çalışkan, A. (1996). *The Study Mapping for Directed Space-like and Time-like Lines in Minkowski 3-space $\mathbb{R}_1^3$* . Math. Comput. Appl., 1(2), 142-148.

TURAN-FUNDAMENTAL SCIENCES SYMPOSIUM
TURAN-TEMEL BİLİMLER SEMPOZYUMU

Honorary Chair/Onursal Başkan: Prof. Dr. Aziz SANCAR

PHYSICS

Dielectric Relaxation and Conductivity Characteristics of Gamma-Irradiated TlGaTe₂ and TlInSe₂ Crystals

Adil ABDULLAYEV¹, Dilara GAFAROVA¹, Imamhasan MUSAZADE¹,
Ismayilov I. M.¹, and Ilhama NURIYEVA²

¹*Azerbaijan University of Architecture and Construction, Baku*²*Scientific Research Center of Azerbaijan
Medical University, Baku*
dgafarova14@gmail.com

A³B³C₂⁶ group compounds and the mixed crystals based on them constitute a large class of semiconductors with strong anisotropic (chain-like and layered) crystal structures. The physical properties of these crystals, which are analogs of disordered systems (solid solutions) with broken translational invariance of the crystal lattice, remain weakly studied despite their strong anisotropic crystal structures. One of the results of strong disorder in the system is the increased contribution of ion accumulation to the electrical conductivity during measurements, and when the contribution of ion accumulators is 10⁶ times greater than that of electrons, the system exhibits the characteristics of superionic conductors.

The electrical and dielectric properties of A³B³C₂⁶ group chain-structured semiconductor crystals in a wide temperature range (80-450 K) have been studied both parallel and perpendicular to the crystallographic "c" axis, with a complex investigation of the effects of radiation.¹ In the framework of the Mott approximation, the density of localized states (N_F), activation energy (E_a), hopping length (R), the energy difference between levels near the Fermi level (ΔE), and the concentration of deep traps (N_t) have been calculated. In the framework of the Pol-Frenkel model, the concentration of ionized centers, the free hopping distance (λ), and the value of the Frenkel coefficient (β) have been calculated, and the potential barrier form has been determined for unirradiated and 250 Mrad irradiated crystals. It has been determined that above room temperature, TlGaTe₂ and TlInSe₂ compounds transition to the superionic state. The critical transition temperature values for this state have been determined separately for each compound. The study of the temperature-frequency characteristics of the dielectric permeability and electrical conductivity of TlGaTe₂ and TlInSe₂ crystals revealed strong dielectric relaxation, and the mechanism of this phenomenon was identified. An "S"-shaped switching and a memory effect, accompanied the transition of the crystal to the superionic state during which long-term variable frequency voltage oscillations occurred.

Keywords: Superionic, hopping, activation, relaxation.

References

[1] Abdullaev A.P. (2011). *Features of Conductivity and Dielectric Permittivity of TlGaTe₂ Crystal*. AMEA Proceedings. Series of Physical-Mathematical and Technical Sciences, 31(2), 96-102.

Electrodeposition and Spray Techniques for Pt/Pd Coated Nickel Foam Electrodes

İsmet İŞGÖR¹, Sibel Eken KORKUT², and Önder YARGI¹

¹Department of Physics, Yildiz Technical University, 34220, Istanbul, Türkiye

²Department of Chemistry, Yildiz Technical University, 34220, Istanbul, Türkiye

ismetisgor.10@gmail.com

Supercapacitor (SC) electrodes were prepared by coating porous nickel foam with platinum (Pt) and palladium (Pd) powders using two different methods: electrochemical deposition and spray coating. Cyclic voltammetry (CV) and galvanostatic charge–discharge (GCD) tests were applied to evaluate supercapacitor electrode performance, focusing on specific capacitance, energy and power densities, and cycling stability. Results showed that electrodes coated via electrodeposition showed a broader CV curve and higher current response, highlighting a larger active surface area and better energy storage capability. Moreover, GCD measurements confirmed that the electrodes coated by electrodeposition has higher specific capacitance and energy-power densities. Energy and power density analyses showed that electrodeposited Pd electrodes has the highest overall performance, outperforming both spray coated and Pt based electrodes. Capacitance retention tests also showed that especially electrodeposited Pt electrodes maintained over 70% retention after 2000 cycles, significantly outperforming spray coated ones, so that shows better long-term stability¹⁻³.

Keywords: Pt, Pd, electrodes, supercapacitor, coating.

References

- [1] Xu, Y. (2024). *Conductive PEDOT:PSS Films via Electrodeposition and Spray Coating*. ACS Omega, 9(9), 8075–8083.
- [2] Chen, X. (2024). *Supercapacitor Electrodes: Is Ni Foam the Right Substrate?* Materials, 17(6), 1292.
- [3] Chen, W. (2016). *Polyaniline Nanofiber/MnO₂ Composite for SC Electrodes*. Sci. Rep., 6, 25684.

Körük Ventili (Levelling Valve) Hava Akış Karakteristiklerinin Karşılaştırmalı Teknik İncelemesi

Fatih PİLAVCI, Mesut TOSUN

Yumak Air Brake Systems, Konya Organize Sanayi Bölgesi

fatih.pilavci@mehmetyumak.com

Bu çalışmada, ağır taşıtlarda hava süspansiyon sisteminin yüksekliğini otomatik olarak ayarlayan körük ventillerinin (levelling valve) performans karakteristikleri deneysel olarak incelenmiştir. Yerli üretim ve OEM olmak üzere iki farklı valf, belirli sıcaklık ve basınç koşullarında test edilerek, valf kol açısına bağlı hava akışı değerleri karşılaştırılmıştır.¹ Ayrıca, valflerin teknik yapısının bu performansa etkisi de analiz edilmiştir. Testler, 6 bar sabit basınç altında ve +40° ile -40° kol açıları arasında gerçekleştirilmiştir. Sonuçlar, OEM valfin negatif açılarda egzoz yönünde, yerli üretim Yumak valfin ise pozitif açılarda basınç yönünde daha yüksek hava akışı sağladığını göstermektedir. Genel performans karşılaştırmasında Yumak valfi, yaklaşık %4,5 daha yüksek hava debisi sunmuştur. Her iki valf de test aralığında kararlı ve doğrusal bir akış karakteristiği sergilemiş, bu da süspansiyon sisteminin tepki süresi ve sürüş konforu açısından önemli bulunmuştur.²

Bu çalışma sonucunda, körük ventili tasarımı ve seçiminde sadece boyut veya montaj uyumu değil, aynı zamanda valfin içyapısında bulunan yay ve benzeri parçalarla belirlenen hava akış karakteristiklerinin de dikkate alınması gerektiği vurgulanmakta ve bu alanda yapılacak standartlaştırma çalışmalarına katkı sağlamayı hedeflemektedir.³

Anahtar Kelimeler: Körük ventili, süspansiyon sistemi, ağır taşıtlar, hava akış karakteristiği.

Referanslar

- [1] Kim, H., Lee, H. (2011). *Height and Leveling Control of Automotive Air Suspension System Using Sliding Mode Approach*. IEEE transactions on vehicular technology, 60(5), 2027-2041.
- [2] Xiao, J., Kulakowski, B. T., and Cao, M. (2007). *Active Air-suspension Design for Transit Buses*. International Journal of Heavy Vehicle Systems, 14(4), 421-440.
- [3] Nguyen, T. A. (2021). *Advance the stability of the vehicle by using the Pneumatic Suspension System Integrated with the Hydraulic Actuator*. Latin American Journal of Solids and Structures, 18, e403.

Sinop İlinde Nükleer Güç Santrali Öncesi Kabalı Nehrinin Sularında Tritiyum Radyoaktivitesinin Araştırılması

Serdar DİZMAN

Recep Tayyip Erdoğan Üniversitesi, Fen Edebiyat Fakültesi, Fizik Bölümü, Rize, Türkiye

serdar.dizman@erdogan.edu.tr

Tritiyum hidrojenin radyoaktif bir izotopudur ve çevredeki yüksek hareketliliği ve biyosferdeki hidrolojik döngüdeki önemi nedeniyle özel dikkat gerektirir¹. Doğal tritiyum, atmosferin üst katmanlarında bulunan gazlarla kozmik radyasyonun etkileşimi sonucu oluşur. Tritiyum ayrıca nükleer güç reaktörlerinde fisyon ve hafif element aktivasyonu ile de üretilir². Bu çalışmada, Türkiye'nin ikinci nükleer santralinin kurulacağı Sinop ilindeki Kabalı deresi sularındaki tritiyum konsantrasyonları araştırılmıştır. Bu kapsamda Kabalı deresinden (51,3 km) toplam 23 su örneği toplanmıştır. Nehir suyu örneklerindeki tritiyum konsantrasyonları sıvı sintilasyon sayacı (LSC, PerkinElmer 2910TR) ile ölçülmüştür. LSC, çevresel örneklerde tritiyumu belirlemek için en yaygın kullanılan tekniktir. Sönümü önlemek için LSC ölçümünden önce numunelere damıtma işlemi uygulanmıştır. Uygulanan yöntem için minimum tespit edilebilir aktivite (MDA) değeri 1,48 Bq/L olarak bulunmuştur. Beş su örneğinin tritiyum konsantrasyonu MDA'nın altında kalmıştır. Kabalı nehrinden alınan su örnekleri için ortalama tritiyum konsantrasyonu $2,37 \pm 0,48$ Bq/L ($20,10 \pm 4,08$ TU) olarak bulunmuştur. Bu çalışma, analiz edilen nehir suyu örneklerinin düşük tritiyum içeriğine sahip olduğunu ve bu sonuçların TSE, WHO ve USEPA tarafından belirlenen izin verilen tritiyum doz limitlerinden çok daha düşük olduğunu doğrulamıştır³.

Anahtar Kelimeler: Radyoaktivite, tritiyum, nehir suyu, Sinop.

Referanslar

- [1] Lucas, L. L., Unterweger, M. P. (2000). *Comprehensive Review and Critical Evaluation of the Half-Life of Tritium*. J. Res. Natl. Inst. Stand. Technol. 105, 541–549.
- [2] Masarik, J., Beer, J. (2009). *An Updated Simulation of Particle Fluxes and Cosmogenic Nuclide Production in the Earth's Atmosphere*. J. Geophys. Res.: Atmos. 114, 1–9.
- [3] WHO. (2011). *Guidelines for Drinking-Water Quality*. World Health Organization, Library Cataloguing-in-Publication Data, Switzerland.

Surface Engineering of Ti13Nb13Zr Implants Using Microarc Oxidation and Hydroxyapatite-Based Thermal Spraying

Bauyrzhan RAKHADILOV¹, Aidar KENGESBEKOV², Ainur SERIKBAIKYZY²,
Kussainov ARYSTANBEK², and Elaman BATANOV²

¹ PlasmaScience LLP, Ust-Kamenogorsk 070000, Kazakhstan

² D. Serikbayev East Kazakhstan Technical University, Ust-Kamenogorsk 070000, Kazakhstan
ainura.serikbaikyzy@gmail.com

This paper presents an analysis of the formation and characteristics of biocomposite coatings based on hydroxyapatite (HAp) and titanium dioxide (TiO₂) deposited on Ti13Nb13Zr alloy by combined treatment — microarc oxidation (MDO) and subsequent gas thermal spraying. Detonation (DN), cold (CN), and high-speed oxygen-fuel (HVOF) spraying were used as spraying methods. The effect of the electrolyte composition and MDO modes on the morphology, roughness, adhesion strength, and microhardness of the obtained coatings has been studied. It is shown that preliminary microarc oxidation significantly increases the adhesion of the deposited layer by increasing the porosity and roughness of the substrate. The maximum values of adhesion strength and hardness were achieved using the detonation spraying method. Morphological and elemental analysis confirmed the uniformity and density of the structures obtained by this method. The results indicate the high efficiency of the combined approach for creating functional biocompatible coatings intended for use in bone engineering and medical implants.

Keywords: Hydroxyapatite coating, microarc oxidation, detonation spraying, HVOF, titanium alloy.

Acknowledgment: This research has been funded by the Committee of Science of the Ministry of Science and Higher Education of the Republic of Kazakhstan. (Grant No. BR24992862).

References

- [1] Vassal, M., Nunes-Pereira, J., Miguel, S., Correia, I., and Silva, A. (2019). *Microstructural, Mechanical and Biological Properties of Hydroxyapatite - CaZrO₃ Biocomposites*. Ceramics International, 45.
- [2] Hu, F. et al. (2022). *Characterization of Porous Titanium-Hydroxyapatite Composite Biological Coating on PEEK by Vacuum Plasma Spraying*. Coatings, 12(4), 433.
- [3] Tang, J. et al. (2019). *A Novel Bioactive Ta/Hydroxyapatite Composite Coating Fabricated by Cold Spraying*. Materials Letters, 250.
- [4] Wang, X. et al. (2018). *Influence of Surface Structures on Biocompatibility of TiO₂/HA Coatings Prepared by MAO*. Materials Chemistry and Physics, 215, 339–345.

A New Force in Nature? The Magnetic Moment of the Muon

Zoltan FODOR

*Roland Eotvos University, Budapest, Hungary, Pennsylvania State University, USA, Wuppertal
University, Germany*

zxf5098@psu.edu

There are four forces in Nature: gravitational, electromagnetic, weak and strong forces. We call it the Standard Model of particle physics. There has been a strong tension between the measured value of the muon's magnetic moment and its Standard Model prediction. This has indicated a new force in physics. The Budapest-Marseille-Wuppertal Collaboration (of which I am the spokesperson) calculated the most important contribution —the one which dominates the uncertainties— to this magnetic moment using lattice QCD. The first result¹ was selected to be one of the breakthroughs of the year 2020 in all Science. Our most recent calculation confirms those findings with an even higher precision and shows that the tension disappeared. As a conclusion the Standard Model is verified up to 12 decimal places².

Keywords: Lattice QCD, muon, magnetic moment.

References

- [1] Borsanyi, Sz. et al. (2020). *Leading Hadronic Contribution to the Muon Magnetic Moment from Lattice QCD*. *Nature* 593 (2021) 7857, 51-55.
- [2] Boccaletti, A. et al. (2024). *High Precision Calculation of the Hadronic Vacuum Polarisation Contribution to the Muon Anomaly*. e-Print: 2407.10913.

Dielectric Properties Of CuTlS₂ Crystals Studied Via Impedance Spectroscopy

Rahim MADATOV^{1,3}, Rakshana MAMISHOVA^{1,2}, Aydan KHALIGZADE¹,
Lamiya SADIGLI¹, and Ulviyya FARADJOVA²

¹*Institute of Radiation Problems MSE, B. Vagabzadeh street 9, AZ 1143, Baku, Azerbaijan*

²*Azerbaijan University of Architecture and Construction, Ayna Sultanova st. 5, AZ-1073, Baku,
Azerbaijan*

³*National Aviation Academy, Bina, 25th km, AZ - 1045, Baku
rexsane@yandex.ru*

Determining the potential of semiconducting materials in electrical and optoelectronic applications requires an understanding of their dielectric behavior at different temperatures and frequencies^{1,2}. A strong technique for thoroughly examining dielectric relaxation mechanisms and the impact of structural flaws is impedance spectroscopy. In this section, we examine the dielectric characteristics of CuTlS₂ single crystals throughout a broad temperature (150–450 K) and frequency (200 Hz to 500 kHz) range, both in their initial state and upon exposure to gamma radiation ($D_\gamma = 25$ Mrad). Insights into charge carrier dynamics, polarization mechanisms, and defect-related inside the crystal lattice are revealed by the measured temperature- and frequency-dependent dielectric responses.

The impedance spectroscopy (IS) measurements were conducted in an evacuated cryostat within the temperature range 150–450 K, using the E7-25 device to detect key parameters such as the real part of the dielectric function, $\epsilon(\omega)$. With increasing temperature, the dielectric constant increases to a maximum value, after which an anomaly is observed: a relaxation-like decrease in $\epsilon(\omega)$ with further temperature rise. Additionally, as the measurement frequency increases, the peak corresponding to the maximum value shifts to higher temperatures and the magnitude of the peak decreases.

These variations in the temperature-frequency characteristics of dielectric permittivity are typical of Debye-type relaxation processes. The obtained $\epsilon(\omega, T)$ dependences suggest the presence of weakly bound electric charges in the crystal lattice of the solid solution, contributing to the observed dielectric behavior.

Keywords: Dielectric behavior, impedance spectroscopy, relaxation processes.

References

- [1] Madatov, R. S., Mamishova, R. M., and Baylarov, G. B. (2021). *Characteristics of Current Injection in The Narrow -Band P-Cutls Single Crystal*. Applied Physics A, 127:364.
- [2] Madatov, R. S., Mamishova, R. M. (2024). *A Study of the Effect Of Γ -Radiation on the Current-Carrying Mechanism in the P-Cutls₂ Single Crystal*. Modern Physics Letters B, 2450295.

Effect of Doping and Radiation on the Conductivity Mechanism in Layered GaS Crystals

Rahim MADATOV^{1,3}, Teymur TAGIEV¹, Rakshana MAMISHOVA^{1,2},
Aydan KHALIGZADE¹, and Lamiya SADIGLI¹

¹*Institute of Radiation Problems MSE, B. Vagabzadeh street 9, AZ 1143, Baku, Azerbaijan*

²*Azerbaijan University of Architecture and Construction, Ayna Sultanova st. 5,
AZ-1073, Baku, Azerbaijan*

³*National Aviation Academy, Bina, 25th km, AZ - 1045, Baku
aydankhaligzade@gmail.com*

Understanding the electrical properties of layered semiconductors is crucial for their application in radiation-sensitive and anisotropy-dependent optoelectronic devices¹⁻². This study aims to investigate the effect of radiation-induced defects on the anisotropic conductivity of Yb-doped GaS crystals and provide a scientific basis for the development of radiation-resistant semiconductors with tunable transport properties. We investigate the temperature-dependent electrical conductivity of GaS single crystals doped with 0.1 at.% Yb in the 125–300 K range, before and following gamma irradiation at doses up to 200 krad. The addition of Yb creates donor (~0.098 eV) and acceptor (~1.14 eV) energy levels, modifying the charge transport mechanism and resulting in anisotropic conductivity behaviour. Gamma irradiation leads to the formation of additional defect states, causing changes in activation energy and the shifting of the thermal quenching region to higher temperatures as the dose increases. Conductivity measurements parallel and perpendicular to the crystallographic c-axis ($E \parallel c$ and $E \perp c$) were carried out to evaluate the directional sensitivity to structural defects. The results suggest that Yb atoms may occupy both intralayer and interlayer positions, consequently contributing to the observed anisotropy. The overall transport phenomenon is determined by thermally activated conduction at higher temperatures and possible tunnelling effects at lower temperatures.

Keywords: Yb-doped GaS, electrical conductivity, gamma irradiation, anisotropy, activation energy.

References

- [1] Madatov, R., Najafov, A., Alakbarov, A., Tagiev, T., and Khaliqzadeh, A. (2019). *Features of Electrical and Photoelectric Properties of GaS(Yb) Onocrystals*. Journal of Physical Sciences, 74(9), 1-5.
- [2] Madatov, R. S., Tagiev, T. B., Khaligzade, A.Sh., and Mamishova, R. (2025). *Effect of Impurity and Radiation Defects on Anisotropy of Yb-Doped GaS Single Crystal*. East European Journal of Physics, 1, 240-244.

Prospects of Thermal Waters Usage in Small Power Industry

Amina MIKAYILOVA, Khagani MAMMADOV

Institutute of Radiation Problems B.Vahabzadeh 9, AZ 1143, Baku, Azerbaijan

mikayilova.a.c@gmail.com

In Azerbaijan, the installed electricity capacity last years was 8,090.8 MW, of which 6,642.0 MW was generated from 22 thermal power plants (TPPs) using organic fuel (natural gas), 1,301.8 MW from 46 hydroelectric power stations (including 35 small river plants), 66.45 MW from wind, 51.9 MW from solar, and 37.7 MW from bioenergy stations. The annual electricity production is 29,004,300 MWh, of which 2,997,500 MWh is exported. However, the country has set a goal to increase the share of renewable energy sources in the total energy balance to 30% by 2030. Estimates indicate that Azerbaijan's renewable energy reserves total up to 27,000 MW, including 23,000 MW from solar, 3,000 MW from wind, 380 MW from bioresources, and 520 MW from small rivers. Intensive efforts are underway to explore other natural energy sources in the country's economy, particularly small-scale energy.¹

Azerbaijan has abundant geothermal water reserves, with a small portion currently being used for various purposes. One way to utilize them is in small-scale energy systems for heating homes, social facilities, medical and health purposes, tourism, and recreation, among others. In Azerbaijan's thermal water sources, the water temperature reaches 90–95°C. In total, 15% of the sources have been geologically explored, and it has been shown that their operational reserves reach 250,000 cubic meters per day.² Higher values of these stocks can be expected and it is necessary to find effective ways of using them. Practically, in all ways of using thermal waters, it is necessary to study their ecological characteristics. This includes the content of radionuclides (such as radon), heavy metals, and mineral components in general.

Keywords: Thermal water, renewable energy.

References

- [1] Ibrahimova, I. Sh., Tagiyev, I. I., and Babayev, A. A. (2001). Resources of Mineral & Thermal Waters of Azerbaijan. Chashiogly, Azerbaijan.
- [2] National Academy of Sciences of Azerbaijan. (2004). *Perspectives of Development of Use of Alternative Energy Sources in Azerbaijan*. Problems of Energy, No. 1.

(E)-2-(2,4-Dihidroksibenziliden)Tiyosemikarbazon ve (E)-2-[(1H-İndol-3 il)Metilen]Tiyosemikarbazon Komplekslerinin Yapısal ve Spektroskopik Özelliklerinin DFT Metoduyla İncelenmesi

Sümeyye BÜYÜKKAZAZ¹, Yusuf ATALAY², and Davut AVCI²

¹Tıbbi Hizmetler ve Teknikler Bölümü, Sağlık Hizmetleri Meslek Yüksekokulu, Bilecik Şeyh Edebali Üniversitesi, 11230 Bilecik, Türkiye

²Fizik Bölümü, Fen Edebiyat Fakültesi, Sakarya Üniversitesi, 54187 Sakarya, Türkiye

sumeyye.buyukkazaz@bilecik.edu.tr

(E)-2-(2,4-dihidroksibenziliden)tiyosemikarbazon (**1**) ve (E)-2-[(1H-indol-3-il)metilen] tiyosemikarbazonun (**2**) komplekslerin optimize moleküler geometrisi, titreşim frekansları, ¹H NMR ve ¹³C NMR spektrumları HSEhIPBE/6-311++G(d,p) ve B3LYP/6-311++G(d,p) yoğunluk fonksiyon teorisi (DFT) metotları kullanılarak hesaplandı. NMR kimyasal kaymaları, moleküllerin moleküler yapısı ve elektronik özellikleri arasındaki ilişkiyi anlamak için oldukça yararlıdır. Bir molekülün moleküler reaktivliği elektrik ve optik özelliklerinin tanımlanmasında Frontier molekül orbitali (FMO) önemlidir. Komplekslerin frontier molekül orbitalleri ve buna bağlı η , χ , μ ve S parametreleri aynı metotlar kullanılarak hesaplanmıştır. Kompleks **1** ve **2**'nin yapısal ve spektroskopik parametrelerinin elde edilen teorik sonuçları deneysel¹ sonuçlar ile karşılaştırıldı. Elde edilen sonuçlar teorik ve deneysel¹ arasında bir uyum olduğunu göstermiştir.

Anahtar Kelimeler: Yoğunluk fonksiyon teorisi, moleküler geometri, IR, nükleer manyetik rezonans.

Referanslar

[1] Yıldız, M. (2009). Synthesis, Spectroscopic Studies and Crystal Structure of (E)-2-(2,4-dihydroxybenzylidene)thiosemicarbazone and (E)-2-[(1H-indol-3-yl)methylene]thiosemicarbazone. Journal of Molecular Structure, 919, 227–234.

Optimization of Ion-Plasma Nitriding Parameters for Ti-6Al-4V Titanium Alloy

Bauyrzhan RAKHADILOV¹, Nurtoleu MAGAZOV², Zarina ARİNGOZHINA²,
and Arnur ASKHATOV²

¹PlasmaScience LLP, Ust-Kamenogorsk 070000, Kazakhstan

²Daulet Serikbaev East Kazakhstan Technical University, Ust-Kamenogorsk 070000, Kazakhstan

zaringozhina@edu.ektu.kz

This study presents the optimization of ion-plasma nitriding parameters for Ti-6Al-4V alloy. The samples were treated at temperatures ranging from 400 to 800 °C for durations of 1–3 hours. The investigation included measurements of microhardness and elastic modulus, as well as microstructural analysis via SEM and XRD methods. It was found that the best mechanical properties were achieved at 650 °C for 3 hours, resulting in a maximum hardness of 737.4 HV and an elastic modulus of 166.9 GPa, which is attributed to the formation of a dense nitride layer. A further increase in temperature above 700 °C led to a decrease in hardness, likely due to grain growth and degradation of the surface structure¹⁻².

Phase analysis confirmed the formation of titanium nitrides (TiN, Ti₂N), which contribute to the hardening effect³⁻⁴. The obtained results allowed the determination of optimal nitriding parameters that significantly improve the surface mechanical properties. This development provides a promising approach for enhancing the durability of Ti-6Al-4V components used in high-load engineering applications and biomedical implants.

Keywords: Ti-6Al-4V, nitriding, ion-plasma treatment, microhardness, elastic modulus.

Acknowledgments: This research is funded by the Committee of Science of the Ministry of Science and Higher Education of the Republic of Kazakhstan (Grant No. AP19680457).

References

- [1] Ma X., Chen J., Liu Y., and Lu K. (2019). *Grain Refinement and Enhanced Mechanical Properties in Ti-6Al-4V Alloy Induced by Surface Severe Plastic Deformation*. Tribology International, 132, 269–277.
- [2] Amanov A., et al. (2018). *Surface Modification of Ti-6Al-4V Alloy Using Ultrasonic Nanocrystalline Surface Modification (UNSM) and its Effect on Fatigue Behavior*. Surface and Coatings Technology, 347, 503–514.
- [3] Sun Y., Bell T. (1991). *Plasma Surface Engineering of Titanium Alloys for Biomedical Applications*. Surface and Coatings Technology, 49(3), 297–302.
- [4] Luo Y., Zhang Y., Wang H., et al. (2022). *Influence of Plasma Nitriding Temperature on Microstructure and Properties of Ti6Al4V Alloy*. Materials Research Express, 9(3), 036526.

Enhancement of Visible-Light-Driven Photocatalytic Activity in Gd-Doped CdZnS Nanocomposites

İpek BALNAN¹, Sabit HOROZ^{1,2}

¹ *Department of Engineering Fundamental Sciences, Faculty of Engineering and Natural Sciences,
Sivas University of Science and Technology, Sivas, Türkiye*

² *Nanophotonics Research and Application Center, Sivas Cumhuriyet University, 58140, Sivas, Türkiye*

balnanipek@sivas.edu.tr

In this study, gadolinium (Gd)-doped CdZnS nanocomposites were synthesized to address these challenges and enhance photocatalytic performance under visible light irradiation. Structural analyses confirmed the successful incorporation of Gd³⁺ ions into the CdZnS lattice, inducing favorable modifications in crystallinity and morphology. Optical measurements revealed a red shift in the absorption edge and a reduction in bandgap energy, indicating improved light-harvesting ability in the visible spectrum¹⁻². Photoluminescence (PL) analysis demonstrated suppressed electron-hole recombination due to the presence of Gd-induced defect states, contributing to enhanced photocatalytic efficiency. The photocatalytic activity was evaluated through the degradation of Rhodamine B dye under visible light, where Gd-doped CdZnS exhibited significantly higher degradation rates compared to the undoped counterpart³. Additionally, the doped nanocomposites maintained their catalytic stability over repeated cycles. These findings highlight the potential of Gd doping as a strategy to tailor the electronic and structural properties of CdZnS-based photocatalysts, making them suitable for various environmental and optoelectronic applications.

Keywords: Photocatalysis, Gd-doping, CdZnS, visible light.

References

- [1] Abdrabou, D., Ahmed, M. K., Khairy, S. A., and El-Sherbini, T. M. (2024). *Gd₂O₃/CdS Nanocomposites were Synthesized for Photocatalytic Elimination of Methyl Blue (MB) Dye Under Visible Light Irradiation*. International Journal of Environmental Research, 18.
- [2] Basaleh, A., Ismail, A. A., and Mohamed, R. M. (2022). *Novel Visible Light Heterojunction CdS/Gd₂O₃ Nanocomposites Photocatalysts for Cr(VI) Photoreduction*. Journal of Alloys and Compounds, 927.
- [3] Panwar, S., Kumar, V., and Purohit, L. P. (2024). *Solar Light Driven Enhanced Photocatalytic Activity of Novel Gd Incorporated ZnO/SnO₂ Heterogeneous Nanocomposites*. Scientific Reports, 14.

TURAN-FUNDAMENTAL SCIENCES SYMPOSIUM
TURAN-TEMEL BİLİMLER SEMPOZYUMU

Honorary Chair/Onursal Başkan: Prof. Dr. Aziz SANCAR

CHEMISTRY

Particle Size Analysis of Polymer-Based Metal Nanoparticles Using DLS Method

Nargiz RAHIMLI, Elchin BABAYEV, Nigar SHIXVERDIYEVA, and Hacı ASLANOVA
Institute of Catalysis and Inorganic Chemistry named after academician of M. Nagiev, Baku,
Azerbaijan
narciss.rehim93@mail.ru

Particle size is a crucial parameter for supported metal catalysts, as it can significantly influence their activity and selectivity^{1,2}. In certain cases, obtaining particle size information quickly and cost-effectively is essential. While chemisorption techniques offer a more affordable alternative, they are not always suitable for supported metal catalysts. Various factors, such as surface contamination, strong metal-support interactions (SMSI), and spillover effects, can alter the stoichiometric ratio between adsorbate molecules and surface metal atoms during chemisorption measurements. Consequently, the development or adaptation of cost-effective, sensitive, and straightforward methods for particle size determination in supported catalysts remains a relevant challenge. One promising approach in this regard is Dynamic Light Scattering (DLS), which provides a rapid and inexpensive means of measuring particle size. Studies have shown that DLS results generally correlate well with those obtained from TEM analysis. In this way it allows to use more simple, sensitive and fast methods for measures of particle sizes in supported catalysts². For this purpose, PVP/Ni³ and PVP/Gelatin/Ni⁴ metal-supported catalysts were synthesized. Notably, DLS analysis revealed that the average particle size of nanoparticles in the PVP/Ni complex and the PVP/Gelatin/Ni complex was 10.6 nm and 7.5 nm, respectively.

Keywords: Supported metal catalysts, particle size, DLS method.

References

- [1] Tuo, YX. (2018). *Insight into the Support Effect on the Particle Size Effect of Pt/C Catalysts in Dehydrogenation*. J Catal, 360, 175–186.
- [2] Larichev, Y. (2021). *Application of DLS for Metal Nanoparticle Size Determination in Supported Catalysts*. Chemical Papers, 75, 2059–2066.
- [3] Rahimli, N. (2025). *Synthesis and Structural Characterization of Nickel Nanoparticles Capped Polyvinylpyrrolidone*. Asian Journal of Chemistry, 37, 309-312.
- [4] Rahimli, N. (2024). *Immobilization of Ni Metal to PVP/Gelatin Copolymer and Investigation of its Structure*. EUCHEMBIOJ 2024: I. International Conference on Chemistry and Biotechnology, Istanbul, Türkiye.

The Study of The Swelling Ratio of Gum Arabic and Acrylamide Hydrogel in Solutions with Different pH

Elmira Aliyeva, Sevda Fatullayeva

*Institute of Catalysis and Inorganic Chemistry named after academician of M. Nagiev, Baku,
Azerbaijan*

elmiraaliyeva84@gmail.com

Hydrogels are polymer-based materials capable of absorbing water, swelling, and holding a significant amount of water within their structure without dissolving. One of their key properties is the swelling rate, which can be evaluated by measuring the swelling capacity over time through free-absorbency tests conducted at regular intervals. This study aims to synthesize a physical hydrogel using gum arabic (GA) and acrylamide (AM), and to examine its swelling behavior in solutions with different pH. The successful synthesis of the grafted hydrogel and its structural characteristics were confirmed using FTIR, SEM, and XRD analyses¹. In the synthesis, gum arabic polymer was used as the primary raw material, acrylamide as the monomer, ammonium persulfate (APS) as the initiator, and N,N-methylenbisacrylamide (MBAA) as the crosslinking agent. Subsequently, the swelling ratio of the resulting copolymer was examined by immersing it in solutions with different pH values for 24 hours, allowing the evaluation of water absorption and the hydrogel's swelling behavior². As this synthesis process is intended for future application in drug delivery systems, the selected pH values (1, 2, 6 and 7) were chosen to reflect the most relevant natural physiological conditions. Overall, all samples demonstrated a substantial absorption capacity within the first 24 hours. The swelling behavior of GA-AM hydrogel in a given solvent determined by calculating the fractional change in weight relative to its initial dry weight. This is expressed as $SRW = (W - W_0)/W_0$, where W represents the weight of the swollen sample and W_0 is the weight of the dry sample³. Among the samples of the GA-AM hydrogel exhibited the highest swelling ratio up to $490 \pm 4\%$ at pH 2. It should also be noted that the amount of crosslinking agent added was equivalent to 15% of the polymer amount.

Keywords: Hydrogels, gum arabic, acrylamide, swelling ratio.

References

- [1] Asy-Syifa, N. (2022). *The Study of the Swelling Degree of the PVA Hydrogel with Varying concentrations of PVA*. J. Phys.: Conf. Ser. 2243 012053.
- [2] Mammadova, S. (2021). *Obtaining a Hydrogel Based on Chitosan, Study its Structure and Sorbtion Ability with Levofloxacin Antibiotics*. Chemical Problems, 4(19), 215-223.
- [3] Maura, C. (2017). *Practical Strategy to Realistically Measure the Swelling Ratio of Poly (Dimethylsiloxane) Without Underestimation Due to the Solvent Volatility*. Polymer, 113, 187-192.

Synthesis, Structure, and Properties of New Alicyclic and Aromatic Amidophosphates

Akbota KUANDYKOVA, Nazgul AKIMBAYEVA, Nurgul ZHUMANOVA, Bulat DZHIEMBAEV,
Banu DIYAROVA, Nurbala MYRZAKHMETOVA, and Ainur YEREZHEPOVA

Kazakh National Woman's Teacher Training University, Almaty, 700420, Kazakhstan

bota.kuandyk1702@gmail.com

Research into methods for synthesizing new functionally substituted derivatives of amidophosphonates of the alicyclic and aromatic series, elucidating their structures, and identifying biologically active compounds among them is largely driven by their unique set of chemical properties and broad spectrum of biological activity. Further development of both theoretical and applied aspects of our research is focused on designing synthesis methods for new amidophosphates and their modified derivatives using the Todd-Atherton reaction under microwave irradiation. As a result, the following compounds were isolated and characterized: dialkyl-N-(1-ethynylcyclohex-1-yl)amidophosphates, N,N'-diethylenimino- and (N,N'-tetramethylamino)-N''-(1-ethynylcyclohex-1-yl)triamidophosphates, dialkylphenylphosphoramidates, and tetraalkyl 1,3-phenylenebis-(phosphoramidates).¹⁻² Furthermore, biological testing of these compounds has identified agents demonstrating high biological activity concerning the growth, development, and yield of agricultural crops.³

Keywords: Amidophosphates, Atherton-Todd reaction, biological activity, microwave irradiation.

Acknowledgement: This research has been funded by the Science Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan AP №19678514.

References

- [1] Dzhiembaev, B. (2023). *Synthesis of Amidophosphates Based on 1-Ethynyl-1-Aminocyclohexane Upon Microwave Activation and Their Biological Activity*. Eur. Chem.-Technol. J., 25(1), 57-64.
- [2] Kuandykova, A. (2023). *Synthesis of New Symmetrical Diamidophosphates Based on Meta-Phenylenediamine Under Microwave Irradiation*. Russ. J. Gen. Chem., 93(6), 1-5.
- [3] Patent KZ 35532. (2022). *Application Of 0,0-Diethyl-1-(3,4-Dimethoxyphenyl)-(1-Ethynylcyclohexylaminomethyl) Phosphonate as a Growth Stimulator of Vegetable Crops*. Dzhiembaev, B. Bull. #8.

Development of Lead-Free Epoxy Resin Composites for X-Ray Shielding

Rana KHANKISHIYEVA^{1,2}, Vugar KHUDAVERDIEV¹, Haji Vahid AKHUNDZADA¹,
Shahla AHMED², and Gunel MAMMADOVA¹

¹ Institute of Radiation Problems, Baku, Azerbaijan

² Azerbaijan University of Architecture and Construction, Baku, Azerbaijan

rana.khankishiyeva@azmiu.edu.az

The increasing demand for sustainable and effective radiation shielding materials has driven the search for alternatives to traditional lead-based shields^{1,2}. This study investigates lead-free epoxy resin composites incorporating Bi₂O₃ and BaSO₄ powders for X-ray shielding. Composites were fabricated using open mold casting and characterized via FTIR, SEM and XRD. Radiation shielding efficiency was assessed using ²⁴¹Am and ¹³⁷Cs gamma sources in a gamma spectroscopy setup at the Institute of Radiation Problems, Azerbaijan. Shielding performance was analyzed through mass attenuation coefficients and half-value layer (HVL) at 60 keV and 120 keV. Mechanical properties, including tensile strength and thermal stability, were also measured. Bi₂O₃/BaSO₄-based composites demonstrated excellent radiation attenuation comparable to lead-based materials, while maintaining lightweight and flexible characteristics. This study underscores the potential of epoxy resin composites as sustainable alternatives for X-ray shielding. Future research will focus on improving dispersion techniques to enhance homogeneity and shielding performance.

Keywords: Epoxy resin, lead-free shielding, radiation protection, X-Ray Shielding.

References

- [1] Karabul, Y., İçelli, O. (2021). *The Assessment of Usage of Epoxy-Based Micro and Nano-Structured Composites Enriched with Bi₂O₃ and WO₃ Particles for Radiation Shielding*. Results in Physics, 26, 104423.
- [2] Alshahri, S., Alsuhybani, M., Alosime, E., Almurayshid, M., Alrwais, A., and Alotaibi, S. (2021). *LDPE/bismuth Oxide Nanocomposite: Preparation, Characterization, and Application in X-Ray Shielding*. Polymers, 13(18), 3081.

Alunit Cevherinin RSM Kullanılarak Liçinin Optimizasyonu

Pervane BABAYEVA¹, Arif HEYDEROV¹, M. Deniz TURAN²,

Hasan NİZAMOĞLU², and Tuğrulhan DEMİR

¹Azerbaycan Cumhuriyeti Bilim ve Eğitim Bakanlığı
M. Nagiyev adına Kataliz ve İnorganik Kimya Enstitüsü

²Fırat Üniversitesi, Elazığ, Türkiye

tugrul235723@gmail.com

Araştırmada, alunit cevherinin NaOH ile muamelesi sırasında çözeltiye geçen maksimum alüminyum miktarı ve minimum fosfor miktarı RSM ampirik modeli kullanılarak incelenmiştir¹. Bu amaçla, Box-Behnken tasarım yöntemi kullanılarak üç farklı parametrenin (işlem süresi, işlem sıcaklığı ve sıvı-katı oranı) etkileri araştırılmıştır². Deneyler 20-80 °C, 30-270 dk ve 5-95 mL/g aralığında yürütülerek, proses için optimum koşullar belirlendi. RSM kullanılarak önerilen model denkleminin deneysel verilerle iyi bir uyum gösterdiği ve model için $R^2=0,7572$, $R^2=0,9792$ korelasyon katsayılarının belirlendiği görülmüştür. Bu aşamada optimum koşullar NaOH konsantrasyonu: %7, yıkama süresi: 30 dk, yıkama sıcaklığı: 69 °C, sıvı/katı oranı: 56 mL/g olarak belirlendi. Bu optimum koşullar altında çözeltiye geçen alüminyum miktarı %89, fosfor miktarı ise %18 olmuştur. . Alunitin mineralojik analizleri X-ışını kırınımı (XRD, Shimadzu XRD-6000) kullanılarak yapıldı ve morfolojik yapısı taramalı elektron mikroskobu (JEOL JSM 7001F FE-SEM) kullanılarak karakterize edildi. Alunitlerin kimyasal bileşimi XRF cihazı (Rigaku Primus IV) kullanılarak belirlendi ve elde edilen oksitlenmiş bileşen sonuçları elementel sonuçlara dönüştürüldü.

Anahtar Kelimeler: Alunit, alüminyum, fosfor, RSM.

Referanslar

- [1] Taghiyev, E. I., Tagijev, E., Agajeva, L. (2019) *Cost Effective Technology of Alunite Ore Processing*. Int. J. Chem., 11, 36. <https://doi.org/10.5539/ijc.v11n1p36>.
- [2] Turan, M. D. (2018). *Statistical Approach to Mineral Engineering and Optimization*, Intech,p.13. <https://www.intechopen.com/books/advanced-biometric-technologies/liveness-detection-in-biometrics>.

Synthesis of Green Chemistry-Based Catalyst for Carbon Monoxide Neutralization

Gunel AZIMOVA, Sima ZULFUGAROVA, and Zuleykha ALESKEROVA

Institute of Catalysis and Inorganic Chemistry, Baku, Azerbaijan

ezimova2015@gmail.com

Literature analysis shows that synthesis and research based on green chemistry have been rapidly developing recently. The basic principle of green chemistry is the study of chemical products and processes that reduce or eliminate the use of environmentally hazardous substances^{1,2}. At the same time, catalytic oxidation of carbon dioxide is very actual issue. As we know, carbon monoxide is one of the toxic substances emitted into the atmosphere. Therefore, its oxidation to carbon dioxide is one of the important problems³.

The salts $\text{Fe}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$ and $\text{Cu}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ were used as precursors for the synthesis of copper iron system. Pomegranate peel extract is additionally used here. Firstly, the salts are dissolved in a small amount of distilled water, then pomegranate peel extract is added. It is mixed by heating to 80-90°C in a magnetic stirrer for 1 hour. Then heated in a drying cabinet, igniting at a temperature of 175-200°C to form a powder. Finally, it is thermally treated for 4 hours in a muffle furnace at 500°C.

The specific surface area of the sample was determined by low-temperature nitrogen adsorption using the BET method on a SORBI-MS device (ZAO META, Russia). The specific surface area of the obtained powder was determined by low temperature nitrogen adsorption by the BET method and as a result showed 36 m²/g. The catalytic activity of the sample in the oxidation of CO in CO₂ in the 180° temperature was determined.

Keywords: Green chemistry, pomegranate peel extract, carbon monoxide.

References

- [1] Y. Tamboli, Q. et al (2023). *Green Synthesis of Cobalt Ferrite Nanoparticles: An Emerging Material for Environmental and Biomedical Applications*. Journal of Nanomaterials, 1-15.
- [2] Naik, M. M. et al (2019). *Green Synthesis of Zinc Doped Cobalt Ferrite Nanoparticles: Structural, Optical, Photocatalytic and Antibacterial Studies*. Nano-Structures & Nano-Objects, 19, 100322-100334.
- [3] Azimova, G. R. et al (2022). *Synthesis of Transition Metal Ferrites (Co, Cu, Ni, Mn) by the Sol-gel Method with Combustion and the Use of Microwave Processing*. Bulletin of the University of Karaganda Chemistry, 108(4), 118-135.

TURAN-FUNDAMENTAL SCIENCES SYMPOSIUM
TURAN-TEMEL BİLİMLER SEMPOZYUMU

Honorary Chair/Onursal Başkan: Prof. Dr. Aziz SANCAR

BIOLOGY

Potential Use of Watermelon Seed Oil as an Alternative Linoleic Acid Source Enhanced by Coffee-Derived Caffeic Acid in Cholesterol Treatment

Deniz Duru DERVİŞ

Yildiz Technical University, Davutpaşa Campus, 34220, Istanbul, Türkiye

duru.dervis16@gmail.com

Cardiovascular diseases remain the leading cause of death worldwide, with dyslipidemia and elevated LDL cholesterol levels playing a central role. One critical nutritional factor contributing to this issue is the excessive consumption of omega-6-rich vegetable oils, particularly corn and sunflower oil, which are widely used as primary dietary sources of linoleic acid (LA), an essential omega-6 fatty acid. While LA is required for the synthesis of arachidonic acid—a molecule involved in inflammation and neural transmission—its disproportionate intake can lead to metabolic imbalances and pro-inflammatory states when the omega-6 to omega-3 ratio is disrupted.¹

This study proposes watermelon seed oil as a lesser-known yet promising alternative source of linoleic acid, offering a more moderate concentration that may help mitigate the risks associated with excessive omega-6 intake. Additionally, the research investigates the potential synergy between LA and caffeic acid, a phenolic compound naturally present in coffee. Literature review suggests that caffeic acid can enhance the solubility and bioavailability of linoleic acid, potentially increasing its absorption efficiency and functional value when co-administered.

We hypothesize that integrating a controlled extract of watermelon seed oil into coffee, a globally consumed beverage, could serve as a novel, functional approach to essential fatty acid supplementation. This model not only utilizes an agricultural byproduct that is often discarded but also provides a nutritional strategy that aligns with public health goals aimed at reducing cardiovascular risk. Although empirical validation through laboratory experimentation is still needed, the conceptual foundation bridges nutritional biochemistry, food science, and preventive medicine.

Keywords: Linoleic acid, watermelon seed oil, LDL cholesterol, caffeic acid, functional nutrition, cardiovascular health.

References

[1] Medjakovic, S., Hobiger, S., Ardjomand-Woelkart, K., Bucar, F., and Jungbauer, A. (2016). Pumpkin seed extract: *Cell Growth Inhibition of Hyperplastic and Cancer Cells, Independent of Steroid Hormone Receptors*. *Fitoterapia*, 110, 150-156.

Bazı Yonca (*Medicago sativa* L.) Genotiplerinin iPBS-retrotranspozon Markörleri ile Genetik Çeşitliliğin Değerlendirmesi

Meliha Feryal SARIKAYA, Muhammad Azhar NADEEM

*Sivas Bilim ve Teknoloji Üniversitesi, Tarım Bilimleri ve Teknoloji Fakültesi, Tarla Bitkileri Bölümü,
Sivas, Türkiye*

feryalsarikaya@gmail.com

Yonca, dünya çapında 32 milyon hektarlık küresel üretim alanı ile dünyada en çok yetiştirilen yem bitkisidir.¹ Ayrıca biyoenerji, insan gıdası ve moleküler tarım gibi farklı kullanım alanları da vardır.² Bu çalışmada, Türkiye'nin farklı bölgelerinden toplanan 88 adet yonca genotipi ve 6 adet ticari yonca çeşidinin genetik çeşitliliğinin 10 adet polimorfik iPBS-retrotranspozon belirteci ile belirlenmesi amaçlanmıştır. Çalışma sonucunda 10 primerden toplam 351 polimorfik bant elde edilmiş olup, 2074 ve 2075 numaralı primerler 41 adet ile en fazla alel sayısını üretmiştir. Ortalama polimorfizm bilgi içeriği (PIC) 0,31 olarak kaydedilmiştir. Genetik çeşitlilik indeksleri ortalama 0,32 gen çeşitliliği (h) ve 0,49 Shannon bilgi indeksi (I) değerlerini göstermiştir. Komşu birleştirme analizi tabanlı kümeleme, tüm genotipi iki gruba ayırmıştır. Türk yonca germ plazması, yeni çeşitlerin dayanıklılığını ve verimliliğini artırmayı amaçlayan ıslah programları için hayati önem taşıyan önemli bir genetik çeşitlilik sergilemektedir.

Anahtar Kelimeler: Yonca, genetik çeşitlilik, iPBS-retrotranspozon markörleri, bitki ıslahı.

Teşekkür Bölümü: Bu çalışma, Sivas Bilim ve Teknoloji Üniversitesi Bilimsel Araştırma Projeleri Koordinasyon Birimi tarafından 2024-DTP-TBT-0002 numaralı proje ile desteklenmiştir.

Referanslar

- [1] Mouttet, R., Escobar-Gutierrez, A., Esquibet, M., Gentzbittel, L., Mugniery, D., Reignault, P., Sarniguet, C., and Castagnone-Sereno, P. (2014). *Banning of Methyl Bromide for Seed Treatment: Could Ditylenchus Dipsaci Again Become a Major Threat to Alfalfa Production in Europe?* Pest Management Science, 70, 1017–1022.
- [2] Zhang, Z., Min, X., Wang, Z., Wang, Y., Liu, Z., and Liu, W. (2017). *Genome-wide Development and Utilization of Novel Intron-length Polymorphic (ILP) Markers in Medicago Sativa*. Molecular Breeding, 37, 1–8.

γ (Gama) Işınlarıyla Muamelenin Bitkilerde Tuz (NaCl) Stresine Tepkisi

Guliyeva NARMİN, Elimkhan JAFAROV

Radyasyon Problemleri Enstitüsü, B. Vahabzadeh 9, AZ 1143, Bakü, Azerbaijan

guliyevanarmin.1991@gmail.com

Bilindiği üzere, günümüzde ekolojik problemlerin arttığı bir dönemde bitkilerin gelişiminin zayıflaması ve hatta tamamen yok olması süreçleri hızla devam etmektedir. Bu süreçlerin artmasında antropojen etkilerin rolü büyüktür. Ekolojik krizin şiddetlendiği bir dönemde, çevrenin stres koşullarında bitkilerin büyüme ve gelişimi zayıflamaktadır. Abiyotik stres faktörleri, kültür bitkilerinin gelişiminin yaklaşık %30 - 35 oranında azalmasına neden olmaktadır. Aynı zamanda insan nüfusunun kitlesel şekilde artması, gıda ürünlerine olan talebin de artmasına yol açmaktadır. Bu nedenle, stres koşullarında yetişip ürün verebilen bitki türlerinin geliştirilmesi uygun görülmektedir. Dünya genelinde tarıma elverişli toprakların yaklaşık %20'si tuzlanmaya (şorlanmaya) maruz kalmıştır. Tüm bunlar göz önüne alındığında, ekolojik açıdan kirlenmiş alanlarda yetişip ürün verebilen bitki türlerinin geliştirilmesi temel hedeflerden biridir.²⁻³ Tuz stresinin etkisine bitkilerde ilk olarak kök hücreleri maruz kalır. Araştırma çalışmasında, γ ışınlarıyla muamele edilmiş tohumların tuz (NaCl) stresi koşullarındaki gelişimi incelenmiştir. γ ışınlarıyla farklı dozlarda muamele edilmiş bitkilerin, düşük dozlarda daha fazla direnç kazandığı belirlenmiştir. Işınlanmış tohumların tuz stresine karşı yanıt reaksiyonlarını incelemek amacıyla, MDA ve H₂O₂ miktarının yanı sıra, antioksidan bir enzim olan katalaz enziminin aktivitesi de değerlendirilmiştir. Araştırmamızın sonuçları göstermektedir ki, tohumların ekiminden önce 10 Gy ışın dozu ile muamele edilmesi durumunda, nispeten düşük tuz konsantrasyonlarında (1-10 mM) malondialdehit, hidrojen peroksit ve katalaz miktarındaki değişim, oksidatif stresin azaldığını gösterebilir. Tahmin ediyoruz ki, tohumların ekiminden önce 10 Gy ışın dozu ile muamele edilmesi, 1-10 mM tuz (NaCl) konsantrasyonunda bitkilerin tuzluluğa karşı direncini artırabilir.¹

Anahtar Kelimeler: γ (Gama) ışınlanması, tuz stresi, MDA, H₂O₂, katalaza.

Referanslar

- [1] Bailly, C., Leymarie, J., Lehner, A., [et al.]. (2004). *Catalase Activity and Expression in Developing Sunflower Seeds as Related to Drying*. Journal of Exp. Bot., 55(396), 475–483.
- [2] Gitelson, A., Solovchenko, A. (2018). *Non-invasive Quantification of Foliar Pigments: Possibilities and Limitations of Reflectance- and Absorbance-based Approaches*. J. Photochemistry and Photobiology B: Biology, 178, 537-544.
- [3] Mancini, A., Restivo, F.M. [et al.]. (2006). *Oxidative Stress as DNA Damage in Different Transgenic Tobacco Plants*. Plant Sci., 170(4), 845-852.

POSTER PRESENTATION ABSTRACTS

TURAN-FUNDAMENTAL SCIENCES SYMPOSIUM
TURAN-TEMEL BİLİMLER SEMPOZYUMU

Honorary Chair/Onursal Başkan: Prof. Dr. Aziz SANCAR

MATHEMATICS

Ağırlıklı Standart Olmayan Banach Uzaylarında Harmonik Fonksiyonların Yapısal Analizi

Bilal T. BİLALOV, Yonca SEZER, and Eylem YAŞAR

Yıldız Teknik Üniversitesi, Fen-Edebiyat Fakültesi, Matematik Bölümü, İstanbul, Türkiye

eylem.yasar@std.yildiz.edu.tr

Bu çalışma, ağırlık yapısına sahip standart olmayan Banach fonksiyon uzaylarında tanımlı harmonik fonksiyonların norm sürekliliği ve sınır değer davranışlarını incelemektedir. Mucenhaupt sınıfına ait ağırlık fonksiyonları ve yeniden düzenleme altında değişmeyen norm yapıları çerçevesinde bu fonksiyonların analitik özellikleri araştırılmıştır. Poisson çekirdeği, Hilbert dönüşümü ve Boyd indisleri gibi klasik analitik araçlar yardımıyla, Hardy tipi uzaylarda tanımlı bazı singüler integral operatörlerinin sınırlılığı analiz edilmiştir. Operatörlerin norm sürekliliği ve fonksiyonel yapılarla etkileşimleri, Boyd indisleri aracılığıyla karakterize edilmiştir. Elde edilen bulgular, ağırlıklı fonksiyon uzaylarında tanımlı sınırlı operatörlerin yapısını ortaya koymakta ve potansiyel teorisi ile sınır değer problemleri gibi alanlarda teorik katkılar sunmaktadır.

Anahtar Kelimeler: Harmonik fonksiyonlar, ağırlıklı Banach uzayları, boyd indisi, Hardy uzayları, operatör teorisi.

Referanslar

- [1] Bennett, C., Sharpley, R. (1988). Operatörlerin Enterpolasyonu. Academic Press.
- [2] Bilalov, B. T. (2022) *Hardy Banach Fonksiyon Sınıfları ve Kenar Değer Problemleri Yöntemi*. Bakü, Elm.
- [3] Kokilaşvili, V. (2018). *Standart Olmayan Fonksiyonel Uzaylarda İntegral Operatörler*. Cilt 1, Birkhäuser.

Structural Properties of the $h_p(X)$ Class of X-Valued Harmonic Functions

Bilal T. BİLALOV^{1,2,3}, Yonca SEZER², Afra BÜYÜKARSLAN², and Ümit İLDİZ²

¹Odlar Yurdu University, Baku, Azerbaijan

²Department of Mathematics, Yıldız Technical University, Davutpaşa, 34220 Istanbul, Turkey

³Department of Non-harmonic Analysis, Institute of Mathematics and Mechanics, The

Ministry of Science and Education of the Republic of Azerbaijan, Baku, Azerbaijan

afra.buyukarslan@std.yildiz.edu.tr

In this work, we study the class $h_p(X)$ of X-valued harmonic functions defined on the unit disc, where X is a UMD space and $1 < p < \infty$. The space is constructed using X-valued trigonometric series in the Bochner space $L_p([- \pi, \pi]; X)$, and the t-basis property of the exponential system is established. This structure provides the foundation for analysing boundary value problems, particularly the Laplace equation with directional derivative conditions. It is proved the Fredholmness of this problem. Additionally, harmonic and analytic X-valued function spaces are examined, including their relationship through generalized Cauchy-Riemann equations derived from a *-involution on X. A vector-valued version of Fatou's theorem is also established. The study introduces derivative-based subspaces of $h_p(X)$, and the index of associated boundary value problems is computed¹⁻⁴.

Keywords: X-valued harmonic functions, Noether, Fredholm, Bochner space.

Acknowledgement: This research was supported by Scientific Research Project Coordination Unit of Yıldız Technical University (Project ID:6700 Project Code: FBA-2025-6700).

References

- [1] Adams, D. R. (2016). Morrey Spaces. Switzerland: Springer.
- [2] Bilalov, B. T., Ahmedzadeh, N. R. and Garayev, T. Z. (2022). *Some Remarks on Solvability of Dirichlet Problem for Laplace Equation in Non-standard Function Spaces. Mediterranean Journal of Mathematics*, 19(3), 133.
- [3] Arendt, W. (2016). *Vector-valued Holomorphic and Harmonic Functions. Concrete Operators*, 3(1): 68–76. <https://doi.org/10.1515/conop-2016-0007>
- [4] Bilalov, B. T., Sadigova, S.R. (2025). *The Concept of t-basis and Vector-valued Hardy Classes. Turkish Journal of Mathematics*, 49(3), 261–286. <https://doi.org/10.55730/1300-0098.3588>.

Exact Solutions of the Time fractional Benney-Luke Equation and the Second Order Benjamin-Ono Equation

Özlem BAKŞİ, Seda KIZILBUDAK ÇALIŞKAN

Department of Mathematics, Faculty of Arts and Science, Yıldız Technical University,

34210, Istanbul, Türkiye

skizilb@yildiz.edu.tr

In this study, we obtain the exact solutions of the time-fractional non-linear Benney-Luke equation and the second order Benjamin-Ono equation, which are difficult to solve analytically by converting them into ordinary differential equations using a conformable derivative transformation and the exponential function method. This method helps us examine the behavior of the solutions and understand the effects of different parameter values on them. We also present graphical illustrations of some solutions for different values of the parameters in our study.

Keywords: Benney-Luke equation, Benjamin-Ono equation, exponential method, conformable derivative, exact solutions.

References

- [1] Khalil, R., Al Horani, M., Yousef, A., and Sababheh, M. (2014). *A New Definition of Fractional Derivative*. J. Comput. Appl. Math., 264, 65–70.
- [2] Benney, D. J., Luke, J. C., (1964). *Interactions of Permanent Waves*. Journal of Mathematics and Physics, 43, 209–213.
- [3] Durur, H., Yokuş, A. (2021). *Exact Solutions of the Benney–Luke Equation via $(1/G')$ -expansion Method*. Bilecik Şeyh Edebali Üniversitesi Fen Bilimleri Dergisi.
- [4] Wazwaz, A. M. (2017). *A Study of a Generalized Benney–Luke Equation with Time-dependent Coefficients*. Nonlinear Dynamics.
- [5] Benjamin, T. B., Ono H. (1967). *Internal Waves of Permanent Form in Fluids of Great Depth*. Journal of Fluid Mechanics, 29, 559–592.
- [6] Ono, H. (1975). *Algebraic Solitary Waves in Stratified Fluids*. Journal of the Physical Society of Japan, 39(5), 1082–1091.
- [7] Tasbozan, O. (2019). *New Analytical Solutions for Time Fractional Benjamin-Ono Equation Arising Internal Waves in Deep Water*. China Ocean Engineering, 33(4), 593–600.
- [8] He, J. H. (2013). *Exp-function Method for Fractional Differential Equations*. Int. J. Nonlinear Sci. Numer. Simul., 14, 363–366.

Lipschitz Stability Analysis of Nonlinear Set-Valued Differential Equations with Initial Time Differences

Tuba SATILMIS

Department of Mathematics, Gebze Technical University, Kocaeli, Türkiye
t.satilmis2022@gtu.edu.tr

In this paper, we examine the connection between an unperturbed set-valued differential system and a perturbed one, where the systems differ in both initial conditions and initial times. By applying the variation of parameters method, we derive integral representations of solutions and establish Lipschitz stability criteria for nonlinear set-valued differential systems. A comparison is also made between general stability and Lipschitz stability to emphasize their differences. The theoretical background and criteria are inspired by earlier contributions on initial time difference stability and Lyapunov-based analysis¹⁻³.

Keywords: Lipschitz stability, variation of parameters, set-valued differential equations, perturbations, initial time difference.

References

- [1] Yakar, C. (2000). *Stability Analysis of Nonlinear Differential Systems with Initial Time Difference*. Florida Institute of Technology.
- [2] Yakar, C., Shaw, M. (2005). *A Comparison Result and Lyapunov Stability Criteria with Initial Time Difference*. Dynamics of Continuous, Discrete and Impulsive Systems Series A, 12(6), 731.
- [3] Shaw, M. D., Yakar, C. *Lipschitz. Stability Criteria with Initial Time Difference*. Journal of Applicable Analysis (to appear).

Orlicz-Sobolev Uzaylarında Yerel Olmayan Bir Problemin Güçlü Çözülebilirliği

Bilal T. Bilalov^{1,2}, Yonca Sezer¹, and Ümit İldız^{1,3}

¹Matematik Bölümü, Yıldız Teknik Üniversitesi, 34220, İstanbul, Türkiye

²Cebir ve Matematiksel Mantık Bölümü, Matematik ve Mekanik Enstitüsü,
Azerbaycan Cumhuriyeti Bilim ve Eğitim Bakanlığı, Bakü, Azerbaycan

³Yönetim Bilişim Sistemleri Bölümü, Marmara Üniversitesi, R.T.E. Külliyesi, Maltepe, İstanbul, Türkiye

umit.ildiz@std.yildiz.edu.tr

Bu çalışmada sınırsız bir bölgede Laplace denklemi için yerel olmayan bir problem ele alınmış ve bu problem için güçlü çözüm kavramı tanımlanmıştır. Spektral problemin kök fonksiyonlar sisteminin Orlicz uzayı için bir baz oluşturmasından yararlanılmıştır. Simetrik uzaylar için Boyd indeksleri kavramı kullanılarak spektral yöntem uygulanmış ve Orlicz normu tarafından tanımlanan Sobolev uzaylarında problemin güçlü çözülebilirliği gösterilmiştir. Bu çalışmada ilk kez Orlicz-Sobolev uzaylarında çözümün varlığı ve tekliği ortaya konmuştur. Orlicz-Sobolev uzaylarında problemi çözmek, daha genel bir yapıya sahip olan simetrik Sobolev uzaylarına çözümün genelleştirilmesinin önünü açacaktır.

Anahtar Kelimeler: Orlicz-Sobolev uzayları, Laplace denklemi, yerel olmayan problem, güçlü çözüm.

Teşekkür Bölümü: Bu çalışma Yıldız Teknik Üniversitesi Bilimsel Araştırma Projeleri Koordinasyon Birimi tarafından desteklenmektedir (Proje ID:6596, Proje Kodu: FBA -2025-6596).

Referanslar

- [1] Bilalov, B. T., Nasibova, N. P., Softova, L. G., and Tramontano, S. (2023). *Global Solvability of the Laplace Equation in Weighted Sobolev Spaces*. arXiv preprint:2307.07776.
- [2] Krasnoselski, M. A., Rutitski, Y. B. (1958). *Convex Functions and Orlicz Spaces*. GIFML, Moskow.
- [3] Moiseev, E. I. (1999). *The Solution of a Nonlocal Boundary Value Problem by the Spectral Method*. Differ. Equations, 35(8), 1105–1112, transl. from Differ. Uravn. (1999). 35(8), 1094–1100.

Ülke İçinde Karayolu ile Yük Taşımacılığı

Parvin QULİYEVA

Azərbaycan Mimarlıq və İnşaat Universiteti, Bakü, Ayna Sultanova Cad., 11

parvin.quliyeva@sabah.edu.az

Makalede karayoluyla eşya taşımacılığı anlatılmaktadır. Ülkemizde gerçekleştirilen hızlı yapılaşma, nüfus artışı, bölgelerimizin gelişmesi ülke içinde yük taşımacılığı hizmetlerine olan ihtiyacı artırmaktadır. Yük taşımacılığında taşımacılık mevzuatlarına uymanın yanı sıra, müşterinin gereksinimlerini karşılamak, beklentilerini karşılamak ve onları memnun etmek de çok önemlidir. Ancak, rekabet ortamında yük taşımacılığı faaliyetinde bulunan girişimci veya işletmelerin (kamu veya özel) tüm bu gereklilikleri yerine getirmenin yanı sıra, kullandıkları araçların teknik işleyişini de denetlemeleri gerektiği unutulmamalıdır. Aynı zamanda aracı kullanan sürücünün, gidilecek yere göre çalışma programı da dikkate alınmalıdır.

Yükün cinsine, minimum maliyetle taşınacak araca, yükün parametrelerinin doğru bir şekilde incelenmesi, acil olup olmadığı vb. bilgilerin kayıt altına alınması ve tüm bunlar göz önünde bulundurularak uygun aracın belirlenmesi gerekmektedir. Acil kargoların özel şartlarda taşınabilmesi esastır. Taşınma işlemi yapılmadan önce, gerekli değerlendirmenin yapılabilmesi için adreste önceden inceleme yapılmalı ve buna göre değerlendirme yapılmalıdır. Türüne göre kargo paketlemesi ve gerekirse işçilik hizmeti de sunulmalıdır. Değerlendirme sırasında tüm bunların dikkate alındığının müşteriye bildirilmesi gerekmektedir. Bu ve bunun gibi birçok parametrenin tek merkezden erişilebilir ve yönetilebilir olmasını sağlayacak bir yük bilgi yönetim programının geliştirilmesi önerilmektedir.

Anahtar Kelimeler: Yük sınıflandırması, taşıma süreci, lojistik yazılımı, taşıyıcı firma, GPS izleme.

Referanslar

- [1] <https://e-qanun.az/framework/18722>
- [2] Cavadov, Ə. Ə. (2009). *Karayolu Yük Taşımacılığı*, Bakü, Təhsil.
- [3] *U.S. Senate Hearing on Motor Carrier Safety Programs*, 1999.
- [4] TURKSAN Center. (2022). The XXVIII International Scientific Symposium "Karabakh is My Native Land", dedicated to the 190 anniversary of Khurshidbanu Natavan 30 July, 2022 ISBN: 978-605-71427-9-5 Amsterdam/ Holland.

Fe-Si-C Amorf Şerit Numunelerinin Araştırılması ve Kristal Fe-Si Muadilleri ile Karşılaştırılması

Valik AHMETOV, Farhad MAMMADOV, Tebriz ŞAMILOV,
Aida İSAYEVA, and Günel ASKEROVA

Azerbaycan Mimarlık ve İnşaat Üniversitesi
isayeva_aida86@mail.ru

Amorf malzemeler, uzun menzilli atomik düzen eksikliği ile karakterize edilir ve benzersiz fiziksel ve mekanik özelliklerinden dolayı kristal muadillerine kıyasla umut verici alternatifler olarak kabul edilir. Fe-Si-C amorf şerit numunelerinin araştırılması, yumuşak manyetik özellikleri, enerji verimliliği ve mühendislik malzemeleri olarak çeşitli koşullardaki potansiyel uygulamaları hakkında önemli bilimsel bilgiler sağlar¹. Bu şeritlerin karakteristik manyetik özellikleri arasında düşük koersif kuvvet, yüksek doyum manyetizasyonu, düşük enerji kaybı vb. bulunur². Fe-Si-C amorf şeritlerinin manyetik özellikleri, elektrik motorlarında, transformatörlerde ve yüksek verimlilik gerektiren cihazlarda kullanımları için cazip hale getirir. Göreceli olarak yüksek Curie sıcaklıkları – manyetik özelliklerini geniş bir sıcaklık aralığında koruma yeteneği – çalışma sıcaklıklarının önemli ölçüde dalgalandığı uygulamalara uygun olmalarını sağlar.

Araştırmalar, Fe-Si-C amorf şerit numunelerinin karbon ve silikon içeriklerini ayarlayarak manyetik özelliklerin yapay olarak ayarlanabileceğini göstermiştir. Bu uygulama, özel alanlar için manyetik malzemelerin geliştirilmesinde benzersiz fırsatlar sunar.

Araştırmanın temel amacı, bu amorf şeritlerin manyetik özelliklerinin sıcaklık kararlılığı ve enerji verimliliği hakkında kristal Fe-Si muadilleriyle karşılaştırıldığında daha kapsamlı bilimsel veri sağlamaktır. Gelecek araştırma yönleri olarak, bu şeritlerin işleme teknolojisinin simüle edilmesi ve yeni örneklerin oluşturulması kapsamında çalışmaların derinleştirilmesi önerilmektedir.

Anahtar Kelimeler: Fe-Si-C amorf şerit, enerji verimliliği, elektrik mühendisliği malzemeleri.

Teşekkür Bölümü: Bu çalışma, Azerbaycan Bilim Fonu'nun (Azərbaycan Elm Fondu) mali desteğiyle gerçekleştirilmiştir. Hibe No: AEF-MQM-QA-2-2023-3(45)-05/01/1-M-01.

Referanslar

- [1] Abdullayev A.P. ve diğerleri. (2024). Fe-Si-C Amorf Şeritlerin Elde Edilmesine Yönelik Teknoloji Geliştirilmesi. *Universum: Teknik Bilimler: Bilimsel Dergi – Moskova*, 8(125), 56-60.
- [2] Abdullayev A.P. ve diğerleri. (2024). Fe₉₂Si₆C₂ ve Fe₉₃Si₆C₁ Amorf Şeritlerin Elektriksel Özelliklerinin İncelenmesi ve Bunların Kristal Fe-Si Muadilleri ile Karşılaştırılması. *Uluslararası Bilimsel Dergi "Bilim ve Eğitim Dünyasında"*, Astana, Kazakistan, 46-52.

Basicity of a Trigonometric System in Morrey-type Spaces

Yonca SEZER, Siavash AZİMİ

*Department of Mathematics, Faculty of Arts and Sciences, Yıldız Technical University,
Istanbul, 34210, Türkiye*

siavash.azimi@std.yildiz.edu.tr

We study the basis property of the trigonometric system $T = \{\cos(nx), x\sin((n+1)x)\}_{n \in \mathbb{N}}$ in the Morrey space $L_{p,\lambda}(0,2\pi)$, focusing on its application to a degenerate elliptic boundary value problem for $m \geq -2$. Morrey spaces are suitable for studying regularity in partial differential equations^{1,2}. Motivated by the application of basis systems to such problems³, we consider a separable subspace $\hat{L}_{p,\lambda}(0,2\pi)$ in which $C_0^\infty(0,2\pi)$ is dense.

We construct the system

$$t_0^+ = 1, t_n^+ = \cos(nx), t_n^- = x\sin(nx),$$

and its biorthogonal functionals $\phi_n^\pm$, and prove that the system is minimal and complete in $\hat{L}_{p,\lambda}(0,2\pi)$. Consequently, we obtain that the system T forms a basis in non-separable Banach space $\mathcal{L}_{p,\lambda}(0,2\pi)$. The system T has also the Riesz Property.

Keywords: Morrey space, Riesz basis, trigonometric system, degenerate elliptic equation.

References

- [1] Morrey, C. B. (1938). On the Solutions of Quasi-linear Elliptic Partial Differential Equations. Trans. Amer. Math. Soc., 43(1), 126-166.
- [2] Sawano, Y., Sugano, S., and Tanaka, H. (2020). Morrey Spaces II: With Applications to Harmonic Analysis. CRCPress.
- [3] Bilalov, B. T., Guliyeva, A. A. (2014). On the Basisness of the System of Eigenfunctions of a Degenerate Elliptic Operator. International Journal of Mathematics, 25(6), 1-10.

TURAN-FUNDAMENTAL SCIENCES SYMPOSIUM
TURAN-TEMEL BİLİMLER SEMPOZYUMU

Honorary Chair/Onursal Başkan: Prof. Dr. Aziz SANCAR

PHYSICS

Electron Screening Meets Space: D+D Fusion-Driven Propulsion Systems

Ali İhsan KILIÇ^{1,2}, Orkun HASEKİOĞLU², Kübra AYSU³, Muhammet Melih KAZANCI³, Hüseyin AYRANCI³, Furkan TAŞOĞLU³, Ahmet Taha KARABULUT³, and İbrahim YÖRÜKSOY³

¹*Eskişehir Osmangazi University, Department of Physics, Meşelik Kampüsü, Eskişehir, Türkiye*

²*TUBITAK, Research Institute of Fundamental Sciences, Barış Mah. Dr. Zeki Acar Cad. 41470
Gebze / Kocaeli, Türkiye*

³*Eskişehir Osmangazi University, Faculty of Engineering and Architecture, Meşelik Kampüsü,
Eskişehir, Türkiye*

turkfusiongroup@gmail.com

Material enhanced electron screening in Deuterium+Deuterium fusion reactions¹ has the potential to reduce the effective Coulomb barrier, significantly increasing fusion rates at low energies. This effect, observed in metallic lattices², may open a novel route toward compact nuclear micro propulsion systems for miniaturized spacecraft. In this study, we explore how the choice of deuterated target materials especially those with high electron densities and structured metallic meshes could influence reaction rates through screening induced enhancements as well as neutron suppression using unusual materials². We assess how such systems could generate usable thrust in space without requiring high-energy acceleration or bulky reactors. By tailoring the fusion environment at the atomic level, screening assisted D+D fusion may offer a clean, scalable, and isotopically safe propulsion method for CubeSat class missions, particularly suited for long-duration or attitude control applications. By enabling compact, fuel-efficient, and neutron-safe thrust, screening-enhanced D+D fusion systems hold great promise for future applications in space mining operations and sustained interplanetary travel.

Keywords: Electron screening effects, enhancement of fusion, space mining, interplanetary travel.

References

- [1] Kilic, A. I., Kustan, F.K. (2019). *Material-dependent $d + d$ Cross-section in Dense and Metallic Mesh Wire Environments at Very Low Energies*. Modern Physics Letters A, 34 (29), 1950234.
- [2] Czerski, K., Weissbach, D., Kilic, A. I., Ruprecht, G., Huke, A., Kaczmariski, M., Targosz-Ślęczka, N., and Maass, K. (2016). *Screening and Resonance Enhancements of the $2H(d, p)3H$ Reaction Yield in Metallic Environments*. Europhysics Letters, 113(2).

Constraining Key Resonances in the $^{40}\text{Ca}(p,\gamma)^{41}\text{Sc}$ Reaction via ANC-Assisted DWBA Analysis of Transfer Reactions and its Astrophysical Reaction Rate

Burak ÖZCAN¹, Ali İhsan KILIÇ^{1,2}, Hasret Gül ÖZTOP¹, Ebru Fadime KILIÇ KUŞTAN³, Derin ÇIVGIN¹, and Yasemin BULAT¹

¹*Eskişehir Osmangazi University, Department of Physics, Meşelik Campus, Eskişehir, Türkiye*

²*TUBITAK TBAE, Research Institute of Fundamental Sciences, Kocaeli, Türkiye*

³*Gazi University, Graduate School of Natural and applied Sciences, 06500 Teknikokullar,
Ankara, Türkiye
turkfusiongroup@gmail.com*

We present a novel investigation of the $^{40}\text{Ca}(p,\gamma)^{41}\text{Sc}$ reaction, which plays a critical role in nucleosynthesis processes in ONe novae. In particular, the low-energy resonances at $E_R=647$ keV and 1842 keV dominate the reaction rate within the stellar temperature range $0.2 < T_9 < 0.7$. Previous direct measurements have reported resonance strengths, but a deeper understanding of the nuclear structure and partial widths of these resonances remains essential. In this work, we revisit the $^{40}\text{Ca}(\alpha,3\text{He})^{41}\text{Ca}$ reaction at 40 MeV, for which angular distributions and spectroscopic factors have been reported for several known and newly observed states in Sc^{41} , including previously unreported levels at 3.356 MeV and 3.905 MeV excitation energy¹. By applying a DWBA framework² enhanced with experimentally extracted ANCs³, we aim to determine the proton and neutron partial widths Γ_p for the key resonances in Sc^{41} . Our approach allows consistent comparison of extracted widths with those inferred from gamma-ray resonance data, offering a nuclear structure-based explanation for their astrophysical relevance. This ANC-based method innovatively applies indirect techniques to constrain (p,γ) reaction rates and resonance properties, improving stellar model predictions⁴.

Keywords: DWBA analysis, ANC method, resonance widths, mirror transfer reactions.

References

- [1] Cohen, L. D., Holmgren, H. D. (1974). *Analog states in ^{41}Ca and ^{41}Sc from the $^{40}\text{Ca}(\alpha,3\text{He})$ and $^{40}\text{Ca}(\alpha,t)$ reactions*. Phys. Rev. C 9, 1023.
- [2] G. R. Satchler. (1983). *Direct Nuclear Reactions*. Oxford University Press.
- [3] Mukhamedzhanov, A. M. et al. (2001). *Asymptotic Normalization Coefficients, Spectroscopic Factors, and Direct Radiative Capture Rates*. Phys. Rev. C 63, 024612.
- [4] José, J., Hernanz, M. (1998). *Nucleosynthesis in Classical Novae: CO versus One White Dwarfs*. Astrophys. J. 494, 680.

Contaminated or Clean? **How Do Surface Conditions Shape Fusion Rates in Deuterated and Boron Targets?**

Ali İhsan KILIÇ^{1,3}, Derin ÇIVGIN², Ebru Fadime KILIÇ KUŞTAN⁴, and Orkun HASEKİOĞLU³

¹*Eskisehir Osmangazi University, Department of Physics, Meşelik Kampüsü, Eskişehir, Türkiye*

²*Eskisehir Osmangazi University, Graduate School of Natural and Applied Sciences, Department of Physics, Meşelik Kampüsü, Eskişehir, Türkiye*

³*TUBITAK, Research Institute of Fundamental Sciences, Barış Mah. Dr. Zeki Acar Cad. 41470 Gebze / Kocaeli, Türkiye*

⁴*Gazi University, Faculty of Sciences, Department of Physics, Emniyet Mahallesi Bandırma Caddesi No: 6/9, 06500, Yenimahalle/ANKARA, Türkiye*
turkfusiongroup@gmail.com

We aim to investigate surface effects on $p+^{11}\text{B}$ and Deuterium+Deuterium fusion reactions in the different metallic environments by combining theoretical and experimental approaches below 20 KeV. In the first phase, SRIM simulations will be used to model deuteron implantation profiles, energy loss, and surface layer formation in metallic targets. Special attention will be given to how oxide or carbon contaminations alter stopping powers and affect fusion yields via electron screening^{1,2}. In the second phase, experimental measurements will be performed at the Çekmece Nuclear Research and Training Center using deuterated metal targets under controlled conditions. The study seeks to quantify surface-induced modifications to fusion cross sections and to optimize target preparation for enhanced neutron, proton and alpha productions.

Keywords: D+D reactions, low energy nuclear reactions, surface effect, fusion.

References

- [1] Kilic, A. I., Kustan, F.K. (2019). Modern Physics Letters A, 34(29), 1950234.
- [2] Czerski, K., Weissbach, D., Kilic, A. I., Ruprecht, G., Huke, A., Kaczmariski, M., Targosz-Ślęczka, N., and Maass, K. (2016). Europhysics Letters, 113(2).

Fusion in the Deep: Pycnonuclear Reactions Under Extreme Stellar Conditions

Ali İhsan KILIÇ^{1,2}, Doğukan BARLAS³, Hasret Gül ÖZTOP³, Derin ÇİVGİN³,

Yasemin BULAT³, and Burak ÖZCAN¹

¹*Eskisehir Osmangazi University, Department of Physics, Meselik Campus, Eskişehir, Türkiye*

²*TUBITAK TBAE, Research Institute of Fundamental Sciences, Kocaeli, Türkiye*

³*Eskisehir Osmangazi University, Graduate School of Natural and Applied Sciences, Department of Physics, Meselik Campus, Eskişehir, Türkiye*

turkfusiongroup@gmail.com

Nuclear reaction rates can increase dramatically by several orders of magnitude, which would be under conditions found in dense and relatively cold astrophysical environments such as white dwarfs, brown dwarfs, and giant planets^{1,2}. Similar environments also occur during supernova explosions, where the ignition conditions are essential for cosmological modeling^{1,3}.

In this study, we introduce a new theoretical framework for calculating nuclear cross sections and reaction rates for light nuclei. This formulation incorporates not only nuclear structure effects but also material dependencies, magnetic field strengths, and the influence of the crystal lattice in dense metallic media². We explore how these factors modify the reaction dynamics and present the impact of our model on both the cross sections and reaction rates^{2,4}. This enhanced understanding may provide valuable insights into astrophysical nucleosynthesis and open up new avenues for energy generation using nuclear fusion reactions under solid-state or magnetically enhanced conditions^{1,2,4}.

Keywords: Reaction rate, astrophysical dense matter, white dwarfs, magnetic field effects.

References

- [1] Ichimaru, S. (1993). *Nuclear Fusion in Dense Plasmas*. Rev. Mod. Phys. 65, 255–299.
- [2] Kılıç, A. I. (2017). *Astrophysical Nuclear Reaction Rates in the Dense Metallic Environments*. APS Division of Nuclear Physics Hawaii Meeting, abstract id. DC.004.
- [3] Kondratyev, V. N. et al. (2020). *Magnetic and Thermal Effects in Neutrino Scattering in Hot and Dense Nuclear Matter*. Bull. Russ. Acad. Sci. Phys. 84, 962–967.
- [4] Haensel, P., Zdunik, J. L. (2008). *Models of Crustal Heating in Accreting Neutron Stars*. Astron. Astrophys. 480, 459–464.

Atmospheric Physics and Natural Transport Paths in the Atmosphere

Elif Seda ÇAĞATAY, Hasan TATLIPINAR

Yildiz Technical University, Esenler, Istanbul, Türkiye

escgty@gmail.com

The Earth's atmosphere can be divided into various layers according to the effects of gravity, density, temperature and pressure changes. The layer that directly concerns living things is the Troposphere, which is between 0-12 km thick.¹ The breathing of living things and photosynthesis events occur in this layer. 99 percent of water vapor and various aerosols, cloud formations are also found in this layer. Natural transport events in the atmosphere occur simply through diffusion, convection and air currents. Air currents are determined by many reasons such as irregular sea and land formations on the Earth's surface, regional temperature differences such as the poles and equator. In addition to air currents, atmospheric rivers and tornadoes are also transport paths. These transport paths, which occur seasonally, are important components of the Earth's climate formation.²⁻³ Thanks to these transport paths, water vapor, rain, desert dust, pollen, various viruses and chemical compounds are transported from one region to another on the Earth's surface and form the climate and life on Earth. In recent years, the impact of global warming on these natural air transport pathways in the atmosphere and the transportation of atmospheric pollution caused by industrialization through these pathways have caused great concern. For this reason, scientific studies are being conducted to determine, map and observe changes in the natural air transport pathways in the Earth's atmosphere.

Keywords: Atmospheric physics, atmospheric models, atmospheric river.

References

- [1] Nasa. (2025). <https://science.nasa.gov/earth/earth-atmosphere/earths-atmosphere-a-multi-layered-cake/>
- [2] Ambaum, M. (2010). Thermal Physics of the Atmosphere, John Wiley.
- [3] Francis, D. Fonseca, R., Bozkurt, D., Nelli N, and Guan, B. (2024). *Atmospheric River Rapids and Their Role in the Extreme Rainfall Event of April 2023 in the Middle East*, 51(12).

Renormalization Group Studies of Continuous-Spin Complex Systems

Erbil Can ARTUN^{1,2}, Ahmet Nihat BERKER^{2,1,3}

¹ TÜBİTAK Research Institute for Fundamental Sciences, Gebze, Kocaeli 41470, Türkiye

² Faculty of Engineering and Natural Sciences, Kadir Has University, Cibali, Istanbul 34083, Türkiye

³ Department of Physics, Massachusetts Institute of Technology, Cambridge,

Massachusetts 02139, USA

can.artun@tubitak.gov.tr

Renormalization group (RG) theory has illuminated critical phenomena across various systems, from discrete-spin models to continuous-spin systems with their distinctive topological transitions. While the XY model in two dimensions established the paradigm of algebraic order through the Berezinskii-Kosterlitz-Thouless transition¹, complex spin systems with multiple degrees of freedom remain less explored. Our study investigates the XY-Ashkin-Teller model, where each site hosts two continuously variable XY spins coupled via two-spin and four-spin interactions. Using position-space RG with bond-moving approximation, we identify a rich phase diagram containing multiple algebraically ordered phases—both ferromagnetic and antiferromagnetic in nature—for both direct and composite spin variables. Notably, these phases correspond to RG fixed surfaces rather than fixed points, characterized by factorizable double Fourier coefficients. This continuous spectrum of critical behavior reveals unprecedented complexity in the universality landscape of continuous-spin systems, providing new frameworks for classifying exotic phases of matter.

Keywords: Renormalization group theory, continuous-spin systems, XY-Ashkin-Teller model, critical phenomena, hierarchical lattices.

References

[1] Kosterlitz, J. M., Thouless, D. J. (1973). *Ordering, Metastability and Phase Transitions in Two-dimensional Systems*. J. Phys. C 6, 1181.

A Holistic Design Approach to Compact Neutron Generators: Adjustable Energies & Enhanced Yield with D-D Fusion Reactions

Hasret Gül ÖZTOP¹, Ali İhsan KILIÇ^{2,3}, Burak ÖZCAN², Ebru Fadime KILIÇ KUŞTAN⁴,

Derin Çİlgin¹, and Yasemin BULAT¹

¹ *Eskisehir Osmangazi University, Graduate School of Natural and Applied Sciences, Department of Physics, Meselik Campus, Eskisehir, Türkiye*

² *Eskisehir Osmangazi University, Department of Physics, Meselik Campus, Eskişehir, Türkiye*

³ *TUBITAK TBAE, Research Institute of Fundamental Sciences, Kocaeli, Türkiye*

⁴ *Gazi University, Graduate School of Natural and Applied Sciences, 06500, Teknikokullar, Ankara, Türkiye*

turkfusiongroup@gmail.com

Compact neutron generators offer a high-yield, portable neutron source by using D-D reactions without radioactive isotopes or reactors with turn on/off property reducing handling risks. High neutron flux improves precision by speeding up data acquisition and raising SNR in applications like security, medical imaging, and industry¹. We present a new generator design with three key features: a tunable RF ion source for real-time control of energy and current; an optimized² beamline for efficient ion extraction and focusing; and a meshed, metallic deuterated target engineered to enhance electron screening, lowering the Coulomb barrier and boosting reaction rates³. Our design improves control over ion-beam transport, and target performance in a compact setup. Simulations show that electron screening significantly enhances D-D fusion. By combining accelerator physics, material science, and nuclear engineering, this design offers a scalable solution for adjustable, high-flux neutron generation, awaiting experimental validation. New calculations shows that new designed generator promise more than 10⁸ neutron/s.

Keywords: Compact neutron generator, meshed-material target, RF ion-beam optimization, electron screening enhancement in D-D reaction, computational neutronics.

References

- [1] IAEA Radiation Technology Reports No1 (2012). Neutron Generators for Analytical Purposes.
- [2] Kilic, A. I., Kustan, F. K. (2019). *Material-dependent d + d Cross-section in Dense and Metallic Mesh Wire Environments at Very Low Energies*. Modern Physics Letters A, 34(29), 1950234.
- [3] Czerski, K., et. al. (2016). *Screening and Resonance Enhancements of the ²H(d, p)³H Reaction Yield in Metallic Environments*. Europhysics Letters, 113(2), 22001.

Re-thinking the ${}^8\text{Li}(\alpha, n){}^{11}\text{B}$ Reaction: Evidence for a Missing Resonance in ${}^{12}\text{B}$

Yasemin BULAT¹, Ali İhsan KILIÇ^{2,3}, Hasret Gül ÖZTOP¹, Derin ÇIVGIN¹, Doğukan BARLAS¹,
Burak ÖZCAN², and Ebru Fadime KILIÇ KUSTAN⁴

¹Eskisehir Osmangazi University, Graduate School of Natural and Applied Sciences Department of
Physics Meşelik Campus, Eskişehir, Türkiye

²Eskisehir Osmangazi University, Department of Physics, Meşelik Campus, Eskişehir, Türkiye

³TUBITAK TBAE, Research Institute of Fundamental Sciences, Kocaeli, Türkiye

⁴Gazi University, Graduate School of Natural and Applied Sciences, 06500,
Teknikokullar, Ankara, Türkiye

turkfusiongroup@gmail.com

The ${}^8\text{Li}(\alpha, n){}^{11}\text{B}$ reaction is highly interesting in both primordial nucleosynthesis scenarios, particularly in inhomogeneous Big Bang models¹ and in constraining the physical conditions relevant to the r-process². Despite its astrophysical importance, notable discrepancies exist between inclusive and exclusive measurements of the cross section, especially at center-of-mass energies below 3 MeV³. To address this issue, we apply the R-matrix formalism for extracting nuclear structure of compound systems such as ${}^{12}\text{B}$ and for extrapolating the astrophysical S-factor to the relevant Gamow window. In this work, we present R-matrix⁴ calculations for the ${}^8\text{Li}(\alpha, n){}^{11}\text{B}$ reaction, providing results for both the reaction rates and the partial S-factors. By incorporating direct reaction contributions into our model, we are able to reproduce the experimental data from inclusive measurements with improved consistency. We are investigating a possible new resonance in the nuclear structure of ${}^{12}\text{B}$, including levels with known energies but uncertain spin-parity that may impact reaction rates. Our extended R-matrix analysis aims to resolve experimental discrepancies and clarify the ${}^{12}\text{B}$ levels.

Keywords: Big bang nucleosynthesis, inhomogeneous big bang model, astrophysical r-process.

References

- [1] Kajino, T. (1995). *Inhomogeneous Big-bang Model, Revived, and Evolution of the Light Elements in Cosmic Rays*. Nucl. Phys. A, 588, 339.
- [2] Sasaqui, T. et al. (2005). *Sensitivity of r-Process Nucleosynthesis to Light-Element Nuclear Reactions*. Astrophys.J, 634, 1173.
- [3] Kilic, A. I. (2023). *The Investigation of the ${}^8\text{Li}(\alpha, n){}^{11}\text{B}$ Cross Section at Low Energy by a Distorted Wave Born Approximation Model*. Physics of Particles and Nuclei Letters, 20(6), 1335–1341.
- [4] Descouvemont, P., Baye, D. (2010). The R Matrix Theory. Rep. Prog. Phys., 73, 036301.

Aktif Matrisli Hibrit Kompozitlerin Piezoelektrik Özellikleri

Rena CEBRAYILOVA

*Azerbaycan Mimarlık ve İnşaat Üniversitesi, Bakü, Ayna Sultanova Cad., 11
ranacabrayilova@sabah.edu.az*

Aktif kompozit malzemeler, piropiezo özellikleri sayesinde çoğunlukla radyo mühendisliği, elektronik ve optoelektronik alanlarında kullanılmaktadır. Bu malzemeler; uzay araçlarının radyasyondan korunması, denizaltı görüntüleme, sismik ve jeolojik araştırmalar, alternatif enerji kaynakları ile tıbbi ve biyolojik alanlar gibi uygulamalarda yer almaktadır.¹⁻² 0-3 ve 1-3 tipi piezoelektrik kompozitler, farklı boyutlardaki seramik biçimli piezoelektrik parçacıkların (örneğin PZT-5H, PZT-5A) polimer matrisler ile rastgele karıştırılmasıyla elde edilir. Bu aktif matrisli kompozitler özel şekillerde üretilebilir ve piezoelektrik özelliklerini korurlar.² Hibrit piezoelektrik kompozitler (HPK), en az iki gruptan oluşan yapılardır. İlk grupta düşük veya yüksek yoğunluklu polimerler (PE—polietilen, PVDF—poliviniliden florür, PP—polipropilen) ile nano boyutlu metal oksitler (BaTiO₃, SiO₂, TiO₂) bulunur. İkinci grupta ise yine aynı polimerler ve mikro boyutta tetragonal yapılı PZT (kurşun zirkon titanat) seramikleri yer alır. Birinci gruptaki kompozitler nanokompozit, ikinci gruptakiler ise mikro piezo kompozit olarak adlandırılır. Hibrit piezoelektrik yapı, bu iki grubun entegrasyonu sonucu elde edilir. PZT ailesindeki bileşim çeşitliliğinin piezo-piro ve elektriksel malzemelerin gelişimi için yeterli olmadığı söylenebilir. Bu tür malzemelerde sadece piezo-piroelektrik özellikler değil, aynı zamanda dielektrik sabitleri de artmaktadır. Ancak bu durum malzemenin verimliliğine katkı sağlamamaktadır. Bu nedenle piezo-piroelektrik ve elektroseramik malzemelerin karakteristik verimliliğinde kayda değer bir artış elde edilememektedir. Bu çalışmada, deşarj plazma sistemleri ile elde edilen hibrit piezokompozitlerin ve mikro boyutlu piezokompozitlerin elektromekanik özellikleri araştırılmıştır. Deşarj plazma sistemi ile elde edilen hibrit yapılı piezokompozitler, 5 Hz-40 kHz aralığındaki akustik uygulamalarda kullanılabilir. Uygulama alanlarına özel olarak şekillendirilip farklı kalıplara yerleştirilen hibrit yapılı piezokompozitler, piezo sensörlere dönüştürülebilir.

Anahtar Kelimeler: Hibrit kompozitler, piezokompozitler, deşarj plazma, nanokompozitler.

Referanslar

- [1] Kerimov, M. K., Kurbanov, M. A., Mekhtili, A. A., Aliev, G. G., Sultanakhmedova, I. S., Tatardar, F. N., Yusifova, U. V., Kulieva, G. K., and Yakhyaev, F. F. (2011). *Piezoelectrics Based on a Hybrid of Piezoelectric Matrix Nano and Microcomposites*. Tech. Phys., 56, 1187–1194.
- [2] Rus, G., García-Sánchez, F., Sáez, A., and Gallego, R. (2010). Damage Detection in Piezoceramics via BEM. Key Eng. Mater., 417, 381–384.

Amorf Şerit Üretiminde Kalite Kontrol İçin Vickers Sertlik Ölçümlerinde Standart Sapma Potansiyeli

Farhad MAMMADOV, TebriZ ŞAMİLOV, Sedaket MUSAYEVA, Nurlan REFİYEV,
and Nuranə QASIMOVA

Azərbaycan Mimarlık ve İnşaat Üniversitesi
nrafiyev@mail.ru

Vickers sertlik ölçümleri, amorf malzemelerin mekanik özelliklerini değerlendirmek için temel yöntemlerden biridir. Sertlik ölçümlerinin standart sapması hesaplanarak numuneler arasındaki değişkenlik tespit edilebilir. Düşük bir standart sapma, numunelerin aynı koşullar altında hazırlandığını ve benzer yapısal veya mekanik özelliklere sahip olduğunu gösterir. Amorf malzemelerdeki rastgele ancak düzgün atomik düzen, mekanik özelliklerin birbirine yakın olmasını sağlar¹. Düşük standart sapma, veri değişkenliğinin minimum olduğunu ve homojenliği doğrular. Özellikle Fe-Si-C amorf malzemelerde, dislokasyonların ve tane sınırlarının olmaması deformasyon ve aşınmaya karşı dayanıklılığı artırır².

Bu çalışmada, TQ-150 mikrosertlik cihazı kullanılarak 10 gf yük altında Vickers sertlik ölçümleri gerçekleştirilmiştir. Her numune için ardışık 10 ölçüm yapılmış ve numuneler arasındaki homojenliği değerlendirmek amacıyla ortalama ve standart sapma gibi istatistiksel göstergeler hesaplanmıştır. Düşük standart sapma ($\leq 1.5\%$) ile, numunelerin yüksek homojenliğini doğrulanmıştır.

Araştırma sonuçları, malzemenin çok yüksek Vickers sertlik değerleri, düşük standart sapma değerleri ve homojen bir yapıya sahip olduğunu göstermiştir. Bu nedenle, düşük standart sapma değerleri, şerit üretiminde süreç istikrarını değerlendirmek için uygulanabilir. Standart sapma, amorf malzemelerde çeşitli mekanik özelliklerin optimize edilmesinde kritik bir rol oynayabilir. Bu yaklaşım, elektrik motorları, transformatörler ve manyetik alanlarda çalışan cihazlarda kaliteyi sürdürmek için değerli olabilir.

Anahtar Kelimeler: Fe-Si-C amorf şerit, kalite kontrol, istatistiksel analiz.

Teşekkür Bölümü: Bu çalışma, Azerbaycan Bilim Fonu'nun (Azərbaycan Elm Fondu) mali desteğiyle gerçekleştirilmiştir. Hibe No: AEF-MQM-QA-2-2023-3(45)-05/01/1-M-01.

Referanslar

- [1] Abdullayev A.P ve diğerleri (2024). *Fe-Si-C Amorf Şeritlerin Elde Edilmesine Yönelik Teknoloji Geliştirilmesi*. Universum: Teknik Bilimler: Bilimsel Dergi – Moskova, 8(125), 56-60.
- [2] Abdullayev A.P ve diğerleri (2024). *Fe₉₂Si₆C₂ ve Fe₉₃Si₆C₁ Amorf Şeritlerin Elektriksel Özelliklerinin İncelenmesi ve Bunların Kristal Fe-Si Muadilleri ile Karşılaştırılması*. Uluslararası Bilimsel Dergi "Bilim ve Eğitim Dünyasında," Astana, Kazakistan, 46-52.

Dielectric Performance and Potential Applications of HDPE/nano MgO Composites

Aida AZIZOVA¹, Vugar KHUDAVERDIEV¹, Akif SALEHOV¹, and Shafiqa SHUKUROVA²

¹*Institute of Radiation Problems of the Ministry of Science and Education of the Republic of Azerbaijan, Baku, Azerbaijan*

²*Azerbaijan State Oil and Industry University, Baku, Azerbaijan*

aidaazizova889@gmail.com

Metal oxide-filled polymer composites represent a significant advancement in insulation materials, offering enhanced physical properties and novel characteristics. This study investigates the dielectric properties of high-density polyethylene (HDPE) composites with varying concentrations of magnesium oxide (MgO) nanoparticles, focusing on their temperature-dependent and frequency-dependent behavior.¹ Composite samples were prepared by mechanically mixing HDPE with nano-MgO at different volume fractions (0%, 10%, 20%, and 30%), followed by hydraulic pressing at 15 MPa and 130°C for 5 minutes. Dielectric measurements were conducted using a "sandwich"-type measurement chamber and an E7-20 impedance analyser across frequencies of 25 Hz to 1 MHz and temperatures of 297–383 K. Results revealed distinct differences in dielectric constant between pure HDPE and HDPE/nano-MgO composites. The 30% nano-MgO composite exhibited the highest dielectric constant, showing a 45% increase compared to pure HDPE. The loss tangent showed positive correlation with temperature, with a notable conductivity increase at 353 K in the 30% nano-MgO sample. This behavior is attributed to enhanced ionic conductivity, polymer crystalline phase transformation, and increased inter-particle spacing of nano-MgO.² The study demonstrates that nano-MgO concentration significantly influences composite dielectric properties, suggesting potential applications in advanced insulation materials and electronic devices.

Keywords: Metal oxide-filled polymer composites, dielectric properties.

References

- [1] Shaojie, W., Zhen, L., Jiajie, L., Jun, H., Naisheng, J., Jinliangü, H., and Qi, L. (2022). *Polymer Nanocomposite Dielectrics: Understanding the Matrix/Particle Interface*. ACS Nano, 16(9), 13612-13656.
- [2] Tan, D. Q. (2020). *The Search for Enhanced Dielectric Strength of Polymer-based Dielectrics: A Focused Review on Polymer Nanocomposites*. Journal of Applied Polymer Science, 137(33), e49379.

Electrical Conductivity of Low-Dimensional Electron Gas in Asymmetric Quantum Well Scattering in Ionized Impurities in the Absence of Screening

Huseynagha HUSEYNOV¹, Mehdi MAHMUDOV²

¹*Azerbaijan University of Architecture and Construction, A.Sultanova st., 11, AZ1073, Baku, Azerbaijan*

²*Baku State University, Z.Khalilov st., 23, AZ1148, Baku, Azerbaijan*

huseynov.h.i@mail.ru

Currently the development of nanotechnology has made it possible to grow various types of low-size systems, such as thin films, quantum wells and others¹. Asymmetric quantum wells, unlike symmetric ones, have some very interesting properties. Resistive effects in a semi-parabolic quantum well were studied mainly in phonon scattering, but due to the difficulty of calculating the relaxation time in the case of scattering on impurity ions, they are weakly studied²⁻³.

This work is devoted to the study of the electrical conductivity of a two-dimensional electron gas in a semi-parabolic quantum well with scattering of conduction electrons on impurity ions. The work considers a semi-parabolic quantum well with potential $V(z)$ ².

Calculations show that the electrical conductivity of an electron gas in a semi-parabolic quantum well increases significantly with increasing potential and with decreasing quantum well width. This growth is due to the effect of electron localization and reduced electron scattering. The electron mobility during scattering on impurity ions becomes an order of magnitude greater than during scattering on phonons².

Keywords: Electrical conductivity, two-dimensional electron gas, semi-parabolic quantum well.

References

- [1] Niculescu, E. C., Eseanu, N. (2011). *Interband Absorption in Square and Semiparabolic Near – surface Quantum Wells Under Intense Laser Field*. The European Physical Journal B, 79, 313-319.
- [2] Figarova, S. R., Huseynov H. I., Figarov, V. R. (2021). *Thermoelectric Power Hysteresis in Semi-parabolic Quantum Well*. Thin Solid Films, 721, 138554.
- [3] Figarova, S. R., Huseynov, H. I. (2024). *Conductivity of a Two-dimensional Electron Gas in a Semi-parabolic Well Under Electron-impurity Scattering*. International Scientific Journal “In the World of Science and Education”, Aktau, Kazakhstan, 195-200.

Influence of Gamma Radiation on the Anisotropy of Electrical Conductivity in TlGaTe₂ Crystals

Adil ABDULLAYEV¹, Dilara GAFAROVA¹, İlaha ABDULLAYEVA^{1,2}, and Fatima BABAYEVA¹

¹Azerbaijan University of Architecture and Construction, Baku, Ayna Sultanova St., 11

²Institute of Radiation Problems, Ministry of Science and Education Republic of Azerbaijan, Baku
dgafarova14@gmail.com

The semiconductor crystals TlGaTe₂ belong to the class of compounds from group A³B³C₂⁶, crystallizing in the tetragonal space group D_{4h}¹⁸. The presence of a gap in the density of states, as well as the highly anisotropic chain-like structure in crystals of this class, suggests the possibility of peculiarities in the mechanism of electrical conductivity¹.

The temperature dependence of the electrical conductivity of the chain-like semiconductor crystal TlGaTe₂ was studied along both the chains and perpendicular to them. It was shown that in the region of the activation mechanism of conductivity, the anisotropy coefficient K_σ: it becomes > 1 (a value of K_σ = 1 indicates no anisotropy), increases with temperature, and at 250 K reaches a value of approximately 4, which is higher than the anisotropy coefficient 1/ K_σ ≈ 3.3 at low temperatures.

This work presents the results of studies on the influence of gamma radiation on the electrical conductivity of TlGaTe₂ single crystals.

The experimental results show that induced radiation defects lead to the formation of impurity energy levels in the forbidden band of the crystal. The thermal filling of these levels occurs at a lower temperature compared to the unirradiated compound. This temperature range is also characterized by the thermal filling of trap centers, and localized charged impurities become neutral. Defects of ionization type (charged defects) play a dominant role in these processes, which arise as a result of gamma irradiation. In our opinion, the most likely mechanism for forming radiation defects in TlGaTe₂ is the repeated ionization of the anionic sublattice.

Analyzing the obtained data, it can be concluded that gamma irradiation expands the temperature range for the existence of the thermally activated conductivity component. It is also shown that the temperature for the thermal filling of trap centers shifts toward lower temperatures.

Keywords: Semiconductor, radiation, trap center.

References

[1] Abdullayev, A. P., Rzayev, R. M., Naghiyev, T. G., Mammadova, J. P., Aliyev, S. S., and Musazade, I.V. (2023). *Determination Shape of the Potential Well of Traps in TlGaTe₂ Crystals*. International Journal of Modern Physics B, 37(28), 2350248, (9 pages). <https://doi.org/10.1142/S021797922350248X>.

Role of the P_z^{1+} State in Ferromagnetism and Topology of $MnBi_2Te_4$

Elbek Javokhir KESKİNOĞLU

Department of Physics, The Hong Kong University of Science & Technology

ejkeskinoglu@connect.ust.hk

$MnBi_2Te_4$ and its layered derivatives exhibit tunable magnetic and topological behaviors, where interlayer spacing^{1,2} and $MnBi_2Te_4/Bi_2Te_3$ ratios govern antiferromagnetic (AFM) and ferromagnetic (FM) orderings^{3,4,5}.

This study identifies the P_z^{1+} state as pivotal in driving FM behavior and enabling non-trivial topology. A tight-binding model, as shown in prior studies^{6,7}, reveals that the M_α/M_β ratio critically controls band inversion and quantum anomalous Hall (QAH) states. At a ratio of 1, band inversion occurs, but even 2-3% deviations yield trivial phases. The P_z^{1+} state enhances spin-dependent band modifications, stabilizing both FM order and topological phases. These results advance strategies for engineering magneto-topological materials for quantum technologies.

Keywords: Ferromagnetism, topological properties, $MnBi_2Te_4$, Tight-Binding Hamiltonian.

References

- [1] Li, J., Xu, Y. (2019). *Intrinsic Magnetic Topological Insulators in van der Waals layered $MnBi_2Te_4$ -Family Materials*. Science Advances, 5, 6.
- [2] Otrokov, M. M., Chulkov, E. V. (2019). Unique Thickness-dependent Properties of the van der waals Interlayer Antiferromagnet $MnBi_2Te_4$ Films. Phys. Rev. Lett., 122, 107202.
- [3] Wu, J., Liu, F., and Hosono, (2019). *H. Natural van der waals Heterostructural Single Crystals with Both Magnetic and Topological Properties*. Science Advances, 5(11), eaax9989.
- [4] Hu, C., Ni, N. (2020). *Realization of an Intrinsic Ferromagnetic Topological State in $MnBi_8Te_{13}$* . Science Advances, 6(30), eaba4275.
- [5] Ovchinnikov, D., Xu, X. (2021). *Intertwined Topological and Magnetic Orders in Atomically Thin Chern Insulator $MnBi_2Te_4$* . Nano Lett., 21(6), 2544–2550.
- [6] Atanov, O., Law, K. T., and Lortz, R. (2024). Proximity-induced Quasi-one-dimensional Superconducting Quantum Anomalous Hall State. Cell Reports Physical Science, 5(1), 101762.
- [7] Chen, C. Z., Law, K. T. (2018). Quasi-one-dimensional Quantum Anomalous Hall Systems as New Platforms for Scalable Topological Quantum Computation. Phys. Rev. B, 97, 104504.

Yapay Zeka Destekli Moleküler Sistemlerin Kuantum Simülasyonu: H₂ Molekülü İçin Qiskit ve OpenFermion ile VQE Uygulaması ve Hata Analizi

Büşra ŞAHİNAL

Ankara, Türkiye, 06450

busrasahinal@gmail.com

Bu çalışma, yapay zeka tabanlı kuantum algoritmalarının moleküler sistemlerin enerji hesaplamalarındaki potansiyelini araştırıyor. H₂ molekülünün temel enerji seviyesini bulmak için Qiskit ve OpenFermion kütüphanelerini kullanarak Varyasyonel Kuantum Özdeğer Çözücü (VQE) algoritması uygulandı. Farklı optimizasyon yöntemleri olan COBYLA, SLSQP ve SPSA ile simülasyonlar yapıldı ve bu sonuçlar klasik Hartree-Fock yöntemiyle karşılaştırıldı.¹⁻⁴ Elde edilen verilere dayanarak hata analizi yapıldı; kuantum simülasyonlarının moleküler kimyada nasıl fayda sağladığı ve kısıtlamaları tartışıldı. Çalışma, kuantum ve yapay zeka teknolojilerinin moleküler modelleme konusundaki önemini vurgulamaktadır.

Anahtar Kelimeler: Kuantum simülasyonu, yapay zeka, VQE, moleküler sistemler, qiskit.

Referanslar

- [1] McClean, J. R., Romero, J., Babbush, R., and Aspuru-Guzik, A. (2016). *The Theory of Variational Hybrid Quantum-classical Algorithms*. New Journal of Physics, 18(2), 023023.
- [2] McArdle, S., Endo, S., Aspuru-Guzik, A., Benjamin, S. C., and Yuan, X. (2020). *Quantum Computational Chemistry*. Reviews of Modern Physics, 92(1), 015003.
- [3] Qiskit Documentation. (2024). IBM Quantum Experience. <https://qiskit.org/documentation/>
- [4] OpenFermion. (2024). Open-source Software for Quantum Simulation of Chemistry and Materials. <https://github.com/quantumlib/OpenFermion>.

TURAN-FUNDAMENTAL SCIENCES SYMPOSIUM
TURAN-TEMEL BİLİMLER SEMPOZYUMU

Honorary Chair/Onursal Başkan: Prof. Dr. Aziz SANCAR

CHEMISTRY

Influence of Water Quality on The Electrochemical Behavior of a Carbon Steel Used in The Combustion Chamber of Boilers in a Petroleum Complex

Nadia HAMMOUDA¹, Abdel Ali BOUKRIKA¹, Mehdi GABLI¹, and Mostefa BACCOUCHE²

¹*Department of Chemistry, Faculty of Sciences, University of 20 Août 1955, Route d'El Hadaiek, Skikda 21000, Algeria*

²*Faculty of Technology, Badji Mokhtar-Annaba University, Annaba 23000, Algeria
hammoudanad@yahoo.com*

In this paper, industrial boilers in a petrochemical industry of Skikda (Algeria) are used as the research object. The aim of this work is based on the study of the efficiency of chemical treatments by chemical analysis (pH, TAC), and the study of the influence of these different parameters on the corrosion behavior of C-1026 carbon steel used in combustion chamber tubes in an industrial high pressure boiler¹, using stationary electrochemical methods (free potential: E-t, polarization curves: E-i, the Tafel rights and the R_p). The results obtained, shows that C-1026 carbon steel has good behavior in steam water and feed water. The pH values are clearly basic and almost stable, boiler water TAC values are quite high²⁻³. The treatment of the feed water and the boiler water with inhibitors has reduced the values of the corrosion current density (i_{corr}) and there is a remarkable displacement of the corrosion potential (E_{corr}) to higher values.

Keywords: Carbon steel, corrosion, feed water, boiler water.

References

- [1] Pincus, L. I. (1962). *Practical Boiler Water Treatment: Including Air-conditioning Systems*. McGraw-Hill, New York.
- [2] Boufelfel, M. (2022). *Etude Préventive sur la Corrosion des Chaudières au Niveau du Complexe RA1K*, Mémoire de fin d'étude, Université 8 Mai 1945 Guelma, Algérie.
- [3] Abbas, W. (2020). *Contrôle de Qualité de L'eau D'alimentation et de la Chaudière BAB-COCK de L'unité Centrale et Utilité II FERTIAL- Annaba*, mémoire de fin d'étude, Algérie.

Inhibitive Effect of an Organic Compound on the Corrosion of Carbon Steel in Demineralized Water

Nadia HAMMOUDA¹, Chima LAKEHAL¹, Esma DAOUAD¹, and Mostefa BACCOUCHE²

¹*Department of Chemistry, Faculty of Sciences, University of 20 Août 1955, Route d'El Hadaiek, Skikda 21000, Algeria*

²*Faculty of Technology, Badji Mokhtar-Annaba University, Annaba 23000, Algeria*

hammoudanad@yahoo.com

The study of corrosion inhibition of C-1026 carbon steel in demineralized water by morpholine was carried out using different techniques: stationary electrochemical methods (free potential: E-t, polarization curves: E-i, the Tafel rights and the Rp) and non-stationary electrochemical methods (EIS)¹. The results obtained show that the addition of morpholine to our medium considerably slows down the corrosion process, this has been confirmed by the noble values of the corrosion potential (E_{corr}) and the low values of the corrosion current (i_{corr})².

The immersion time only decreases the inhibitory efficiency. The inhibitory action of our organic compound is related to the formation (by adsorption) of a more or less continuous barrier, but of finite thickness, which prevents access of the solution to the metal. The characterization of the surface was carried out using an optical microscope³.

Keywords: Carbon steel, demineralized water, corrosion, inhibition.

References

- [1] Imran, M. (2014). *Effect of Corrosion on Heat Transfer Through Boiler Tube and Estimating Overheating*. International Journal of Advanced Mechanical Engineering, 4, 629-638.
- [2] Luo, H., Guan, Y. C., and Han, K. N. (1998). *Corrosion Inhibition of a Mild Steel by Aniline and Alkylamines in Acidic Solutions*, Corrosion science, 54, 721.
- [3] Mahlia, T. M. I., Abdulmuin, M. Z., Alamsyah, T. M. I., and Mukhlisshien, D. (2002). *Dynamic Modelling and Simulation of a Palm Wastes Boiler*, Renewable Energy, 28, 1235-1256.

Synthesis and Photopolymerization of Imidazole-based Photoinitiator

Alper OĞUZ, Sevnur KESKİN DOĞRUYOL

Yildiz Technical University, Department of Chemistry, 34220, Istanbul, Türkiye

alper.oguz@std.yildiz.edu.tr

Photo curing technology, as an environment-friendly technology, has been widely applied to chemical, machinery, electronics, light industry, communications, and other industries. The most important component of photo curing systems is photoinitiator, which can absorb UV light and therefore generate active species, leading to the photopolymerization of unsaturated monomers or resins. Most research has focused on type 1 initiators that undergo α -cleavage upon excitation and produce two radicals. In type 2 photoinitiators, triplet excited states react with hydrogen donors to form an initiator radical¹. Imidazoles and their scaffold are an extraordinarily essential class of nitrogen bearing azole heterocyclic compounds. They have different place in wide area of organic synthesis, which can be utilized in a variety of applications in diverse fields including agriculture, medicine, polymer and various industries².

In the presented study, imidazole based 2-(1H-imidazol-1-yl)-1-(naphthalen-2-yl) ethan-1-one (IPY) was synthesized and characterized as a new photoinitiator in the light of α -amino acetophenone derivatives. The influence of IPY concentration and the addition of the co-initiator N-methyldiethanolamine (MDEA) on methyl methacrylate (MMA) polymerization were also investigated.

Keywords: Photopolymerization, photoinitiator, imidazole.

References

- [1] Altikatoglu Yapaoz, M., Dogruyol, S. (2024). *Pyrazole-Based Photoinitiator with Antibacterial Activity*. Polimery, 69, 3, 167-172.
- [2] Qiu, J., Wei, J. (2014). *Novel Imidazole-containing Thioxanthone Derivative for Photo Polymerization*. Journal of Polymer Research, 21, 546.

Comparative Study of the Performance of Activated Carbon Obtained By Activation and Carbonization of Adsorbent Material Obtained From Avocado Peel with Different Chemicals in Removing Textile Dyes

Elif Zeynep ÇATMA, Türkan BÖRKLÜ BUDAK

Yildiz Technical University, Department of Chemistry, 34220, Istanbul, Türkiye
zeynep.catma@std.yildiz.edu.tr

Basic Yellow 28 (BY28) is a cationic dye with high water solubility and is one of the persistent pollutants frequently encountered in textile industry wastewater. Due to its cationic nature, it cannot be effectively removed by conventional treatment methods, leading to increased environmental pollution.¹ In this study, the adsorption method was employed for the removal of BY28 from aqueous solutions, utilizing waste avocado peel biomass as a low-cost and sustainable solution.

In addition to raw avocado peels, various types of activated carbon produced by chemical activation and carbonization processes were also used in comparative experiments. The porous structure and high surface area of activated carbon enhance the adsorption efficiency by allowing BY28 molecules to effectively adhere to the surface.² Experimental results showed that both raw biomass and chemically activated carbons exhibited varying performances in BY28 removal, with some of them identified as among the most effective adsorbents.

Avocado peels, classified as agricultural waste, are targeted to be used as environmentally friendly adsorbent alternatives. In this context, the study aims to provide sustainable solutions for the treatment of textile wastewater, offering both economic and ecological benefits.

Keywords: Basic Yellow 28, adsorption, activated carbon, textile wastewater, sustainable treatment.

References

- [1] Yagub, M. T., Sen, T. K., Afroze, S., and Ang, H. M. (2014). *Dye and its Removal from Aqueous Solution by Adsorption: A Review*. Advances in Colloid and Interface Science, 209, 172-184.
- [2] Babel, S., Kurniawan, T. A. (2003). *Low-cost Adsorbents for Heavy Metals Uptake from Contaminated Water: A Review*. Journal of Hazardous Materials, 97(1-3), 219-243.

Treatment of Synthetic Textile Wastewater Containing Both Basic Yellow 28 and Basic Blue 3 Dyes by Electrochemical Method

Halide İrem KOÇ, Ali Kaan YORGANCI, and Türkan BÖRKLÜ BUDAK

Yildiz Technical University, Department of Chemistry, 34220, Istanbul, Türkiye

halide.koc@std.yildiz.edu.tr

Nowadays, the treatment of textile wastewater has become increasingly important due to growing concerns about environmental pollution. Toxic dyes such as Basic Yellow 28 and Basic Blue 3 pose serious risks to ecosystems and human health. This study aims to treat synthetic textile wastewater containing both dyes simultaneously using an electrochemical method. The electrochemical approach offers advantages such as low energy consumption, high efficiency, and sustainability.¹ Using pure aluminum and graphite electrodes, direct current was applied to initiate the treatment. Treatment efficiency was determined via UV-spectrophotometry,² while parameters like pH and conductivity were monitored. This study aims to contribute to the literature by simultaneously removing two toxic dyes using an environmentally friendly and innovative technique³.

Keywords: Basic Yellow 28, Basic Blue 3, electrochemical method, wastewater treatment.

References

- [1] Martínez-Huitle, C. A., Brillas, E. (2009). *Decontamination of Wastewaters Containing Synthetic Organic Dyes by Electrochemical Methods: A General Review*. Applied Catalysis B: Environmental, 87(3–4), 105–145.
- [2] Chen, Y., Zhang, Y., Liu, C., Lu, A., and Zhang, W. (2012). *Photodegradation of Malachite Green by Nanostructured Bi₂WO₆ Visible Light-Induced Photocatalyst*. International Journal of Photoenergy, 510158.
- [3] Koparal, A. S., Yavuz, Y., Gürel, C., and Öğütveren, Ü. B. (2007). *Electrochemical Degradation and Toxicity Reduction of C.I. Basic Red 29 Solution and Textile Wastewater by Using Diamond Anode*. Journal of Hazardous Materials, 145(1–2), 100–108.

The Synthesis And Mesomorphic Investigation of New Chiral Calamitic Liquid Crystal Derived From Cyanobiphenyl Core

İnci Tuğba ÖRDEKBAY, Hale OCAK GÜMRÜKÇÜ

Yıldız Technical University, Department of Chemistry, 34220, Istanbul, Türkiye

tugba.ordekbay@std.yildiz.edu.tr

Liquid crystals are an intermediate phase between liquid and solid states, exhibiting fluidity like liquids while maintaining a certain degree of order and directed physical properties like solids. Calamitic liquid crystals are the most prominent and extensively studied class of LC¹, and these structures with aromatic ring-based cores and various terminal groups, have a structural stability through molecular interactions ensured by their polar end groups. When chiral units are introduced into the calamitic molecular structure, the resulting chiral liquid crystals exhibit strong optical properties and high sensitivity to environmental stimuli, making them highly suitable for applications in display technologies and sensors^{2,3}. In this study, the synthesis of a novel liquid crystal molecule based on cyanobiphenyl core and carrying an (*S*)-2-*n*-dodecyloxypropoxy chiral terminal chain was successfully carried out. The molecular structure was characterized by spectroscopic techniques such as FTIR, ¹H-NMR, ¹³C-NMR and HR-MS. Liquid crystal properties and thermal characteristics were investigated by using polarized optic microscope (POM) and differential scanning calorimeter (DSC). The new molecule exhibited an enantiotropic SmA mesophase (SmA) characterized by a fan-shaped texture in a wide temperature range.

Keywords: Liquid crystals, molecular chirality, (*S*)-2-*n*-dodecyloxypropyl chain.

Acknowledgement: This work was supported by TUBITAK 2209-A Research Project Support Program with the project number 1919B012311947.

References

- [1] Dierking, I. (2014). *Chiral Liquid Crystals: Structures, Phases, Effects*. Symmetry, 6(2), 444-472.
- [2] Dmochowska E., Herman J., Czerwiński M., Stulov S., Bubnov A., and Kula P. (2021). *Self-Assembling Behaviour of Chiral Calamitic Monoacrylates Targeted for Polymer Stabilisation of Polar Smectic Phases in Chiral Liquid Crystals*. Journal of Molecular Liquids, 331, 115723.
- [3] Srinivasa H. T. (2019). Chiral Liquid Crystals: Synthesis and Characterization for Thermal and Mesomorphic Properties. Molecular Crystals and Liquid Crystals, 680(1), 10-19.

One-Step Preparation of Composite Phase Change Films via Photoinitiated Oil-in-Oil Emulsion Templating

Meltem SÖZBİR¹, Hatice Hande MERT², Mehmet Selçuk MERT³, Burcu KEKEVİ⁴, and Emine Hilal MERT⁵

¹*Institute of Graduate Education, Department of Polymer Materials Engineering,
Yalova University, Yalova, Türkiye*

²*Faculty of Engineering, Department of Chemical Engineering, Yalova University, Yalova, Türkiye*

³*Faculty of Engineering, Department of Energy Systems Engineering,
Yalova University, Yalova, Türkiye*

⁴*Materials and Material Processing Technologies Department, Yalova University,
Yalova Community College, Yalova, Türkiye*

⁵*Faculty of Engineering, Polymer Materials Engineering,
Yalova University, Yalova, Türkiye*

meltem.sozbir@gmail.com

Latent heat storage technology, which offers an important solution for the sustainable use of energy, has been met with increasing interest in recent years and stands out as one of the promising methods in the field of energy storage. In this context, phase change materials (PCMs) are preferred in many thermal energy applications due to their high energy storage properties in a narrow temperature range.¹

Herein, composite phase change material (PCM) films were prepared via a photo-initiated, oil-in-oil (o/o) emulsion templating approach. SiO₂ nanoparticle loaded o/o emulsions were prepared by dispersing HD in a continuous monomer phase. The amount of nanoparticle loading was varied to investigate the influence of nanoparticles on the film properties and latent heat storage capacity. The latent heat storage capacity of the obtained films was measured via DSC analysis.

Keywords: Phase change material, energy storage, photopolymerization.

References

[1] Mert, M. S., Mert, E. H., Pulko, I., Krajnc, P., and Mert, H. H. (2024). *Form-Stable Oleic Acid Based Polyhipe/Nanoclay Framework Supported Composite Phase Change Materials for Low-Temperature Latent Heat Storage*. Thermal Science and Engineering Progress, 50, 102569.

The Preparation and Mesomorphic Investigation of Binary Mixture of a New Chiral Rod-Like Molecule and Benzoic acid Derivative

Nazlı Özgü KIRAL, Hale OCAK GÜMRÜKÇÜ

Yıldız Technical University, Department of Chemistry, 34220, Istanbul, Türkiye
ozgu.kiral@std.yildiz.edu.tr

Liquid crystals exhibit characteristics between a conventional liquid and a solid crystal. The molecular segments of liquid crystals such as a rigid core, linking groups, terminal groups and lateral substituents play a critical role in determining their phase transition temperatures and thermal stability. Chiral liquid crystals have wide technological applications due to their mesomorphic and electro-optic properties.¹ The role of hydrogen-bonding interactions in the formation and/or stabilization of liquid crystalline phases as well as their stimulus-responsive capability have a significant interest by reserachers.² In this study, firstly the synthesis, structural and mesomorphic characterization of a new benzoate based calamitic molecule with an ester linkage group containing (S)-2-(dodecyloxy)propyloxy group at one of the end positions and a 6-(hydroxyhexyl)oxy chain at the other end were carried out. The new rod-like molecule was characterized by FTIR, ¹H-NMR, ¹³C-NMR, and HR-MS spectroscopic methods and the obtained mesomorphic data from polarized optic microscope (POM) and differential scanning calorimeter (DSC) showed that it is non-mesogenic. The occurrence of crystal behavior is thought to be due to the increased order due to the effect of intermolecular hydrogen bonds between the hydroxyl groups at terminals. In the second step of the study, an binary mixture of the rod-like molecule with mesogenic 4-(hexyloxy)benzoic acid was prepared in order to restruct the molecular order and investigate the effect of intermolecular hydrogen bonding on inducing the liquid crystal phase. The mixture was characterized FTIR and ¹H-NMR, the mesogenic properties were determined by POM and DSC.

Keywords: Liquid crystals, hydrogen bonding, binary mixture.

References

- [1] Ocak H., Bilgin Eran B., Nuray S., Özkonstanyan A., Poppe S., and Tschierske C. (2021). *Extraordinary Magnetic Field Effects on the LC Phases of Homochiral and Racemic 4-Cyanoresorcinol-Based Diamagnetic Bent-Core Mesogens*, J. Mater. Chem. C, 9, 1895-1910.
- [2] Liu B., Yang T., Mu X., Mai Z., Li H., Wang Y., and Zhou G. (2021). *Smart Supramolecular Self-Assembled Nanosystem: Stimulus-Responsive Hydrogen-Bonded Liquid Crystals*, Nanomaterials, 11, 448.

Influence of Monomer Composition on the Morphological and Mechanical Properties of Terpene Derivative Macroporous Polymers

Rumeysa BEYAZ¹, Burcu KEKEVİ², and Emine Hilal MERT³

¹*Institute of Graduate Education, Department of Polymer Materials Engineering,
Yalova University, Yalova, Türkiye*

²*Material and Material Processing Technologies Department, Yalova University,
Yalova Community College, Yalova, Türkiye*

³*Faculty of Engineering, Department of Polymer Materials Engineering,
Yalova University, Yalova, Türkiye*

rmysbeyaz@gmail.com

The environmental and economic consequences of dependence on petroleum-based resources have promoted the use of renewable raw materials in polymer industry. In this context, the use of bio-based monomers has emerged as a key research focus in the advancement of sustainable polymer systems¹. Terpenes, which are among the naturally occurring organic molecules, exhibit chemical reactivity owing to their isoprene-derived structure and abundance of their functional groups. In this regard, d-limonene and β -myrcene, which are belong to terpene family, have potential of participating in polymerization reactions, due to the presence of multiple double bonds¹. Accordingly, terpenes are considered promising monomer candidates for the synthesis of sustainable polymers².

Herein, macroporous thermoset polymers were prepared through the crosslinking of high internal phase emulsion (HIPE) templates consisting of β -myrcene and d-limonene in the continuous emulsion phase. To achieve a cross-linked material 1,6-hexanediol diacrylate (1,6-HDDA) was used as crosslinking comonomer. The morphological and mechanical properties of the obtained polymers were investigated in detail depending on the monomer composition. It was revealed that the ratio of d-limonen and β -myrcene in the continuous emulsion phase plays a critical role to obtain highly open-porous polymers. Moreover, mechanical strength was also found to be varied due to terpene ratio.

Keywords: Terpene, β -myrcene, limonene, macroporous polymer.

References

- [1] Belgacem, M. N., Gandini, A. (Ed.). (2011). *Monomers, Polymers and Composites from Renewable Resources*. Elsevier.
- [2] Mosquera, M. E., et al. (2021). *Terpenes and Terpenoids: Building Blocks to Produce Biopolymers. Sustainable Chemistry*, 2(3), 467–492.

Preparation and Characterization of Alumina Enhanced Kaolinite Supported Capric Acid Based Composite Phase Change Materials

Semanur SAĞLAM¹, Hatice Hande MERT², and Mehmet Selçuk MERT³

¹Yalova University, Institute of Graduate Studies, Department of Chemical Engineering,
77200 Yalova, Türkiye

²Yalova University, Faculty of Engineering, Department of Chemical Engineering,
77200 Yalova, Türkiye

³Yalova University, Faculty of Engineering, Department of Energy Systems Engineering,
77200 Yalova, Türkiye
semanur.saglam@yalova.edu.tr

Phase change materials (PCMs) are chemical substances that are capable of storing thermal energy as latent heat during the melting-crystallization phase change period.¹ PCMs can be used in many areas, which require thermal management depending on their phase transition temperatures and thermal energy storage capacity. In recent years, composite forms of PCMs are preferred instead of their direct use due to the drawbacks of them such as leakage and low heat conduction.²

In this study, a capric acid based shape-stabilized composite PCM was prepared and characterized. Capric acid ($\text{CH}_3(\text{CH}_2)_8\text{COOH}$) is a saturated fatty acid, which is one of the promising organic type of PCMs for low-temperature energy storage applications. The composite PCM was prepared with gamma-alumina enhanced kaolinite support matrix by the solvent assisted vacuum impregnation process. The characterization of composite PCM was performed by differential scanning calorimetry (DSC), Fourier transform infrared spectroscopy (FT-IR) and Scanning Electron Microscopy (SEM) analyses. It was anticipated that the composite PCM has a significant potential for low-temperature energy storage applications with thanks to its melting temperature as 31.0 °C and melting enthalpy as 53.4 J/g.

Keywords: Phase change material, capric acid, thermal energy storage.

References

- [1] Can, G., Mert, H. H., and Mert, M. S. (2022). *Shape-Stabilized N-Heptadecane/Polymeric Foams With Modified Iron Oxide Nanoparticles for Thermal Energy Storage*. *Thermochimica Acta*, 714, 179266.
- [2] Küçüker, B. N., Mert, H. H. (2025). *Preparation of CuO/Al₂O₃ and NiO Loaded Form-Stabilized Composite Phase Change Materials with Improved Thermal Properties and Comparison of their Thermal Energy Storage Characteristics*. *Macromol. Mater. Eng.*, 310, 2400379.

Synthesis of TX-Based Photoinitiator with Antibacterial Activity

Şuhedanur KOCATEPE, Sevnur KESKİN DOĞRUYOL

Yildiz Technical University, Department of Chemistry, 34220, Istanbul, Türkiye

suhedanur.kocatepe@std.yildiz.edu.tr

Photopolymerization is an important industrial process that is frequently used in many fields such as inks, coatings, adhesives, contact lenses, biomaterials, etc. due to its various advantages. The most important component of the photopolymerization process is the photoinitiator, which initiates polymerization by absorbing light. Photoinitiators are classified as type I, type II and one component type II depending on the initiation mechanism. Type II photoinitiator require the use of lower energy light sources than type I photoinitiators as they exhibit absorption properties at longer wavelengths and this behavior reduces the cost of the processes. Thioxanthone (TX) and its derivatives are examples of type II photoinitiators. They are interesting in photochemistry because they have high photoinitiation efficiency and absorb most strongly in the 380-420 nm UV-Visible range. In recent years, a lot of research has been done to synthesize photoinitiators operating in the visible region on the spectrum. This is because cheap and safe for health¹. The use of nitrogen-containing heterocyclic compounds is important for the development of clinically bioactive compounds. The pyrazole ring is one of these heterocyclic compounds it exhibit a wide range of pharmacological effects. The prevention of bacterial buildup on biomedical materials depends on the surface properties of the materials. It is very important develop antibacterial coatings to prevent bacteria from spreading. Combining antibacterial polymers and coatings has been shown to be an effective way to fight bacterial pathogens².

The aim of the study focuses on the synthesis and characterization of thioxanthone-based photoinitiator with antibacterial activity.

Keywords: Photoinitiator, thioxanthone, pyrazole, antibacterial.

References

- [1] J. Bessa L., Palmeira A., Gomes S.A., Vasconcelos V., Sousa E., Pinto M., and Martins da Costa P. (2015). *Synergistic Effects Between Thioxanthenes and Oxacillin Against Methicillin-Resistant Staphylococcus Aureus*, Macro. Drug Resis., 21(4).
- [2] Altikatoglu Yapaoz, M., Dogruyol, S. (2024). *Pyrazole-based Photoinitiator with Antibacterial Activity*. Polimery, 69(3), 167-172.

Assessment of the Impact of Climate Change Projects on the Sustainable Development of the Country Based on Multi-Criterion Analysis Method

Sanam VALİYEVA, Muslum GURBANOV

*Institute of Radiation Problems of the Ministry of Science and Education of the Republic of Azerbaijan,
AZ1143, Baku, Azerbaijan
senem.shukurli@mail.ru*

The Republic of Azerbaijan joined international efforts to mitigate the negative impacts of global climate change by ratifying the UN Framework Convention on Climate Change in 1995, the Kyoto Protocol to the Convention in 2000, and the Paris Agreement in 2016. Azerbaijan, with a population of approximately 10 million, accounts for only 0.15% of global greenhouse gas emissions¹.

Azerbaijan is determined to become a leading country in the field of renewable energy and is using its rich wind and solar potential to this end. It is also a crucial component of the country's plan to reduce greenhouse gas emissions by 40% by 2050². It is necessary to develop projects to reduce greenhouse gas emissions in Azerbaijan in the following areas: energy, industry, agriculture, land use and forestry, and waste management are priority sectors. The assessment of the project's impact on Sustainable Development is carried out according to the following 4 criteria: social criteria, economic criteria, environmental criteria and influencing policy in relevant areas^{3,4}. The assessment is made using numbers between -1 and +1, taking into account positive and negative effects. Calculations should be made based on the corresponding impact values, with score values ranging between -1 and +1 for each project under consideration. The calculations should be continued in accordance with the National Criteria system and concluded with a result indicating the impact of the Project under consideration on the Sustainable Development of the Country. If the result obtained is positive, the Project under consideration has a positive impact on the Sustainable Development of the Country and can be implemented as a Clean Development Mechanism project.

Keywords: Greenhouse gas emissions, renewable energy, industry, sustainable development.

References

- [1] Fourth National Communication to the United Nations Framework Convention on Climate Change Republic of Azerbaijan, <https://unfccc.int/sites/default/files/resource/FNC%20report.pdf>.
- [2] <https://eco.gov.az/az/nazirlik/xeber?newsID=20466>.
- [3] How-to Guide: National Institutional Frameworks for the Kyoto Protocol Flexible Mechanisms in Eastern Europe and the Commonwealth of Independent States, UNDP, 2006.
- [4] Albert Zweering, Proposed Method for Sustainability Assessment of CDM Projects, 2005.

Improving Flame Resistance and Polymer Compatibility of Ethylene Propylene-Diene Rubber (SKEPT-60) Through Polyvinyl (PVX) Modification

Fariz Ali AMİRLİ¹, İbrahim Gumbat MOVLAYEV¹, Rana Faig KHANKSHİYEVA²,
and Aynur Fazil MAMMADOVA¹

¹ Azerbaijan State Oil and Industry University, Baku, Azerbaijan

² Institute of Radiation Problems, Baku, Azerbaijan

aynur.memmedova@asoiu.edu.az

This study aims to enhance the flame resistance and compatibility of ethylene-propylene-diene rubber (SKEPT-60) with other polymers by modifying it with polyvinyl (PVX) at different ratios. SKEPT-60, an unsaturated copolymer with high ozone and temperature resistance (Mooney viscosity: 60 Pa·s, dicyclopentadiene content: 1.8%), exhibits poor flammability resistance and limited polymer compatibility. To address these drawbacks, SKEPT-60 was mechanically blended with PVX at 40–50°C for 3–4 minutes, forming binary (SKEPT-60/PVX) systems.

The rheological properties of these blends were investigated under varying loads (11.75, 20.85, 26.10, 32.60 kg) and temperatures^{1,2}. Results indicated that the optimal PVX content for improved performance was 2–4 phr (parts per hundred rubber)³. Further validation involved preparing filled composite materials based on SKEPT-60/PVX blends, which were vulcanized at 155°C for 25 minutes. The modified composites exhibited significant enhancements in flame resistance, polymer compatibility, tensile strength, metal adhesion, tear resistance, fatigue durability, and aging coefficients compared to unmodified SKEPT-60.

Keywords: Ethylene-propylene-diene rubber (EPDM), polyvinyl modification, flame resistance, polymer compatibility, rheological properties.

References

- [1] Movlayev I. G., Mammadova A. F. (2024). *Determination of the Main Parameters of the Modified Epoxide Oligomer*. International Science Group. International Science Journal of Engineering & Agriculture, 3(6), 95-100.
- [2] Amirli F. A., Movlayev I. H., and Mammadova A. F. (2025). *Study of the Rheology Properties of the Mixture of Terminal Ethylene-Propylene Rubber with Benzydamine-Modified Phenol-Formaldehyde Oligomer PPOR*. 24(4), 229-239.
- [3] Amirli, F.A., Movlayev, I. H., Aliyeva, G. A., and Mammadova, A. F. (2023). *Compositions Based on Modified and Filled Epoxy Oligomer*. PPOR, 24(4), 689-696.

Synthesis of Reagents Based On Natural Oil Acids and Ethylenediamine and Their Study As Components For Preservation Fluids

Vaqif ABBASOV, Ulviyye ABBASOVA, Sara ABBASZADE, and Durna AGHAMALIEVA
*Institute of Petrochemical Processes named after acad. Y.M. Mammadaliyev, Ministry of Science and
Education of the Republic of Azerbaijan, Baku, Azerbaijan*
abbasova.u.1986@gmail.com

Atmospheric corrosion remains a critical issue in countries with advanced oil and gas industries, particularly exacerbated by growing anthropogenic pollution¹. One of the effective preventive strategies is the use of conservation fluids with corrosion-inhibitive components. In this study, reagents based on natural petroleum acids (NPAs) and ethylenediamine were synthesized, including amidoamine derivatives and their complexes with transition metal salts (Mn, Fe, Zn, Ni). These reagents were incorporated into T-46 oil distillate both individually and as composite mixtures to formulate protective conservation fluids².

The synthesized conservation fluids were evaluated under accelerated corrosion conditions in humidity chambers, marine water, and 0.001% sulfuric acid solution. The corrosion resistance period of steel-3 samples ranged up to 308 days, depending on the composition and concentration of the additive³. Notably, the complex of amidoamine with the iron salt of NPAs demonstrated the longest protective effect (up to 270 days in humidity chambers), while also offering superior performance in marine and acidic environments. Given the economic advantage and availability of iron salts compared to other metal counterparts, this composition is considered a promising candidate for practical applications in atmospheric corrosion protection.

Keywords: Corrosion inhibitors, amidoamine, ethylenediamine, natural petroleum acids, conservation fluids.

References

- [1] Afandiyeva, L., Abbasov, V. M., Aliyeva, L. I., Ahmadbayova S., Azizbeyli, E., and Ahmed, H. M. E. (2018). *Investigation of Organic Complexes of Imidazolines Based on Synthetic Oxy- and Petroleum Acids as Corrosion Inhibitors*. Iranian Journal of Chemistry & Chemical Engineering, 37(2), 137-144.
- [2] Kosmyrna, M., Nosachova, Y., and Shabliy, T. (2024). *Evaluation of the Efficiency of Inhibitor Protection of Metal Surfaces in Aggressive Water and Oil Environments*. Visnyk NTUU KPI, Chemical Engineering, Ecology and Resource Saving, 4, 26 December.
- [3] Research of Conservation Liquids Made on The Basis of Residual Fatty Acids, Nitro Compounds and Additives Based on Derivatives of Natural Petroleum Acids. (2023). Azerbaijan Journal of Chemical News, 2, 98-105.

Chitosan-Tannin Based Flocculant for Heavy Metal (Ni^{2+} , Cu^{2+} , and Zn^{2+}) Removal from Industrial Wastewater: Investigating the Influence of pH, Reaction Time, and Stirring Speed

Sara HASANOVA¹, Rana KHANKİSHİYEVA^{1,2}

¹Azerbaijan University of Architecture and Construction, Baku, Azerbaijan

²Institute of Radiation Problems, Baku, Azerbaijan
hasanova.sara@azmiu.edu.az

The removal of heavy metal ions from wastewater is a great importance for environmental cleanliness and human health. Wastewater discharged from various sources, such as industry, agriculture, and municipalities, contains a large amount of organic and inorganic pollutants, which causes serious consequences for the ecosystem as a whole¹. There is a growing need for a treatment method that is not only effective but also eco-friendly and cost-efficient. While natural coagulants have been widely researched for water treatment, the potential of chitosan modified with tannins as a natural coagulant is still not fully explored. This study investigates the removal of Cu^{2+} , Zn^{2+} , and Ni^{2+} from industrial wastewater using a chitosan-based composite flocculant derived from tannins extracted from pomegranate peel. The study looks into various factors that influence the removal process, such as solution pH, adsorbent dosage, initial metal concentration, temperature, and contact time, and discusses how each of these factors affects the performance of the flocculant. Tannins were extracted by drying and grinding pomegranate peels into an 80-mesh powder, followed by Soxhlet extraction using 80% methanol². Their presence was confirmed through a colorimetric ferric chloride test. Chitosan and tannins were combined under controlled conditions, followed by crosslinking with glutaraldehyde. The resulting composite was filtered, washed, and dried. The material demonstrated potential for heavy metal removal, offering a sustainable approach to wastewater treatment. Synthesis of bioflocculant significantly improved key water parameters, including pH and turbidity. The removal efficiencies achieved were up to 90% for Cu^{2+} , 75% for Zn^{2+} , and 70% for Ni^{2+} . The addition of the flocculant, along with proper pH adjustment, has improved the efficiency of the metal removal processes. pH was identified as a critical variable, with specific optimum values determined for different metals. Compared to traditional methods such as chemical precipitation and conventional coagulation-flocculation processes, the pomegranate peel-chitosan flocculant offers advantages due to its natural origin, ease of production, and simplified pH adjustment requirements.

Keywords: Chitosan, tannin, bioflocculant, heavy metal removal, wastewater treatment.

References

- [1] Basem, A., Jasim, D. J., Majdi, H. S., Mohammed, R. M., Ahmed, M., Al-Rubaye, A. H., Kianfar E. (2024). *Adsorption of Heavy Metals from Wastewater by Chitosan: A Review. Results in Engineering*, 23, 102404.
- [2] Saad, H., Bouhtoury, F.C.-E., Pizzi, A., Rode, K., Charrier, B., and Ayed, N. (2012). *Characterization of Pomegranate Peel Tannin Extractives. Industrial Crops and Products*, 40, 239–246.

Intelligent Identification of Multicomponent Mixture Components Using an Artificial Intelligence

Arzu İBRAHİMOVA

Azerbaijan State Oil and Industry University, Baku, Azadliq ave., 20
arzu.ibrahimova@sabah.edu.az

This article presents an advanced method for the qualitative and quantitative identification of components in multicomponent mixtures using two densitometers and artificial intelligence (AI). The proposed approach eliminates dependence on chromatographic operating conditions by incorporating a standardized dual-detector system and a mathematical model for determining molecular mass. The core equations are enhanced through regression analysis and neural network techniques, enabling automated and high-precision identification without manual calibration.

The article introduces ChromAI ID Pro, a dedicated software platform that integrates signal acquisition, real-time analysis, AI-based predictions, visualizations, and report generation. The system utilizes machine learning algorithms such as Random Forest and MLPRegressor to predict molecular weights based on detector signals and known standard parameters. Comparative simulations demonstrate a significant reduction in prediction error and improved reliability under variable conditions.

Experimental data and illustrative examples, including model accuracy comparisons and graphical outputs, are provided to validate the efficiency of the method. The proposed solution is applicable to industrial and laboratory settings where fast, accurate component identification is essential.

Keywords: Gas chromatography, densitometer, regression analysis, neural network, artificial intelligence, molecular mass prediction, ChromAI ID Pro.

References

- [1] Farzane, N. G., Ibragimova, A. E. (2010). Development of an Intellectual Chromatograph. Baku: Elm Publishing.
- [2] Farzane, N. G., Ibragimova, A. E. (2013). *Two-Parametric Identification of Complex Mixtures on a Standard Column*. The Azerbaijan Oil Industry Journal, 27, Baku: SOCAR.
- [3] Bishop, C. M. (2006). Pattern Recognition and Machine Learning. New York: Springer.
- [4] Hastie, T., Tibshirani, R., and Friedman, J. (2009). The Elements of Statistical Learning: Data Mining, Inference, and Prediction (2nd ed.). New York: Springer.

Investigation of Concentration Liquids Formulated by Incorporating Amidoamine and Various Fatty Acids into T-30 Oil Distillate

*Elgun HASANOV, Sara ABBASZADE, Rashad AGAKİSHİYEV,
Rufana ALİZADE, and Vugar GASİMOV*

*Institute of Petrochemical Processes named after acad. Y.M. Mammadaliyev, Ministry of Science and
Education of the Republic of Azerbaijan, Baku, Azerbaijan
rufana.alizada93@gmail.com*

Atmospheric corrosion refers to the gradual degradation of metals and other materials due to the interaction with atmospheric elements through chemical or electrochemical mechanisms. It is one of the most prevalent forms of corrosion, primarily influenced by environmental factors such as humidity, oxygen, carbon dioxide, and airborne salt particles¹⁻³. To mitigate this issue, the application of preservation fluids and lubricants offers a more cost-effective and practical solution. In the current study, corrosion tests were conducted in accordance with GOST 9054-75 standards within a "Q-4" thermo-humidity chamber, utilizing seawater and a 0.001% amidoamine-based preservation fluid derived from vegetable fatty acids. The results aim to evaluate the effectiveness of these preservation fluids in preventing atmospheric corrosion under controlled conditions.

Amidoamine compositions synthesized from cottonseed and sunflower fatty acids in a 1:1 molar ratio were prepared, followed by the formulation of preservation fluids through the addition of T-30 oil distillate. Based on the conducted tests, it can be concluded that the corrosion protection efficacy of the preservation fluids, formulated with amidoamine and organic acids, is superior to that of the T-30 oil distillate used as a solvent, across all three tested environments on metal substrates.

The corrosion protection effectiveness of preservation fluids prepared by adding 10% and 20% of these inhibitors to T-30 oil distillate on metal plates was observed to be 310 and 326 days, respectively, in the "G-4" thermo-humidity chamber, 142 and 145 days in seawater, and 139 and 142 days in a 0.001% H₂SO₄ solution.

Keywords: Amidoamine, fatty acid, hydrochamber, corrosion protection.

References

- [1] Abdullaeva, N. R., Abbasov, V. M., and Ismailov, T. A. (2009). *Antistatic Properties of Oxyethers of Organic Acids Obtained from Vegetable Oils in Composition with Diesel Fuel*. Processes of Petrochemistry and Oil Refining, 10, 2(38), 112-115.
- [2] Bessonov, J. I. A. (2006). Theoretical Foundations of Electrical Engineering: Electric Circuits. M.: Gardariki, 701.
- [3] Irodov, I.E. (2002). Electromagnetism: Basic Laws. M.: Lab. Basic Knowledge, 352.

Nitration Behavior of Aromatic- and Olefin-Enriched Coking Light Fraction with Ionic Catalytic Systems

Vaqif ABBASOV, Sara ABBASZADE, Teyyub İSMAİLOV, Feride TAHİROVA,
Tofiq ALİYEV, and Khayala ABBASOVA

*Institute of Petrochemical Processes named after acad. Y.M. Mammadaliyev, Ministry of Science and
Education of the Republic of Azerbaijan, Baku, Azerbaijan*

sara.abbaszade578@gmail.com

The direct nitration of the light fraction of coking distillates — composed of aromatic, olefinic, and aliphatic hydrocarbons — presents significant technological challenges due to the similar boiling points of its constituents and the high, non-selective reactivity of olefins. In this study, a coking light fraction, previously characterized via sulfolane extraction to contain approximately 32% aromatics and olefins, was subjected to nitration without prior fractionation or separation. The reactions were carried out using 59% nitric acid in the presence of catalytic systems comprising sodium nitrite (NaNO_2), N-methylpyrrolidone (NMP), and imidazolium-based ionic liquids^{1,2}. By varying the reaction temperature and catalyst combinations, product yields ranging from 54% to 66% were achieved. IR and NMR analyses confirmed that nitration occurred predominantly on aromatic structures, with minimal formation of O- NO_2 groups and oxidative by-products. The enhanced selectivity and non-volatile nature of the ionic liquids simplified product separation, contributing to a more efficient and energy-saving process. These findings demonstrate that ionic liquid-based nitration offers an effective and scalable strategy for the direct functionalization of petroleum fractions rich in aromatics and olefins, eliminating the need for prior separation steps³.

Keywords: Coking fraction, aromatic nitration, ionic liquids, selective catalysis, green nitration.

References

- [1] Laali, K. K., Gettwert, V. J. (2001). *Electrophilic Nitration of Aromatics in Ionic Liquid Solvents*. The Journal of Organic Chemistry, 66(1), 35-40.
- [2] Michele, C., Caputo, D., Fusco, C., Cotugno, P., Rizzi, V., Dellanna, M. M., D'Accolti, L., and Nacci, A. (2020). *Ionic – Liquid Controlled Nitration of Double Bond: Highly Selective Synthesis of Nitrostyrenes and Benzonitriles*. European Journal of Organic Chemistry.
- [3] Liu, S. (2003). *Nitration of Simple Aromatics and Epoxidation of Alkenes in Ionic Liquids*. Swansea University E – Theses.

Phosphoryl-Substituted α -Chloro- and α -Thiocyanatoacetaldehydes in the Reaction with α -Pyridone

Sevda ASADOVA, Khammad ASADOV, and Mahsati GAFAROVA

*Institute of Petrochemical Processes named after acad. Y.M. Mammadaliyev, Ministry of
Science and Education of the Republic of Azerbaijan, Baku, Azerbaijan*

sevdabedelova34@gmail.com

In the reactions of phosphorylated α -chloro- and α -thiocyanatoacetaldehydes with α -pyridone, two classes of heterocyclic compounds, 2-phosphoryl-oxazolo[2,3-a]pyridinium chlorides and 4-pyridyl-substituted 2-oxothiazolidines, were synthesized. These compounds feature a rare combination of phosphoryl, oxazolyl, and thiazolidine groups, making them promising for potential biological activity and coordination chemistry applications^{1,2}. The product structures were confirmed by NMR spectroscopy, which revealed characteristic coupling constants and chemical shifts. The study highlights α -pyridone as a versatile nucleophile in constructing C-phosphorylated heterocycles with potential functional applications³.

Keywords: Phosphoryl aldehydes, α -Pyridone, oxazolo compounds, heterocycles.

References

- [1] Hurtado-Rodríguez, D., Salinas-Torres, A., Rojas, H., Becerra D., and Castillo J.C. (2022). Bioactive 2-pyridone-containing Heterocycle Syntheses Using Multicomponent Reactions. *RSC Advances*, 54.
- [2] Volkova, Y., Zavarzin, I. (2023). Synthesis of Phosphorus(V)-Substituted Six-Membered N-Heterocycles: Recent Progress and Challenges. *Molecules*, 28(6).
- [3] Larina, L. (2023). C- and N-Phosphorylated Enamines-An Avenue to Heterocycles: NMR Spectroscopy. *International Journal of Molecular Sciences*, 24(11).

TURAN-FUNDAMENTAL SCIENCES SYMPOSIUM
TURAN-TEMEL BİLİMLER SEMPOZYUMU

Honorary Chair/Onursal Başkan: Prof. Dr. Aziz SANCAR

BIOLOGY

Analysis of Codon Usage Bias of Chloroplast Genomes in *Camellia* Species

Yeşim AKTÜRK DİZMAN

Recep Tayyip Erdoğan University, Faculty of Arts and Sciences, Department of Biology, Rize, Türkiye

yesim.akturk@erdogan.edu.tr

Camellia species are significant ornamental shrubs and flowering trees, valued for their economic importance in various sectors, including medicine, food, and beverages.¹ The chloroplast genome is commonly utilized in chloroplast gene expression, studies of species evolution, and genetic transformation within the chloroplast.² Codon usage bias (CUB), which describes the non-random preference for certain synonymous codons over others, provides valuable insights into gene function, species evolution, and foreign gene expression.³ Although CUB in chloroplast genomes of *Camellia* species has not been extensively studied, this research systematically analyzed the chloroplast genomes of three *Camellia* species (*C. sinensis*, *C. nitidissima*, and *C. ptilophylla*) to investigate the factors influencing CUB patterns. Our findings demonstrated that the chloroplast genomes of the three *Camellia* species exhibit a strong bias toward AT-rich sequences and a preference for codons ending in A or T. Across these species, 17 high-frequency codons were consistently identified, along with 10 to 19 optimal codons. The ENC-plot, PR2-plot, neutrality plot, and correspondence analysis collectively indicated that CUB in the three *Camellia* species is shaped by both natural selection and mutation pressure, with natural selection playing a more prominent role. Comparison of codon usage frequencies with those of four common model organisms revealed that *Nicotiana tabacum* and *Saccharomyces cerevisiae* are suitable candidates for heterologous gene expression. Collectively, this study elucidated the patterns of synonymous codon usage in *Camellia* species, offering important insights that could facilitate future genetic engineering efforts in these plants.

Keywords: *Camellia* species, codon usage bias, chloroplast genome, natural selection.

References

- [1] Teixeira, A.M., Sousa, C. (2021). A Review on the Biological Activity of *Camellia* Species. *Molecules*, 26(8), 2178.
- [2] Wang, Y., Jiang, D., Guo, K., Zhao, L., Meng, F., Xiao, J., Niu, Y., et al. (2023) Comparative Analysis of Codon Usage Patterns in Chloroplast Genomes of Ten *Epimedium* Species. *BMC Genom Data*, 24(1), 3.
- [3] Yang, X., Wang, Y., Gong, W., and Li, Y. (2024). Comparative Analysis of the Codon Usage Pattern in the Chloroplast Genomes of *Gnetales* Species. *Int J Mol Sci.*, 25(19), 10622.

TURAN-Fundamental Sciences Symposium (TURAN25) Delivered Oral Presentations

June 23, 2025 – MONDAY

08.30-10.00	REGISTRATION	
10.00-12.00	OPENING SPEECHES <i>Prof. Dr. Salim YÜCE</i> , Chair of the Symposium, Dean of the Faculty of Arts and Sciences, Yıldız Technical University <i>Prof. Dr. Misir MARDANOV</i> , Azərbaycan Cumhuriyeti Bilim və Təhsil Nazirliyi Matematik və Mekanik İnstitutu Müdürü <i>Prof. Dr. Eyüp DEBİK</i> , Rector, Yıldız Technical University <i>Prof. Dr. Aziz SANCAR</i> , Honorary Chair of the Symposium	
12.00-13.30	LUNCH	
13.30-14.15	INVITED PLENARY TALK 1: <i>Prof. Dr. İbrahim Halil KAVAKLI</i> , Can We Target the Core Clock Mechanism for the Treatment of Different Diseases?, Chair: <i>M. Kasım ŞENER</i>	
14.15-14.45	INVITED TALK: <i>Kemal Yavuz ATAMAN</i> , Maveraünnehir, Horasan, Harezmi Havzasında Muhtelif İlimlerin Birlikte Gelişmesi, Meşhur Alimlerin Yetiştirilmesi, Kıymetli Eserlerin Yazılması Üzerine Bir Analiz, Chair: <i>Senem ÖNER BULUT</i>	
14.45-15.00	Coffee and Tea Break	
	HALL 1, Session 2, Mathematics, Chair: <i>Fatma KARACA</i>	HALL 2, Session 2, Mathematics, Chair: <i>Aysel TURGUT VANLI</i>
15.00-15.15	<i>Zehra İŞBİLİR</i> , 2-Parameter Generalized Quaternions with Generalized Tribonacci Numbers Components	<i>Suzan IŞIK</i> , Improved Fixed Point Approximations via Picard's Three-Step Iteration for Suzuki-Type Non-Expansive Mappings
15.15-15.30	<i>Edanur ERGÜL ARSLAN</i> , Calculus on Real-variable Split Quaternion-valued Functions	<i>İpek Ebru KARAÇAY</i> , Investigation of Fractal Dimension Algorithms in Different Geometries
15.30-15.45	<i>Pelin DURSUN</i> , Characteristic Roots, Eigenvalues and Eigenvectors of a Certain Type of 4x4 Dual Tridiagonal Matrices	<i>Radva ANSARY</i> , Machine Learning-Based Classification of Chaotic Behavior In Discrete Dynamical Systems
	HALL 3, Session 2, Mathematics, Chair: <i>Seda KIZILBUDAK ÇALIŞKAN</i>	HALL 4, Session 2, Physics, Chair: <i>Banu Süngü MISIRLIOĞLU</i>
15.00-15.15	<i>Deniz GÜNDOĞAN</i> , Modelling Electromagnetic Wave	<i>İsmet İŞGÖR</i> , Electrodeposition and Spray Techniques for Pt/Pd Coated Nickel Foam Electrodes
15.15-15.30	<i>Handan ÇELİK</i> , Numerical Modeling of Stars and Mathematical Analysis	<i>Fatih PİLAVCI</i> , Körük Ventili (Levelling Valve) Hava Akış Karakteristiklerinin Karşılaştırmalı Teknik İncelemesi
15.30-15.45	<i>Abdüllatif YALÇIN</i> , New Midpoint and Trapezoid-Type Inequalities for Co-ordinated m-Convex Functions via Variable-Order Fractional Calculus	<i>Serdar DİZMAN</i> , Sinop İlinde Nükleer Güç Santrali Öncesi Kabalı Nehrinin Sularında Tritiyum Radyoaktivitesinin Araştırılması
14.45-15.00	Coffee and Tea Break	

TURAN-FUNDAMENTAL SCIENCES SYMPOSIUM
TURAN-TEMEL BİLİMLER SEMPOZYUMU

Honorary Chair/Onursal Başkan: Prof. Dr. Aziz SANCAR

	HALL 1, Session 3, Mathematics, Chair: <i>Fatma KARACA</i>	HALL 2, Session 3, Mathematics, Chair: <i>Aysel TURGUT VANLI</i>
16.00-16.15	<i>Sema Nur ÇEPE</i> , Comparative Analysis of Türkiye's Road Traffic Accidents Using Data Mining Techniques	<i>Dilara KARSLIOĞLU</i> , Variation of parameters and Hölder stability of differential
16.15-16.30	<i>Kübra ÇETİNERBERK</i> , The Moment of Inertia of Generated Curves: A Holditch-Type Theorem	<i>Gamze ALKAYA</i> , W5-Curvature Tensor on a Complex Contact Space Form
16.30-16.45	<i>Merve ÇOLAK</i> , Do Oil Prices Have a Chaotic Effect on Unemployment and Inflation	<i>Yonca SEZER</i> , Schauder Estimates for Elliptic Operators in Banach-Sobolev Spaces
16.45-17.00	<i>Nisa ÇELİK</i> , Genel DSW Sistemine Genişletilmiş Jacobi Eliptik Fonksiyon Yaklaşımı: Tam Çözümler, Kararlılık ve Simülasyonlar	<i>Ömer DURAN</i> , Identities on the generalized harmonic numbers of rank with order using generating functions
	HALL 3, Session 3, Mathematics, Chair: <i>Neslihan ÖZDEMİR</i>	HALL 4, Session 3, Mathematics, Chair: <i>Mustafa DÜLDÜL</i>
16.00-16.15	<i>Çiğdem Zeynep YILMAZ</i> , A Study on Dual Quaternions with k-Pell and k-Pell-Lucas Coefficients	<i>Hamide Feyza AYKUT</i> , Hyperbolic Functions on Split Quaternions and Their Characteristics
16.15-16.30	<i>Duygu ÇAĞLAR ÇAY</i> , A Study on Vietoris' Hybrinomials	<i>Sevilay ÇORUH ŞENOCAK</i> , Surface Pencil With Constant Gaussian Curvature Along A Non-null Curve
16.30-16.45	<i>Emel KARACA</i> , On multiplicative magnetic flux ruled surfaces	<i>Damla BUDAK</i> , Rectifying Rose Ruled Surfaces
16.45-17.00	<i>Gülsüm Yeliz SAÇLI</i> , A Study on the Dual Quaternionic Sequence with Vietoris' Components	<i>Zehra Nur BAKIRBAŞ</i> , The Cross products on Dual and Dual Lorentzian Spaces
17.00-18.00	POSTER	
18.30	DINNER	
June 24, 2025 – TUESDAY		
10.00-10.45	INVITED PLENARY TALK 2: <i>Prof. Dr. Bilal BİLALOV</i> , Some Questions of X-valued Elliptic Equations, Chair: Misir MERDANOV	
10.45-11.15	INVITED TALK: <i>Dr. Amir MOSAVÍ</i> , A Decade of Machine Learning Research: Evaluation Metrics, Taxonomies and Bibliometrics Analysis, Chair: Yonca SEZER	
11.15-11.30	Coffee and Tea Break	
	HALL 1, Session 1, Mathematics, Chair: <i>Neslihan ÖZDEMİR</i>	HALL 2, Session 1, Mathematics, Chair: <i>Fatma KARACA</i>
11.30-11.45	<i>Betül ÖZTÜRK</i> , Bazı Antiviral Moleküllerin Topolojik Karakterizasyonu	<i>Günay ASLAN</i> , Solitary Wave Solutions of the Two-Dimensional Fractional-Order Nonlinear Schrödinger Equation
11.45-12.00	<i>Emine KOPARAL</i> , Affine Factorable Surfaces in Euclidean 4-Space E4	<i>Elman İBRAGIMOV</i> , Boundedness criteria for the G-fractional integral and G-fractional maximal operator on Gegenbauner-morrey spaces
12.00-12.15	<i>Beyza SARUHAN</i> , Monge Surfaces With Their Applications	<i>Bedriye Ecem AKGÜÇ CİMBİZ</i> , Hemen Hemen Neo Kobalans Sayıları

TURAN-FUNDAMENTAL SCIENCES SYMPOSIUM
TURAN-TEMEL BİLİMLER SEMPOZYUMU

Honorary Chair/Onursal Başkan: Prof. Dr. Aziz SANCAR

	HALL 3, Session 1, Chemistry, Chair: <i>Göknur YAŞA ATMACA</i>	HALL 4, Session 1, Biology, Chair: <i>Burgu Gündüz ERGÜN</i>
11.30-11.45	<i>Nargiz RAHIMLI</i> , Particle Size Analysis of Polymer-Based Metal Nanoparticles Using DLS Method	<i>Deniz Duru DERVİŞ</i> , Potential Use of Watermelon Seed Oil as an Alternative Linoleic Acid Source Enhanced by Coffee-Derived Caffeic Acid in Cholesterol Treatment
11.45-12.00	<i>Elmira ALIYEVA</i> , The Study of The Swelling Ratio of Gum Arabic and Acrylamide Hydrogel in Solutions with Different Ph	<i>Meliha Feryal SARIKAYA</i> , Bazı Yonca (<i>Medicago sativa</i> L.) Genotiplerinin iPBS-retrotranspozon Markörleri ile Genetik Çeşitliliğin Değerlendirmesi
12.00-12.15	<i>Akbota KUANDYKOVA</i> , Synthesis, structure, and properties of new alicyclic and aromatic amidophosphates	<i>Narmin GULIYEVA</i> , γ (Gama) Işınlarıyla Muamelenin Bitkilerde Tuz (NaCl) Stresine Tepkisi
12.30-13.30	LUNCH	
13.30-14.15	INVITED PLENARY TALK 3: Prof. Dr. Salim ÇERÇİ, CERN Today: Pushing the Boundaries of Particle Physics, Chair: <i>Zeynep GÜVEN ÖZDEMİR</i>	
14.15-14.45	INVITED TALK: Prof. Dr. Farman MAMEDOV, On the L1data Dirichlet Problem for the Nonuniform Parabolic Equations of Second Order, Chair: <i>Misir MERDANOV</i>	
14.45-15.00	Coffee and Tea Break	
	HALL 1, Session 2, Mathematics, Chair: Serkan ÇAKMAK	HALL 2, Session 2, Mathematics, Chair: Yaprak DERİCİOĞLU
15.00-15.15	<i>Abdurrahman DAYIOĞLU</i> , Bazı Özel Afın Yapıların ve Duallerinin Kromatik İndislerinin Bulunması Üzerine	
15.15-15.30	<i>Neslihan ÖZDEMİR</i> , Optical bullets for the (3+1)-Dimensional Perturbed Nonlinear Schrödinger Equation having the Cubic-Quintic Laws and the Spatio-Temporal Dispersion	<i>Muhammet KNEFATI</i> , Δ -Convergence For Proximal Point Algorithm And Fixed Point Problem in Generalized CAT(0) Spaces
15.30-15.45	<i>Basri ÇELİK</i> , TBMM Maarif Vekâletinin 1920 yılında aldığı geometri öğretimi kararları üzerine	
	HALL 3, Session 2, Chemistry, Chair: <i>Özlem YAZICI</i>	HALL 4, Session 2, Physics, Chair: <i>Macide CANTÜRK</i>
15.00-15.15	<i>Rana KHANKISHIYEVA</i> , Development of Lead-Free Epoxy Resin Composites for X-Ray Shielding	<i>Ainur SERIKBAIKYZY</i> , Surface Engineering of Ti13Nb13Zr Implants Using Microarc Oxidation and Hydroxyapatite-Based Thermal Spraying
15.15-15.30	<i>Tuğrulhan DEMİR</i> , Alunit Cevherinin RSM Kullanılarak Liçinin Optimizasyonu	<i>Zoltan FODOR</i> , A New Force in Nature? The Magnetic Moment of the Muon
15.30-15.45	<i>Günel AZİMOVA</i> , Synthesis of Green Chemistry-Based Catalyst for Carbon Monoxide Neutralization	<i>Rakhshana MAMISHOVA</i> , Dielectric Properties Of CuTiS ₂ Crystals Studied Via Impedance Spectroscopy
15.45-16.00	Coffee and Tea Break	

TURAN-FUNDAMENTAL SCIENCES SYMPOSIUM
TURAN-TEMEL BİLİMLER SEMPOZYUMU

Honorary Chair/Onursal Başkan: Prof. Dr. Aziz SANCAR

	HALL 1, Session 3, Mathematics, Chair: <i>Basri ÇELİK</i>	HALL 2, Session 3, Mathematics, Chair: <i>Neslihan ÖZDEMİR</i>
16.00-16.15	<i>Hande USLU TUNA</i> , Competitive Systems via Hamiltonian Framework and Artificial Neural Estimation	<i>Nail TUKTAMYSHOV</i> , Impulse-Hydrodynamic Model of Explosive Impact on a Half-Plane with Circular Cavity
16.15-16.30	<i>Yaprak DERİCİOĞLU</i> , Data-Driven Estimation of Krylov Iterations in Riccati Solvers	<i>Abdusobir SAIDOV</i> , Model And Theorem Of Existence Of The Problem Of Optimal Control Of The Customs Clearance Process
16.30-16.45	<i>Serkan ÇAKMAK</i> , q-Türev Kullanılarak Tanımlanan Yeni Bir Harmonik Fonksiyon Sınıfının Geometrik Özellikleri	<i>Narmina GUBATOVA</i> , Necessary conditions for weak and strong local minimum the problem of the calculus of variations with higher order derivatives
16.45-17.00	<i>Fatma KARACA</i> , Some Characterizations of Gradient Ricci-Yamabe Solitons	<i>Djamal HAMADI</i> , Numerical Analysis of Stiffened Shell Structures Using an Efficient Three-Dimensional Finite Element
17.00-17.15	<i>Burcu ALTINTAŞ</i> , Matematik Lisans Öğrencilerinin Seçmeli Ders Seçimlerinin Oyun Teorisi İle Matematiksel Modellenmesi	<i>Shahin GULIYEV</i> , Development of a General Algorithm for Solving Stability Problems of Plates on Elastic Bases
	HALL 3, Session 3, Chair: <i>Misir MERDANOV</i>	HALL 4, Session 3, Physics, Chair: <i>Altan BOZDOĞAN</i>
	<i>Adil ABDULLAYEV</i> , Dielectric Relaxation and Conductivity Characteristics of Gamma-Irradiated TlGaTe2 and TlInSe2 Crystals	<i>Aydan KKHALIGZADE</i> , Effect of Doping and Radiation on the Conductivity Mechanism in Layered GaS Crystals
	<i>Aida TAGIYEVA</i> , On an Algorithm for a Morphological Analyzer of the Azerbaijani Language	<i>Amina MIKAYILOVA</i> , Prospects of Thermal Waters Usage in Small Power Industry
	<i>Surkay AKBAROV</i> , The Influence of the Interaction with Fluids of the Hollow Cylinder with Inhomogeneous Initial Stresses on the ZGV Modes that Occur with Axisymmetric Wave Propagation	<i>Sümeyye BÜYÜKKAZAZ</i> , (E)-2-(2,4-dihidroksibenziliden) tiyosemikarbazon ve (E)-2-[[1H-indol-3-il) metilen] tiyosemikarbazon komplekslerinin yapısal ve spektroskopik özelliklerinin DFT metoduyla incelenmesi
		<i>Zarina ARINGOZHINA</i> , Optimization of Ion-Plasma Nitriding Parameters for Ti-6Al-4V Titanium Alloy
	<i>Baharchin AKHMADLI</i> , On the strong solvability of a nonlocal boundary value problem for the Laplace equation in weighted grand Sobolev spaces in rectangle	<i>İpek BALNAN</i> , Enhancement of Visible-Light-Driven Photocatalytic Activity in Gd-Doped CdZnS Nanocomposites
17.15-18.00	POSTER	
18.30	DINNER	
June 25, 2025 – WEDNESDAY		
10.00-11.30	BILATERAL COOPERATION PANEL	
11.30-12.30	LUNCH	
12.30	EXCURSION (17.00 BOAT)	
20.00	DINNER	

TURAN-Fundamental Sciences Symposium (TURAN25)

Delivered Poster Presentations

Name Surname	Section	Presentation Name	
Kübra AYSU	Physics	Electron Screening Meets Space: D+D Fusion-Driven Propulsion Systems	JUNE 23, MONDAY
Burak ÖZCAN	Physics	Constraining Key Resonances in the $40\text{Ca}(p,\gamma)41\text{Sc}$ Reaction via ANC-Assisted DWBA Analysis of Transfer Reactions and its Astrophysical Reaction Rate	
Derin ÇİVGİN	Physics	Contaminated or Clean? How Do Surface Conditions Shape Fusion Rates in Deuterated and Boron Targets?	
Doğukan BARLAS	Physics	Fusion in the Deep: Pycnonuclear Reactions Under Extreme Stellar Conditions	
Elif Seda ÇAĞATAY	Physics	Atmospheric Physics and Natural Transport Paths in the Atmosphere	
Erbil Can ARTUN	Physics	Renormalization Group Studies of Continuous-Spin Complex Systems	
Hasret Gül ÖZTOP	Physics	A Holistic Design Approach to Compact Neutron Generators: Adjustable Energies & Enhanced Yield with D-D Fusion Reactions	
Yasemin BULAT	Physics	Re-thinking the ${}^8\text{Li}(\alpha, n){}^{11}\text{B}$ Reaction: Evidence for a Missing Resonance in ${}^{12}\text{B}$	
Rana CABRAYILOVA	Physics	Aktif Matrisli Hibrit Kompozitlerin Piezoelektrik Özellikleri	
Nadia HAMMOUDA	Chemistry	Influence of Water Quality on The Electrochemical Behavior of a Carbon Steel Used in The Combustion Chamber of Boilers in a Petroleum	
Nadia HAMMOUDA	Chemistry	Inhibitive Effect of an Organic Compound on the Corrosion of Carbon Steel in Demineralized Water	
Alper OĞUZ	Chemistry	Synthesis and Photopolymerization of Imidazole-based Photoinitiator	
Elif Zeynep ÇATMA	Chemistry	Comparative study of the performance of activated carbon obtained by activation and carbonization of adsorbent material obtained from avocado	
Halide İrem KOÇ	Chemistry	Treatment of Synthetic Textile Wastewater Containing Both Basic Yellow 28 and Basic Blue 3 Dyes by Electrochemical Method	
İnci Tuğba ÖRDEKBAY	Chemistry	The Synthesis And Mesomorphic Investigation of New Chiral Calamitic Liquid Crystal Derived From Cyanobiphenyl Core	
Meltem SÖZBİR	Chemistry	One-Step Preparation of Composite Phase Change Films via Photoinitiated Oil-in-Oil Emulsion Templating	
Nazlı Özgü KIRAL	Chemistry	The Preparation and Mesomorphic Investigation of Binary Mixture of a New Chiral Rod-Like Molecule and Benzoic acid Derivative	
Rümeysa BEYAZ	Chemistry	Influence of Monomer Composition on the Morphological and Mechanical Properties of Terpene Derivative Macroporous Polymers	
Semanur SAĞLAM	Chemistry	Preparation and Characterization of Alumina Enhanced Kaolinite Supported Capric Acid Based Composite Phase Change Materials	
Eylem YAŞAR	Mathematics	Ağırlıklı Standart Olmayan Banach Uzaylarında Harmonik Fonksiyonların Yapısal Analizi	
Yeşim Aktürk DİZMAN	Biology	Analysis of Codon Usage Bias of Chloroplast Genomes in Camellia Species	

Tebriş SHAMILOV	Physics	Amorf Şerit Üretiminde Kalite Kontrol İçin Vickers Sertlik Ölçümlerinde Standart Sapma Potansiyeli
Aida AZIZOVA	Physics	Dielectric Performance and Potential Applications of HDPE/nano MgO Composites
Huseynov HUSEYNAGHA	Physics	Electrical Conductivity of Low-Dimensional Electron Gas in Asymmetric Quantum Well Scattering in Ionized Impurities in the Absence of Screening
Dilara GAFAROVA	Physics	Influence of Gamma Radiation on the Anisotropy of Electrical Conductivity in TlGaTe ₂ Crystals
Elbek KESKİNOĞLU	Physics	Role of the Pz ₁₊ State in Ferromagnetism and Topology of MnBi ₂ Te ₄
Büşra ŞAHİNAL	Physics	Yapay Zeka Destekli Moleküler Sistemlerin Kuantum Simülasyonu: H ₂ Molekülü İçin Qiskit ve OpenFermion ile VQE Uygulaması ve Hata Analizi
Şuheda Nur KOCATEPE	Chemistry	Synthesis of TX-Based Photoinitiator with Antibacterial Activity
Sanam VALIYEVA	Chemistry	Assessment Of The Impact Of Climate Change Projects On The Sustainable Development Of The Country Based On Multi-Criterion Analysis
Aynur MAMMADOVA	Chemistry	Improving Flame Resistance and Polymer Compatibility of Ethylene-Propylene-Diene Rubber (SKEPT-60) Through Polyvinyl (PVX) Modification
Ulviyye ABBASOVA	Chemistry	Synthesis Of Reagents Based On Natural Oil Acids And Ethylenediamine And Their Study As Components For Preservation Fluids
Sara HASANOVA	Chemistry	Chitosan-Tannin Based Flocculant for Heavy Metal (Ni ²⁺ , Cu ²⁺ , and Zn ²⁺) Removal from Industrial Wastewater: Investigating the Influence of pH, Reaction Time, and Stirring Speed
Arzu İBRAHİMOVA	Chemistry	Intelligent Identification of Multicomponent Mixture Components Using an Artificial Intelligence
Rufana ALIZADE	Chemistry	Investigation of Concentration Liquids Formulated by Incorporating Amidoamine and Various Fatty Acids into T-30 Oil Distillate
Sara ABBASZADE	Chemistry	Nitration Behavior of Aromatic- and Olefin-Enriched Coking Light Fraction
Sevda ASADOVA	Chemistry	Phosphoryl-Substituted α -Chloro- and α -Thiocyanatoacetaldehydes in the Reaction with α -Pyridone
Afra BÜYÜKARSLAN	Mathematics	Structural Properties of the $h_p(X)$ Class of X-Valued Harmonic Functions
Seda KIZILBUDAK ÇALIŞKAN	Mathematics	Exact Solutions of The Time Fractional Benney-Luke
Tuba SATILMIŞ	Mathematics	Lipschitz Stability Analysis of Nonlinear Set-Valued Differential Equations with Initial Time Differences
Ümit İLDİZ	Mathematics	Orlicz-Sobolev Uzaylarında Yerel Olmayan Bir Problemin Güçlü Çözülebilirliği
Parvin GULIYEVA	Mathematics	Ülke İçinde Karayolu ile Yük Taşımacılığı
Farhad MAMMADOV	Mathematics	Fe-Si-C Amorf Şerit Numunelerinin Araştırılması ve Kristal Fe-Si Muadilleri ile Karşılaştırılması
Siavash AZİMİ	Mathematics	Basicity of a Trigonometric System in Morrey-type Spaces

JUNE 24, TUESDAY

MAIN SPONSOR



SPONSORS

uyum'soft

